

| <i>Tolerance of peony crops to pre-emergence soil acting herbicides</i> |       |
|---|-------|
| Active substance  | Peony |
| clomazone   | T     |
| diflufenican  | S     |
| dimethenamid-P  | T     |
| flufenacet  | T     |
| imazamox  | MS    |
| isoxaben  | T     |
| lenacil   | T     |
| metamitron  | T     |
| metazachlor   | T     |
| metobromuron  | T     |
| metribuzin  | T     |
| napropamide   | T     |
| pendimethalin   | T     |
| prosofocarb   | T     |
| S-metolachlor   | T     |

| <i>Tolerance of peony crops to post-emergence foliar acting selective contact herbicides</i> |       |
|--|-------|
| Active substance   | Peony |
| 2,4-D  | T     |
| amidosulfuron  | MS    |
| bentazone  | T     |
| clethodim  | T     |
| clopyralid   | T     |
| cycloxydim   | T     |
| ethofumesate   | T     |
| florasulam   | MS    |
| fluroxypyr   | MS    |
| nicosulfuron   | T     |
| phenmedipham   | T     |
| rimsulfuron  | MS    |
| triflusulfuron-methyl  | MS    |

| <i>Tolerance of peony crops to post-emergence foliar acting non-selective contact herbicides</i> |       |
|--|-------|
| Active substance   | Peony |
| carfentrazone-ethyl  | S     |
| glyphosate   | S     |
| pelargonic acid  | S     |

T - Tolerant

MS - Moderately susceptible

S - Susceptible