Project title: Field Vegetables: An evaluation of

autumn/winter cauliflower, spring cabbage cultivars and other winter Brassica crops.

Project number: FV 202g

Project leader: Bill Herring, Duchy College

**Report:** Final report July 2017

**Previous report:** Annual report 2016

**Key staff:** Bill Herring

Malcolm Millar Ellis Luckhurst

**Location of project:** Trevarnon Farm, Gwithian, Cornwall.

**Industry representative:** Ellis Luckhurst

Date project commenced: 1st April 2014

Date project completed

(or expected completion date):

30<sup>TH</sup> June 2017

given in respect thereof and, to the maximum extent permitted by law the Agriculture and Horticulture Development Board accepts no liability for loss, damage or injury howsoever caused (including that caused by negligence) or suffered directly or indirectly in relation to information and opinions contained in or omitted from this document.

© Agriculture and Horticulture Development Board 2017. No part of this publication may be reproduced in any material form (including by photocopy or storage in any medium by electronic mean) or any copy or adaptation stored, published or distributed (by physical, electronic or other means) without prior permission in writing of the Agriculture and Horticulture Development Board, other than by reproduction in an unmodified form for the sole purpose of use as an information resource when the Agriculture and Horticulture Development Board or AHDB Horticulture is clearly acknowledged as the source, or in accordance with the provisions of the Copyright, Designs and Patents Act 1988. All rights reserved.

All other trademarks, logos and brand names contained in this publication are the trademarks of their respective holders. No rights are granted without the prior written permission of the relevant owners.

The results and conclusions in this report are based on an investigation conducted over a one-year period. The conditions under which the experiments were carried out and the results have been reported in detail and with accuracy. However, because of the biological nature of the work it must be borne in mind that different circumstances and conditions could produce different results. Therefore, care must be taken with interpretation of the results, especially if they are used as the basis for commercial product recommendations.

[Name] [Position] [Organisation] Team Lead / Trials Coordinator Duchy College [Name] [Position] [Organisation] Date ..... Signature ..... Report authorised by: [Name] [Position] [Organisation] [Name] [Position]

Date .....

We declare that this work was done under our supervision according to the procedures described herein and that the report represents a true and accurate record of the results

obtained.

[Organisation]

Signature .....

### **CONTENTS**

#### **Grower summary** 5 Headline 5 Background 5 Results 5 Main conclusions 8 **Science Section** 10 Introduction 10 Trial site details 15 Production details 15 Trial design 18 Trial records and data collected 18 Discussion 20 Conclusions 21 Technology transfer 21

### **GROWER SUMMARY**

### Headline

Several new cultivars introduced for 2016–17 have yielded over 15% higher than standard cultivars. Potential income can be increased by £95/ha for every additional 1% of Class 1 produce marketed.

### Background

Duchy College, working in partnership with AHDB, eight national seed houses and local growers, have carried out the evaluation of commercially available autumn, winter cauliflower and spring greens cultivars over the past 22 years. This has allowed independent comparison of new and established genetics over several seasons, giving growers' confidence when selecting new cultivars.

Initially introduced into the confidential/screening trials and then within the commercially available trials, each cultivar is recorded and evaluated year on year. With the improvement in plant breeding, new cultivars need to be assessed alongside current and established varieties in order to compare yields and quality of produce.

With improved uniformity, disease resistance, and curd protection the need to compare new introductions with current cultivars is even more important in order to plan the production of the crop to meet changing market requirements and provide the best returns for the grower.

### Results

The following cultivars are of most interest to the industry. Some are new introductions to compliment cultivars that have been grown as the standards for specific production periods.

The cauliflower trials were transplanted on the 12<sup>th</sup> and 21<sup>st</sup> July 2016, similar to previous seasons. The majority of commercial growers transplant from late June to early August.

Many of the late autumn cultivars performed well, as can be seen in the top table below.

Results over several seasons will need to be considered when selecting cultivars for production. Information on all cultivars can be found in the 'Full Trial Report'.

Evaluations of curd protection, leaf colour, disease resistance/susceptibility, uniformity, curd colour and density, suitability for various markets, defects and ease of harvesting are found in the Full Trial report in the Appendices.

The spring greens trial was planted into good conditions on the 12<sup>th</sup> September 2016 and produced high quality greens that were harvested in early March 2017. Results are found in the Full Trial report in the Appendices.

### Autumn Cauliflower 2016 Top performing cultivars

Transplanted at 25,315 plants/hectare (10,249/acre) on both the 12<sup>th</sup> and 21<sup>st</sup> July 2016

The majority of cultivars produced curds 11cms in diameter (8 per tray) however a few cultivars produced larger curds (6 per tray) from a very high percentage of Class 1 produce.

Please refer to the 'Full Trial Report' for further details.

Seed House	Cultivar	% Class 1	Trays/Hectare	(Acre)	Heading period
			Class 1		10 – 90% dates
			(8/tray )	(6/tray)	
Sakata	Jericho	92	2090 (846)	1091 (442)	17.10.16 – 23.10.16
Sakata	Amandine	94	2260 (915)	968 (392)	21.10.16 - 30.10.16
Syngenta	C4040	95	1378 (558)	2158 (874)	23.10.16 – 29.10.16
Monsanto	SV 5965	87	2374 (961)	506 (205)	25.10.16 - 08.11.16
Clause	Optimist	90	2309 (935)	716 (290)	30.10.16 – 10.11.16
Rijk Zwaan	26 - 407	80	2349 (951)	225 (91)	30.10.16 – 12.11.16
Monsanto	SV 5982	90	2793 (1131)	74 (31)	06.11.16 – 22.11.16

### Winter Cauliflower 2016 – 17 Top performing cultivars.

Winter cauliflower is typically harvested from early November to April. Most commercial cultivars grown produce a minimum of 70% marketable crop with some cultivars producing 70% + Class 1 cauliflower at the beginning and end of this harvest period, in a typical season.

Yields will vary as there were two transplanting dates (12th and 21st July 2016). Plant spacings were the same for each cultivar within the trials. Summary of both transplanting dates used in table below.

Some cultivars also produced larger curds that are included on the table below.

Transplanted at 21,193 plants/hectare (8580/acre)

Please refer to the 'Full Trial Report' for further details.

Seed House	Cultivar	%	Trays/Hecta	re (acre)	Heading period
Seed House	Cultival	Class	Class 1	ie (acie)	Heading period
		1	(8/tray)	(6/tray)	10 - 90% dates
Enza	Anique	95	1531 (620)	1319 (534)	27.10.16 - 01.11.16
Clause	Navalo	91	1193 (483)	1625 (658)	27.10.16 - 07.11.16
Clause	Diwan	85	1996 (808)	333 (135)	08.11.16 - 20.11.16
Enza	EW40622	96	1699 (688)	1131 (458)	08.11.16 - 26.11.16
Clause	Galiote	87	2225 (901)		17.11.16 - 10.12.16
Syngenta	Leocen	93	2243 (908)	140 (57)	28.11.16 - 17.12.16
Hazera	Lecatis	86	2146 (869)	177 (72)	10.12.16 - 21.12.16
Clause	Triomphant	87	2237 (906)	59 (24)	15.12.16 - 31.12.16
Syngenta	C5049	87	2319 (939)		21.12.16 - 11.01.17
Clause	Jerboam	82	2176 (881)		31.12.16 - 12.01.17
Elsoms	Paciano	91	2455 (994)	106 (43)	07.01.17 - 17.01.17
Elsoms	Baterno	83	2208 (894)		15.01.17 - 28.01.17
Syngenta	5064	87	2119 (858)	247 (100)	25.01.17 - 01.02.17
Hazera	Trevaskis	91	2450 (922)	178 (72)	03.02.17 - 08.02.17
Monsanto	5697	87	2311 (936)		03.02.17 - 10.02.17
Clause	Fleet	87	2285 (925)		05.02.17 - 12.02.17
Clause	Redoutable	82	2174 (880)		05.02.17 - 14.02.17
Syngenta	Britten	85	2198 (890)	59 (24)	06.02.17 - 17.02.17
Syngenta	C5052	87	2274 (921)	30 (12)	11.02.17 - 21.02.17
Monsanto	Carantic	91	2411 (976)		17.02.17 - 23.02.17
Clause	CLX 33611	92	2331 (944)	140 (57)	20.02.17 - 23.02.17
Clause	Mascaret	95	2514 (1018)	·	20.02.17 - 27.02.17

Elsoms	Madiot	92	2440 (988)		22.02.17 - 03.03.17
Syngenta	C5067	87	2453 (993)	40 (16)	24.02.17 - 04.03.17
Elsoms	Mystique	91	2361 (956)	59 (24)	25.02.17 - 05.03.17
Elsoms	Tempest	88	2334 (945)		04.03.17 - 15.03.17
Hazera	AE 9307	93	2465 (998)		06.03.17 - 13.03.17
Clause	CLX 33518	91	2453 (933)		15.03.17 - 25.03.17

### Late heading Winter Cauliflower 2016-17 Top performing varieties

These plots were transplanted on the 12th July 2016 aiming to head in the period of April through to late May 2017. Over recent seasons there has been a shortage of cauliflower for this period, which has resulted in the increase of imported produce to maintain supply to the various market outlets. Although fixed price returns to the grower over this heading period are not guaranteed there is scope to aim for this market share.

The very good growing conditions throughout the autumn and winter months did not have any impact on the later heading cauliflower in regards initiation of the curds and this resulted in the majority of cultivars being harvested up to 14 - 21 days earlier than their expected harvesting slots. Yields, as for the winter heading cultivars, were at least 15% above average.

Transplanted at 21,193 plants/hectare (8,580/acre)

Some cultivars also produced larger curds which are included in the table below.

Please refer to the 'Full Trial Report' for further details.

Seed House	Cultivar	% Class 1	Trays/Hecta	are (Acre)	Heading period
110000		Oldoo I	(8/tray)	(6/tray)	10 - 90% dates
Syngenta	C5044	93	583 (236)	2507 (1015)	06.03.17 - 13.03.17
Syngenta	C5033	93	1113 (451)	1591 (644)	14.03.17 - 16.03.17
Monsanto	Tenfold	91	1474 (597)	1360 (551)	14.03.17 - 17.03.17
Elsoms	Jerome	89	2358 (955)		14.03.17 - 23.03.17
Hazera	VT 4013	78	2067 (837)		15.03.17 - 23.03.17
Elsoms	Vogue	90	2287 (926)	106 (43)	23.03.17 - 29.03.17
Seminis	SV0890AC	83	2003 (811)	274 (111)	25.03.17 - 03.04.17
Clause	CLX 33116	92	2327 (942)	143 (58)	27.03.17 - 11.04.17
Syngenta	Delon	93	2339 (947)	185 (75)	03.04.17 - 16.04.17
Clause	CLX33612	89	2336 (946)		05.04.17 - 15.04.17
Elsoms	Fletcher	94	2504 (1014)		10.04.17 - 19.04.17

The trials were undertaken at Trevarnon Farm, Gwithian, Cornwall. The soil type is a sandy clay loam. The site is south facing and is part of a farm rotation based around Brassicas, cereals and grass break crops.

The farm has traditionally grown Brassica crops supplying both the multiple and local markets.

The trial itself is treated as a commercial crop undergoing similar field operations as the commercially grown crops. The cultivars are harvested twice a week on a similar basis as commercial crops on the farm.

### Spring Green cultivars 2016 – 17 Main Results

Seed house Cultivar

Monsanto Winter Supreme

A L Tozer Winter Green

Nickerson Kempsy

Monsanto Winter Special

Seed House Cultivar	Comments and observations on 19.01.2016	Transplanting Date	Harvest dates	Average weight of marketable green
Monsanto Winter Supreme	Very leafy. Slightly leggy frame. Paler inner leaf. Dark green outer leaves. Very Uniform. Excellent greens. 93% of transplanted crop marketable.	12.09.16	08.03.2017	197 grams
A. L. Tozer Wintergreen (Traditional open pollinated variety)	Slightly Crinkled leaf type. Variable size plants/greens. Dark green leaf. Medium height. 90% of transplanted crop marketable.	12.09.16	08.03.2017	230 grams
Nickerson Kempsy	Very Uniform. Medium height. Close internodes on stem. Slightly crinkled leaf. Mid green in colour, slightly paler inner leaves. Good greens. 90% of transplanted crop marketable.	12.09.16	08.03.2017	241 grams
Monsanto Winter Special	Some variation in green sizes. Dark green leaf. Heavy cabbage greens. Easy to cut. Not as tall as Winter Supreme. 87% of transplanted crop marketable.	12.09.16	08.03.2017	305 grams

### Main conclusions

All of the cultivars that have performed well during 2016 – 17 are commercially available to growers, each with their own characteristics, traits and qualities. The large number of cultivars available enables growers to select suitable cultivars for their own production systems, which vary from farm to farm and area to area. The cultivars available ensure continuous supply, complimenting each other throughout the production period. However, there is a continuous need to improve output and quality especially in the production period between late December and early February, when yields can be lower. There have been a number of new introductions over this period which have performed very well alongside established cultivars.

Trials have also been undertaken where a number of confidential cultivars from several seed companies have been assessed alongside new named cultivars being introduced into the commercially available cultivar trials for 2017 -18, with more to follow in subsequent years. There is renewed interest in the late heading cultivars for the heading period of April and into mid-May.

### **SCIENCE SECTION**

#### Introduction

Working in partnership with AHDB and a number of national seed-houses, these trials have evaluated the commercially available Autumn, Winter Cauliflower, Late Winter Cauliflower and Spring Green cultivars, so that improved cultivars can be assessed alongside current and established cultivars. The aim is to improve yields, quality of produce and continuity of supply.

### **Autumn, Winter and late Winter Cauliflower**

The Autumn Cauliflower trial compared established cultivars alongside new introductions for the heading period of early October to early November. Several new cultivars have been introduced this year.

The main Winter Cauliflower trial compares cultivars transplanted in early and late July and provides detailed information on the advantages/disadvantages of staggered transplanting dates, which is commercially practiced in order to maximize the potential of some cultivars. These cultivars mature (head) from November through to April.

Increased interest and opportunities to market later heading winter cauliflower has resulted in a number of new introductions for the May - June period, with several cultivars performing well.

The cauliflower trials provide detailed information on each cultivar including:

- Harvesting period 10%, 50% and 90% harvest dates.
- Percentage Grade 1, Grade 2 and unmarketable produce.
- Marketable yields.
- Comments on any defects, including disease tolerance especially for Xanthomonas and mildew, pest presence, and the suitability for the various market outlets.
- Evaluation on the ease of harvesting, packing, uniformity, depth and quality of curds.
- Comments on curd protection, leaf quality and crop growth.

### Varieties and numbered selections included

The Late Variety trial consisted of 12 cultivars replicated twice, transplanted at 21,193 plants/hectare (8,580/acre), transplanted on the 12th July 2016.

The Autumn Cauliflower trial consisted of 9 cultivars replicated twice, transplanted at 25,315 plants/hectare (10,249/acre), transplanted on the 12th July 2016. This was an increase in density from 21,193 plants/hectare (8,580/acre) in 2014-15 to reflect current commercial practice locally.

The Winter Cauliflower trial consisted of 36 cultivars transplanted at 21,193 plants/hectare (8,580/acre), transplanted on the 12th and 21st July 2016.

### **Spring Green Trial**

The Spring Greens Trial compares commercially grown hybrid cultivars transplanted alongside the traditional open pollinated Wintergreen cultivar. These are then assessed for suitability as greens being harvested December to March depending on the cultivar and season.

The field plan is found in the Appendices.

### Late Winter Cauliflower Cultivars 2016-17

### Transplanted 12th July 2016

<u>Plot</u>	Seed House	<u>Variety</u>	Heading Period
1	Elsoms	Cartagena	Mid May
2	Hazera	VT4013	Early/Mid May
3	Elsoms	Fletcher	Early May
4	Clause	CLX33612 (CHF12-27	
5	Elsoms	Cartagena	Mid May
6	Hazera	VT4013	Early/Mid May
7	Elsoms	Fletcher	Early May
8	Clause	CLX33612 (CHF12-27	
9	Elsoms	Douglas	Late April
10	Monsanto	Tenfold	Late April
11	Syngenta	Delon	Mid/Late April
12	Elsoms	Jerome	Mid April
13	Elsoms	Douglas	Late April
14	Monsanto	Tenfold	Late April
15	Syngenta	Delon	Mid/Late April
16	Elsoms	Jerome	Mid April
16B	Seminis	SV0890AC	Mid April
17	Syngenta	C5033	Early April
17B	Seminis	SV0890AC	Mid April
18	Elsoms	Vogue	Early April
19	Syngenta	C5044	End March
20	Clause	CLX 33116 (Avelmore)	Mid April
21	Syngenta	C5033	Early April
22	Elsoms	Vogue	Early April
23	Clause	CLX 33116 (Avelmore)	-
24	Syngenta	C5044	End March

### Autumn Cauliflower cultivars

First Planting Transplanted 12<sup>th</sup> July 2016

<u>Plot</u>	Seed House	<u>Variety</u>	Heading Period
81 82 83 84 85	Sakata Elsoms Monsanto Elsoms Sakata	Jericho Benidorm SV5965AC Cantabria Amandine	Mid October Mid October Late October Mid/Late October Late October
86 87 88 89	Rijk Zwaan Clause Monsanto Syngenta		November Early November Early November Early November

2016-17

Autumn Cauliflower cultivars 2016-17

Second Planting Transplanted 21st July 2016

Seed House	<u>Variety</u>	<b>Heading Period</b>
Sakata	Jericho	Mid October
Elsoms	Benidorm	Mid October
Monsanto	SV5965AC	Late October
Elsoms	Cantabria	Mid/Late October
Sakata	Amandine	Late October
Rijk Zwaan	26-407 Early	November
Clause	Optimist	Early November
Monsanto	SV5982AC	Early November
Syngenta	C4040	Early November
	Sakata Elsoms Monsanto Elsoms Sakata Rijk Zwaan Clause Monsanto	Sakata Jericho Elsoms Benidorm Monsanto SV5965AC Elsoms Cantabria Sakata Amandine Rijk Zwaan 26-407 Early Clause Optimist Monsanto SV5982AC

### Winter Cauliflower cultivars

## Time of planting Trial First planting 12<sup>th</sup> July 2016

<u>Plot</u>	Seed House	<u>Variety</u>	Head	ling Period
90	Elsoms	Telde		November
91	Enza	ANIQUE (20259)	6)	November
92	Clause	Navalo	,	Mid November
93	Enza	EW40622 (2025	99)	Mid November
94	Clause	Diwan `	,	Late November
95	Elsoms	Belot		Late November
96	Enza	AGILITY (20259)	7)	Late November
97	Clause	Galiote	,	Early December
98	Clause	Triomphant	Mid [	December
99	Syngenta	C5049		Mid December
100	Hazera	Lecatis		Mid December
101	Syngenta	Leocen		Late December
102	Clause	CLX 33422 (Jerb	oam)	Late December
103	Elsoms	Paciano		Early January
104	Elsoms	Baterno (BJ2589	9)	January
105	Clause	Brick		Late January
106	Syngenta	C5064		Late January
107	Hazera	Trevaskis		Late Jan/February
108	Clause	CLX 33610		Early Feb
109	Clause	Redoutable	Early	<sup>,</sup> February
110	Monsanto	5697		February
111	Syngenta	Bretten		February
112	Clause	Fleet		Mid February
113	Clause	CLX33611		Late February
114	Syngenta	C5067		Late February
115	Clause	Mascaret		Late February
116	Syngenta	C5052		Late February
117	Elsoms	Gerona (BJ 2860	))	Late February
118	Syngenta	Caylen		Feb/Early March
119	Monsanto	Carantic		Feb/Early March
120	Clause	CLX 33518 (CHF	- 12-2	73) Early March
121	Elsoms	Madiot		March
122	Elsoms	Mystique		Mid/End March
123	Elsoms	Isadora		Mid/End March
124	Hazera	AE 9307 (Parotis	s)	Late March
125	Elsoms	Tempest		Late March

### Winter Cauliflower cultivars 2016-17

Time of planting Trial Second planting 21st July 2016

<u>Plot</u>	Seed House	<u>Variety</u>	Head	ding Period
140 141 142 143 144 145 146 147 148 149 150	Elsoms Enza Clause Enza Clause Elsoms Enza Clause Elsoms Enza Clause Clause Syngenta Hazera	Telde ANIQUE (20259 Navalo EW40622 (2025) Diwan Belot AGILITY (20259) Galiote Triomphant C5049 Lecatis	Nove 6) 99) 7)	<del>_</del>
151 152 153 154	Syngenta	Leocen CLX 33422 (Jerl Paciano Baterno (BJ2589 Brick C5064 Trevaskis CLX 33610 Redoutable 5697 Bretten Fleet	9)	Late December
162 163 164 165 166 167 168 169 170 171 172 173 174 175	Clause Syngenta Clause Syngenta Elsoms Syngenta Monsanto Clause Elsoms Elsoms Elsoms Elsoms Elsoms Elsoms Hazera Elsoms	CLX33611 C5067 Mascaret C5052 Gerona (BJ 286) Caylen Carantic CLX 33518 (CH Madiot Mystique Isadora AE 9307 (Parotic Tempest	Late Late 1	Late February Late February Late February Late February Late February Feb/Early March Feb/Early March

### SPRING CABBAGE TRIALS 2016 -17

4 rows / variety 25cm spacing in row

Seed house Cultivar

Monsanto Winter Supreme .

A L Tozer Winter Green

Nickerson Kempsy

Monsanto Winter Special

Transplanted 12th September 2016 into moist soil conditions. 1000 of each cultivar. Module grown transplants.

Winter Green is an open pollinated cultivar, with the remainder being F1 hybrids.

### Trial site details

The trials were conducted at Trevarnon Farm, Gwithian, Cornwall, by kind permission of Mr. J. Hosking and Son.

### **Production details**

The trials followed a crop of Winter Greens 2015 – 16 and left fallow prior to field operations in early September 2016.

### CULTURAL INFORMATION 2016 – 17

### AUTUMN and WINTER CAULIFLOWER TRIALS

7 to 1 0 1 1 1 1 t a 1 1	<u> </u>
<u>Fertiliser</u>	
24.06.16	Base Dressing KESERITE 25% Magnesium @ 247Kg/ha (2cwt/acre) NITRAM 34.5% Nitrogen @ 300Kg/ha (2.42 cwt/acre)
17.08.16	PERLKA 271Kg/ha (2.2 cwt/acre) (Applied inter row through shoots)
03.11.16	Top Dressing Yara CALCIUM NITRATE 15.5% Nitrogen 247Kg/ha (2.0 cwt/acre)
20.12.16	140 Kg/ha or 1.13 CWT / AC CALCIUM NITRATE 15.5 % N
08.01.17	90 Kg/ha or 0.72 CWT / AC CALCIUM NITRATE 15.5 % N
Weed contro	<u>bl</u>
16.07.16	GAMMIT @ 0.15Ltr/ha SULTAN @0.6Ltrs/ha (First planting). Inter row cultivations.
29.07.16	GAMMIT @ 0.15Ltr/ha SULTAN @0.6Ltrs/ha MOLYTRAC @ 0.250Ltrs/ha in 300 Litres of water. (Second planting). Inter row cultivations.
Pest control	Wire netting prior to transplanting as rabbit control

### Chemical plan

- BISCAYA @ 0.4Ltr/ha ACTIVATOR 90 @ 0.3Ltr/ha 14.08.16 MOLYTRAC @ 0.250Ltr/ha in 300 Litres of water.
- 10.09.16 **RUDIS** @ 0.2Ltr/ha, DIFCOR @ 0.2Ltr/ha. HALLMARK ZEON@0.05Ltr/ha, BREAKTHRU@ 0.3Ltr/ha,FARM FOS @ BISCAYA @ 0.4Ltr/ha 2Ltr/ha, MAGFLO@1.5Ltr/ha + TDS MAJOR (Slug Pellets)@4Kg/ha. All Trials 1st FUNGICIDE
- White Blister control. FOLIO GOLD@0.75Ltr/ha, SIGNUM 30.09.16 @O.75Kg/ha. ACTIVATOR 90 @ 0.3Ltr/ha in 300 Litres of water. All Trials.
- 07.11.16 RUDIS@ 0.2Ltrs/ha, PLOVER@0.2Ltrs/ha, ACTIVATOR 90 @ 0.3Ltr/ha, MAGFLO@3.0Ltr/ha in 300 Litres of water on selected plots due to heading. (3<sup>rd</sup> FUNGICIDE)
- 16.12.16 In 400 Litres water/ha all trial 4th FUNGICIDE (SELECTED PLOTS

# ONLY due to heading) RUDIS @ 0.4 Ltr/ha, ACTIVATOR 90 @ 0.3Ltr/ha, MAGFLO @ 2 Ltr/ha, COPTREL @ 0.5 Ltr/ha, MOLYTRAC @ 0.25Ltr/ha, FARMFOS @ 2 Ltr/ha

### <u>CULTURAL INFORMATION 2016 – 17</u>

Duchy Trial	Spring Greens	Planted 12/09/16
Fertiliser		
01.09.16	Base Dressing	6-18-36@494Kg/ha (4.0cwt/acre)
02.11.16	Top Dressing	CALCIUM NITRATE 15.5%@ 200Kg/ha (1.6cwt/acre)
06.01.17	Top Dressing	CALCIUM NITRATE 15.5%@ 222Kg/ha (1.79cwt/acre)
Weed control	_	
15.09.16		GAMMIT @ 0.15Ltr/ha SULTAN @1.5Ltrs/ha in 300 Litres of water. SLUXX@4Kg/ha (Slug Pellets)
Chemical Pla	an	
09.10.16		DECIS@0.3Ltr/ha, ACTIVATOR 90 @ 0.3Ltr/ha, MAGFLO@2.5Ltr/ha, MOLYTRAC @ 0.250Ltrs/ha in 300 Litres of water.
01.11.16		BISCAYA @ 0.4Ltr/ha, ACTIVATOR 90 @ 0.3Ltr/ha DIFCOR @ .3Ltr/ha,MAGFLO@3.0Ltr/ha. MOLYTRAC @ 0.250Ltrs/ha in 300 Litres of water.
02.12.16		PLOVER @ 0.3 Ltr/ha, ACTIVATOR 90 @ 0.3Ltr/ha, MAGFLO @ 2 Ltr/ Ha, COPTREL @ 0.5 Ltr/ha FARMFOS @ 3 Ltr/ha in 300 Litres of water.

### Trial design

Each plot consisted of 4 rows with the outer rows being guards. Fifty plants of each cultivar were recorded with harvesting occurring every Monday and Thursday throughout the harvesting period of the trials (October – July 2016-17).

### Trial records and data collected

For each autumn and winter cauliflower cultivar the following record sheet is completed at each harvest and on completion of harvesting summaries are made of all data collected supported by visual observations as appropriate. These to be found in the Full Trial report in the Appendices.

### Cauliflower record

	Title of Exper		nt					planting		1	Ref. V	FC	)3							
	LAPOI	,				,			Year	16	17								···	
	Plot								i eai	10	17				P	lot				
		C	Col.	(6-8	3)		·						·				(6-	·8)		
	ple		Defe	ects		=	+	t	Missi	ORDERS Fing Plots	- When w	hole p	olot		ple		Def	ects	+ +	l to
	Class or Unmarketable	Size	Ricey	Bracted	Loose	Curd Colour	Curd Depth + Immature	Space A not punched	missi * und	ing put OO ler class. ing Plan	under hai	rvest a	and	Harvest	Class or Unmarketable	Size	Ricey	Bracted	Curd Colour	Space A not
<b>-</b>		0)		ш			0 =	υ <u>α</u>	missi throu		nt unre arietal fac	ecordal ctors p	ble			0)	<u> </u>	ш _		
									date/o curd i Class O – E 2 – C	code again recorded of s or Unma Extra Class Class 2, F – inmarketab	nst 1 <sup>st</sup> , 11 on one harv o <mark>rketable</mark> o, 1 – Class - Facepack	I <sup>th</sup> + 2 vest. s 1,	est 21 <sup>st</sup>							
										- O Not sized 1 Button 8 Facepack										
										12 16 8 Blind 9 Frost kille	d in vegetati	ve stage	e							
									appro	y or Loc opriate colu ts	<u>ose</u> – P umn or leav	ve blar	nk. =							
									2 Gr	hite fine br reen bracts <u>I <b>Colour</b></u>		blank.								
										1 Yellov 2 Pink 3 Disco	w loured d curd									
										5 Slight soake	frosted ed. re frosted		-							
										X Leaf s	Scorch	t write								
									or lea Curd Reco	ave blank. I <b>Depth + I</b> ord curd d	mmature lepth on	all siz	ed _							
									curds 1 1	Depth ½ s <sub> </sub> Depth ⅓ s <sub> </sub>	phere	snhere	<b>.</b>							
									3   4   	Depth <sup>1</sup> / <sub>3</sub> s Immature leave blant	phere at final hak k.	arvest	or							
									inforn data	ce A may mation held (e.g. bird an addition)	pful in int <i>damage,</i>	terpreti stem	ing <i>rot</i>							

### Discussion

The weather pattern from transplanting in July through to the end of December played a significant part in the production of cauliflower planned for harvest from late October through to March.

Following favorable planting conditions in July 2016 there was excellent crop establishment in August despite some late cabbage root-fly being present in some crops. With good growing conditions in September, October and November, and above average day and night temperatures, this resulted in some early winter cauliflower cultivars being harvested up to 10 days later than would be normally expected, due to later vernalisation of the curds. By mid December cultivars were back on track in regards harvesting periods with the season finishing earlier than normal for growers concentrating on the December – early April harvest window.

The introduction of Verimark as a new control of cabbage root-fly, along with a broad spectrum activity on a range of other pests clearly impacted on crop establishment. By preventing feeding of early season pests this resulted in strong robust crops going into the winter period which contributed to a very productive season.

The season was exceptional in the fact that nearly all cultivars grown for the harvest period of early November to early June produced yields of Grade 1 produce well over 10% above seasonal averages and in some cases over 20% higher yields. However this did not result in a glut period, rather a much larger quantity of high quality cauliflower available through the harvest period, which found a ready market due to the quality of the product.

Within the trials and in commercial production, the average marketable yields across the harvest period were the highest recorded over the 25 years.

The introduction of some new autumn and winter cauliflower cultivars in 2014 -15 have replaced some of the mainstay cultivars grown over recent years and these again have performed consistently well, as seen in 2015-16. Some new introductions for 2016 – 17 have harvested over 15% above standard cultivars based on 25,315 plants/hectare (10,249/acre) for the autumn cultivars and 21,193 plants/hectare (8,580/acre) for the winter cultivars. Potential income can be increased by £95/ha for every additional 1% Class 1 marketed, which is a real possibility with many producers now on a fixed price for their produce.

The top performing cauliflower cultivars highlighted within the results section are cultivars that have performed above the average yield of 75% Grade 1.

Many of the autumn heading cultivars produced well above 70% Class 1 with some cutting out over 85%. Although the price to the grower is much less than the winter cauliflower price, a greater planting density per hectare would increase yields and returns even at a lower unit price to the grower.

### Conclusions

Within the autumn and winter cauliflower cultivars, several seed houses have introduced numbered cultivars, which have performed well alongside established and recently introduced cultivars. As with all new introductions, these need to be evaluated over several seasons to ensure consistent performance under different weather conditions. Those that have done well this season include the following:-

Number	Heading period
C4040	Late October
SV5965	Early November
SV5982	Mid November
EW40622	Mid November
C5049	Late December
5064	Late January
5697	Early February
C5052	Mid February
C5067	Late February
AE9307	Early March
C5033	Mid March
VT4013	Late March
	C4040 SV5965 SV5982 EW40622 C5049 5064 5697 C5052 C5067 AE9307 C5033

Each cultivar performs differently from season to season, however data is available over the past 20 seasons with each entry tested for several seasons. This provides a basis upon which growers can make an informed decision on which cultivars to grow in order to provide a continuous supply.

### **Technology transfer**

The trials are harvested twice a week and results are updated weekly at <a href="http://www2.cornwall.ac.uk/research/herring/default.asp">http://www2.cornwall.ac.uk/research/herring/default.asp</a>

Open days were held in December 2016, January and March 2017, where the trials were available to be viewed and up to date information was made available to growers and industry representatives. A growers evening also took place on 18<sup>th</sup> January meeting to update growers on current Brassica developments <a href="https://horticulture.ahdb.org.uk/event/winter-cauliflower-spring-cabbage-20162017-trials-open-day">https://horticulture.ahdb.org.uk/event/winter-cauliflower-spring-cabbage-20162017-trials-open-day</a>

٠

### **Appendices**

The following appendices summarise results of all cultivars in the:

Trials field Plan

Autumn Cauliflower Variety Trials 2016

Winter Cauliflower Variety Trials for 2016–17

Late Winter Cauliflower Trials 2017

With additional comments regarding each cultivar.

		BRAS	SICA	TRIA	LS FIE	ELD P	LAN	2016	-17		Spacing	g 34" x 2	1"	CAU	LIFLO	WER		
	SECONI	D PLANT	ING						FIRST F	LANTIN	G							
Winter	Winter	Winter	Winter	Winter	Autumn	Winter	Winter	Winter	Winter	Winter	Autumn	Autumn Winter	Winter		Late va	rieties	Winter	
4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows	4 rows		
	9 beds	9 beds	9 hads	9 beds	9 beds		9 beds	9 beds	9 hads	9 beds	9 beds	9 beds	Confider 8 heds		8 beds	8 hads		
Guards	170	161	152	143	134		125	116	107	98	89	80	35	o beas	o beas	o beas		
Guards	169	160	151	142	133		124	115	106	97	88	79	single	24	16	8		
Guards	168	159	150	141	132	G	123	114	105	96	87	78	lines	23	15	7	G	(
Guards	167	158	149	140	131	U	122	113	104	95	86	77		22	14	6	U	I
175	166	157	148	139	130	Α	121	112	103	94	85	76		21	13	5	Α	
174	165	156	147	138	129	R	120	111	102	93	84	16 single		20	12	4	R	
173	164	155	146	137	128	D	119	110	101	92	83	lines		19	11	3	D	
172	163	154	145	136	127		118	109	100	91	82	Plots	Plots	18	10	2		
171	162	153	144	135	126		117	108	99	90	81	60 - 75	25 - 59	17	9	1		
													L	AY B	Y			

		AUTU	MN CA	ULIFL	OWER	VARI	ETY T	RIAL				
		2016						SPACI	NG 34"	X 18"	10249/	Acre
A= Transpla	anted 12th July	B= Tran	splanted	21st July								
										Crates / Acre	<u>;</u>	
VARIETY	PLOT	CUT	TING PER	IOD	DAYS	Class	Class	Unmarket	Facepack			Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Sakata	81A	13.10	17.10	20.10	7	90	2	8	641	683	13	0
Jeri <u>c</u> ho _	126B	_ 22. <u>1</u> 0	_22.10 _	27.10	5	94	2	4	1050	201	13	0
A= Transpla	anted 12th July Average	B <sub>17</sub> , T <sub>0</sub> an	spkyyted.	21st July	6	92	2	6	846	442	13	0
Smooth curds	Slightly wide base	e to c <mark>urds. Paler i</mark>	nner leaf. I	niform. Ver	y good Face	ack materi	ıl. High %	large curds.	Good curd o	Crates / Acre	,	
VARIETY	PLOT		TTING PER		DAYS	Class	Class	Unmarket	Facepack			Class 2
Flooms	82A	10% 31.10	50% 31.10	90% 5.11	5	18	<del>2</del> 6	%	5 <b>8</b> 9	<u>54</u>	16 282	12
Sakata	85A 127B	13.10 5.12	17.10 8.12	20.10	7	90	9	10	<u>826</u>	717	0 26	8
Amandine	Average	3 <del>9</del> .19	19.41	29:11	61	88	25	125	<del>173</del> 84	68	194	8
- war 1 0 11 days 1 5 1	Average	21.10	27.10	30.10	9	94	0	6	915	392	0	0
Very uniform.	Good weight. Slig	htly wide base to c	urds. Very	good Facepa	ck material				oks very go	od in trays.		
D'1 7	0.64											
Rijk Zwaan	83A 131B	22.10 20.10	24.10 24.10 14.11	31.10 31.10	9 ! <u>1</u>	68 90 91	16 2	16	820 1025 1082	68 171 114	102 13	0
26-407	. 100=	7.11		24.11	17/		0	9			0	0
	Average Average	30.10 25.10	3.11	12.11 8.11	13	80	8	12	951 961	91 205	51 32	8
<del>Upright frame</del> Uniform, Tali	Average . Slightly wide bas frame. Slightly wide	<del>e to curds. Paler t</del> de base to curds, n	<del>ight wrapp</del> ot easy to n	<del>er leaf. Medi</del> ack. Mediun	<del>um depth cu</del> n depth. Twi	rds. Slightl isted leaf ove	<del>y crinkled l</del> r curds. Ve	<del>eaf type. Lool</del> erv good Face	ks very good pack mater	<del>l in trays.</del> iai. Verv fev	defects.	
Very Good Fac	frame. Slightly wie epack material es	<del>pecially from the</del>	<del>later planti</del>	ng. Some sm	all white br	acts.		7 8	<b>F</b>	J		
ורז	844	17.10	20.10	31.10	1 <u>4</u>	55	31	14	703	0	201	0
Clause	84A 87A 129B	22:18 10:11	20.10 24.10 14.11	27.10	5	55 100 76	11	0	845	581 114	0	<u> </u>
Optimist	129B 132B	7.11	14.11	24.11 24.11	14 17	76 80	0	13 20	898 1024	0	28	0
<b>X</b>	Average	30.10	3.11	10.11	11	90	0	10	935	290	0	0
	Some small white Easy to see, cut a		ev franspla Very good	rting Very	peavy chics	Shisphy M	de base to c	HERS GOOD CI	ird cover. G	ood Facepac	k material.	
Smooth curus.	Easy to see, cut al	iu pack.										
Monsanto	88A	27.10	31.10	7.11	11	92	0	8	1179	0	0	0
SV5982	133B	17.11	24.11	8.12	21	88	2	10	1082	59	11	0
	Average	6.11	12.11	22.11	16	90	1	9	1131	30	5	0
Very tall fram	e. Strong midrib to	o leaf. Upright with	h spreading	g outer leave	s. Very unif	orm smooth	curds. Lool	ks very good	in trays. Go	od Facepack	•	
Syngenta	89A	13.10	13.10	17.10	4	94	0	6	688	696	0	0
C4040	134B	3.11	7.11	10.11	7	95	2	3	427	1053	11	0
	Average	23.10	25.10	29.10	6	95	1	4	558	874	6	0
Dark green le		curd cover. Slight		e to curds. V	ery good Fa	cepack amte	rial. Very f	ew defects. N	ledium dept	h. Paler inne	er leaf.	

COMP	ARING	LATE A	UTUM	N CAU	LIFLOV	ARIETIES	10249plar	nts/acre	(8785plan	ts/acre in	2010-11)
SEEDHOU	ISE	CLAUSE				SEEDHO	JSE	MONSAN	TO/SEMIN	IS	
VARIETY		<b>OPTIMIST</b>				VARIETY		5982			
	Planting	50%	Days	%	Trays 8		Planting	50%	Days	%	Trays 8
	Date	cut date	-	Class 1	per Acre		Date	cut date		Class 1	per Acre
2016 - 17	21st July	14th Nov	116	80	1024	2016 - 17	21st July	24th Nov	126	88	1126
2016 - 17	12th July	24th Oct	104	100	1280	2016 - 17	12th July	31st Oct	111	92	1179
2015 - 16	9th July	19th Oct	102	86	974	2015 - 16	9th July	29th Oct	112	97	1042
2014 - 15	10th July	4th Oct	86	69	764	2014 - 15	10th July	14th Oct	96	98	1157
2013 - 14	10th July	22nd Oct	104	72	871	2013 - 14	10th July	4th Nov	117	60	737
2012 - 13	19th July	11th Nov	115	88	1128	2012 - 13	19th July	17th Nov	121	88	1123
2011 - 12	13th July	17th Oct	96	69	877	2011 - 12	13th July	27th Oct	106	86	1076
2010 - 11	10th July	26th Oct	108	72	891	2010 - 11	10th July	3rd Nov	116	84	922
2009 - 10	10th July	2nd Nov	115	81	1067	2009 - 10	10th July	18th Nov	131	81	1076
						SEEDHO	JSE	SAKATA			
						VARIETY		<b>JERICHO</b>			
							Planting	50%	Days	%	Trays 8
							Date	cut date		Class 1	per Acre
						2016 - 17	21st July	22nd Oct	93	94	1200
						2016 - 17	12th July	17th Oct	97	90	1153
SEEDHOU	ISE	SAKATA				2015 - 16	9th July	13th Oct	96	83	1062
VARIETY		AMADINE									
	Planting	50%	Days	%	Trays 8	SEEDHO	JSE	MONSAN	TO/SEMIN	IS	
	Date	cut date	•	Class 1	per Acre	VARIETY			5965	5	
2016 - 17	21st July	7th Nov	109	98	1204		Planting	50%	Days	%	Trays 8
2016 - 17	12th July	17th Oct	97	90	1163		Date	cut date		Class 1	per Acre
2015 - 16	9th July	13th Oct	96	84	1075	2016 - 17	21st July	14th Nov	116	84	1076
2014 - 15	10th July	2nd Oct	84	82	880	2016 - 17	12th July	24th Oct	104	90	1153
2013 - 14	10th July	29th Oct	111	79	944	2015 - 16	Not in Tria				
						2014 - 15	10th July	24th Oct	106	74	886
						2012 - 13	19th July	10th Nov	114	90	1152
						2011 - 12	13th July	17th Oct	96	90	1152
						2010 - 11	10th July	28th Oct	110	78	856
						2000 40	10th July	30th Oct	112	100	1280

	2016-17	WINTE	R CAU	LIFLOW	/ER VA	RIETY	ΓRIAL					
		First Trans	splanting	12th July	2016	(First line)		SPACIN	G 8580	Plants/A	cre	
		Second T	ransplanti	ng 21st J	uly 2016	(Second lin	ne)		(	Crates / Acı	re	
VARIETY	PLOT	CU	TTING PERI	IOD	DAYS	Class	Class	Unmarket	Facepac	k		Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Elsoms	95	7.11	5.12	8.12	31	72	24	4	751	29	129	0
Belot	145	23.11	12.12	19.12	26	84	9	7	761	182	49	0
Average		15.11	8.12	13.12	29	78	17	5	756	106	89	0
	ck material. Dark											
Will turn off	white if not cut	every 4 days. N	ledium de	pth curds.	Some Xa	nthomonas	on outer	leaves of f	irst transp	lanting.		
Enza	96	22.10	31.10	14.11	23	72	22	6	686	114	118	0
Agility	146	28.11	5.12	19.12	21	73	10	17	787	0	54	0
Average		9.11	17.11	1.12	22	72	16	12	737	57	86	0
Uniform. Xan	thomonas on oute	r leaves, clean in	ner leaves.	Good weig	ht. Good cu	ırd cover. Lo	oks good i	n trays. Good	l Facepack	material.		
26% with sm	all white bracts	in earlier trans	planting o	date. Some	small pla	ants/curds v	with later	transplanti	ng.			
Clause	97	31.10	23.11	8.12	38	80	12	8	858	0	64	0
Galiote	147	5.12	8.12	12.12	7	93	0	7	944	0	0	0
Average		17.11	30.11	10.12	23	87	6	7	901	0	32	0
	lium sized frame.	Good wrapper lea	f. A few wh	nite bracts la	ate Octobe	r. Good Face	pack mate	rial. Quick h	eading peri	od for later	transplanti	ng.
	white if not cut								<u> </u>			
Clause	98	8.12	15.12	27.12	19	85	8	7	875	48	45	0
Triomphant	148	22.12	27.12	5.1	14	88	7	5	938	0	40	0
Average		15.12	21.12	31.12	16	87	7	6	906	24	42	0
	sted leaf over curc		epcak mat	erial. Looks	very good	in trays. Sor	ne Xanthor	monas on ou	ter leaves,	clean inner	leaves.	
	ne. Easy to cut a		•						•			
Syngenta	99	15.12	22.12	5.1	21	85	10	5	912	0	54	0
C5049	149	27.12	9.1	18.1	22	90	0	10	965	0	0	0
Average		21.12	31.12	11.1	21	87	5	8	939	0	27	0
	cepack material.	Solid curds. Tight			ner leaves.	Easy to cut	and bag, Ui	niform. Large	e frame. Ve	rv few defe		

	2016-17		WINTE	R CAU	LIFLOW	VER VA	RIETY	ΓRIAL					
			First Trans	splanting	12th July	2016	(First line)		SPACIN	G 8580	Plants/A	cre	
			Second T	ransplanti	ng 21st J	uly 2016	(Second lin	ne)		(	Crates / Acr	е	
VARIETY	PLOT		CU	TTING PERI	IOD	DAYS	Class	Class	Unmarket	Facepac	k		Class 2
			10%	50%	90%		1	2	%	8	6	16	12
Hazera	100		5.12	5.12	15.12	10	84	10	6	794	143	54	0
Lecatis	150		15.12	15.12	27.12	12	88	4	8	944	0	21	0
Average			10.12	10.12	21.12	11	86	7	7	869	72	37	0
Uniform. Very	y good Facep	ack mater	ial. Good cu	rd cover. S	lightly knobl	bly curds. N	ledium dept	h curds. Da	ark green lea	f. Clean lea	ıf.		
Easy to cut	and bag. Ve	ery few de	efects.										
Syngenta	101		14.11	5.12	15.12	31	93	0	7	815	114	0	0
Leocen	152		12.12	12.12	19.12	7	93	0	7	1001	0	0	0
Average			28.11	8.12	17.12	19	93	0	7	908	57	0	0
Dark green le	eaf. Tight wra	pper leave	s. Medium o	depth curds	s. Smooth c	urds. Xanth	omonas on	outer leave	es. Good Face	epack mate	erial.		
Uniform. Lo													
		•				·			•				
Clause	102		31.12	9.1	12.1	12	82	7	11	881	0	36	0
Jerboam	152		31.12	12.1	12.1	12	82	3	15	881	0	24	0
Average			31.12	10.1	12.1	12	82	5	13	881	0	30	0
Twisted leaf	over curds. F	Paler inner		_			_	all plants/c					
					<u> </u>								
Elsoms	103		27.12	31.12	12.1	16	94	2	4	1008	0	11	0
Paciano	153		18.1	19.1	23.1	5	88	2	10	879	86	11	0
Average			7.1	10.1	17.1	10	91	2	7	944	43	11	0
Deep curds.	Very uniform	. Upright le						_	ck material	-			
_ top oa. ao.	y willion	pr. 19.11 10	, goc	- Jul u JOV									
			40.4	23.1	26.1	14	80	10	10	858	0	54	0
Fleome	104		1 171				. ()()		1 10 1	UJU		1 77	ı U
Elsoms Baterno	104 154		12.1 19.1		+		+		a				0
Elsoms Baterno Average	104 154		12.1 19.1 <b>15.1</b>	26.1 26.1	30.1 28.1	11	87 <b>83</b>	4	9	929 <b>894</b>	0	24	0

	2016-17	V	WINTE	R CAU	LIFLOV	VER VA	RIETY	TRIAL					
		F	irst Trans	splanting	12th July	2016	(First line)	)	SPACIN	G 8580	Plants/A	cre	
		s	Second T	ransplanti	ng 21st J	uly 2016	(Second lin	ne)			Crates / Acı	е	
VARIETY	PLOT		CU	TTING PERI	OD	DAYS	Class	Class	Unmarket	Facepac	k		Class
7744211	. 20.		10%	50%	90%	27110	1	2	%	8	6	16	12
Clause	105		31.1	3.2	6.2	6	74	8	18	729	86	43	0
Brick	155		2.2	2.2	9.2	7	72	10	18	751	29	54	0
Average			1.2	3.2	8.2	7	73	9	18	740	58	48	0
	. Tendency to	become lo	ose and to	ırn off white	e. Twisted l	eaf over cu	rds. Fairly u	niform.			•		
Syngenta	106		23.1	26.1	31.1	8	82	10	8	729	200	43	14
5064	156		26.1	30.1	2.2	7	92	4	4	987	0	43	0
			25.1	28.1	1.2	7	87	7	6	858	100	43	7
Very soft leav	es. Paler gree	en inner lea	f. Tall fran	ne. Twisted	l leaf over c	urds. Unifo	rm. Some la	rge curds	from earlier t	ransplantii	ng.		
Very Good Fac	epack mater	ial.											
Hazera	107		3.2	3.2	9.2	6	92	4	4	901	114	21	0
Trevaskis	157		3.2	6.2	6.2	3	90	4	6	944	29	21	0
			3.2	4.2	8.2	5	91	4	5	922	72	21	0
Excellent Fa	cepack mate	erial. Very	y Uniform	. Easy to	see, cut ar	nd bag. Go	ood weight	Good cu	rd cover. C	lean leaf.	1		
Clause	108		9.1	23.1	26.1	17	72	6	22	772	0	21	14
CLX 33610	158		19.1	3.2	3.2	15	68	16	16	729	0	86	0
			14.1	28.1	30.1	16	70	11	19	751	0	54	7
Some variatio	n in plant size	s. 12% sma	all loose c			Facepack ı	material. Uni	form. Solid	curds. Good	l leaf cover	. Looks vei	y good in tr	ays.
Clause	109		2.2	6.2	9.2	7	88	4	8	944	0	21	0
Dadautabla	159		9.2	14.2	20.2	11	76	6	18	815	0	32	0
Redoutable							82						

	2016-17	WINTE	R CAU	LIFLOW	VER VA	RIETY	ΓRIAL					
		First Tran	splanting	12th July	2016	(First line)	1	SPACIN	G 8580	Plants/A	cre	
		Second T	ransplanti	ng 21st J	uly 2016	(Second lin	ne)		(	Crates / Acr	е	
					1							
VARIETY	PLOT		TTING PERI		DAYS	Class	Class	Unmarket	Facepac		10	Class 2
	1112	10%	50%	90%		1	2	%	8	6	16	12
Monsanto	110	31.1	31.1	6.2	6	90	2	8	965	0	11	0
5697	160	6.2	6.2	13.2	7	84	2	14	906	0	12	0
Average		3.2	3.2	10.2	7	87	2	11	936	0	12	0
		Solid curds. Easy	to see, cut a	and bag. Ve	ry uniform.	Good cover	leaves. A f	ew small pla	nts/curds.	1		
No other de	efects.					1		1				<del> </del>
Syngenta	111	3.2	9.2	14.2	11	86	8	6	922	0	43	0
Britten	161	9.2	14.2	20.2	11	83	5	12	858	48	27	0
Average	101	6.2	12.2	17.2	11	85	6	9	890	24	35	0
	an leaf Good wra	pper leaves. Solid					-	, ,				
Ormorm. Oic	The state of the s	pper ieuves. oona	Curus. VCI				lian plants	l l	icw office c			+
												+
Clause	112	3.2	3.2	9.2	6	90	4	6	965	0	21	0
Fleet	162	6.2	9.2	14.2	8	83	7	10	884	0	40	0
Average		5.2	6.2	12.2	7	87	5	8	925	0	30	0
Very Uniforn	n. Clean Leaf. Goo	od curd cover, Very	Good Face	pack mater	rial. Easy to	see, cut an	d bag. Lool	ks very good	in trays. Ve	ery few defe	cts.	
Clause	113	20.2	20.2	23.2	3	88	4	8	944	0	21	
CLX 33611	163	20.2	20.2	23.2	3	96		2	944	114	11	0
Average	103	20.2	20.2 20.2	23.2	3	90	2 3	5	944	57	16	0
	Very good Face	pack material. Eas				_		-	J T T	31	10	
very official	i. Very good Face	pack material. Eas	sy to cut an	pack. 300		Ivieu	uiii/iicavy	Luius.				+
Syngenta	114	20.2	23.2	2.3	10	88	8	4	944	0	43	0
C5067	164	27.2	2.3	8.3	9	86	10	4	922	32	29	0
Average		24.2	27.2	4.3	10	87	9	4	933	16	36	0
Heavy leaf d	op. Some yellow	ing of leaves at ba	se of curds	. Very Good	Facepack	material.						

	2016-17	WINTE	R CAU	LIFLOW	<b>VER VA</b>	RIETY	ΓRIAL					
		First Tran	splanting	12th July	2016	(First line)	)	SPACIN	G 8580 I	Plants/A	cre	
		Second T	ransplanti	ng 21st J	uly 2016	(Second lin	ne)		(	Crates / Acr	е	
VARIETY	PLOT	CI	ITTING PER	IOD	DAYS	Class	Class	Unmarket	Facepaci	•		Class 2
VAIGETT	1201	10%	50%	90%	DATO	1	2	%	8	6	16	12
Clause	115	20.2	23.2	27.2	7	96	2	2	1029	0	11	0
Mascaret	165	20.2	23.2	27.2	7	94	4	2	1008	0	11	14
Average		20.2	23.2	27.2	7	95	3	2	1018	0	11	7
Pale inner lea	f. Difficult to see c	urds due to tight	twisted w	rapper leave	es. Very go	od Facepack	material.	Very few def	ects. Solid	curds.		
								ļ				
Syngenta	116	9.2	14.2	20.2	11	83	7	10	876	24	36	0
C5052	166	14.2	20.2	23.2	9	90	2	8	965	0	9	0
Average		11.2	17.2	21.2	10	87	4	9	921	12	23	0
Smooth curds	s. Upright leaf. Ver	y uniform. Slight	twist to wr	apper leaf o	over curds.	Very good F	acepack m	naterial.				
Elsoms	117	2.2	20.2	20.2	18	55	36	9	548	0	191	0
Gerona	167	20.2	20.2	27.2	7	84	10	6	901	0	54	0
Average		11.2	20.2	23.2	13	70	23	7	725	0	123	0
	Upright leaf. Small	curds. Strap lik	e leaves. To	endency to t	turn off whi	te. Open fra	me. Much k	etter from a	later planti	ng.		
Syngenta	118	14.2	20.2	27.2	13	80	2	18	787	95	9	0
Caylen	168	27.2	2.3	6.3	7	80	0	20	815	57	0	0
Average	100	21.2	25.20	3.30	10	80	1	19	801	<b>76</b>	5	0
	lants/curds other						k Slightly				_	
- Jino Jinan p			(10/9.0			Jac and pac	Ongridy			opuon ille		
Monsanto	119	14.2	20.2	23.2	9	90	8	2	965	0	43	0
Carantic	169	20.2	20.2	23.2	3	92	0	8	987	0	0	0
		17.2	20.2	23.2	6	91	4	5	976	0	22	0

	2016-17	WINTE	R CAU	LIFLOV	VER VA	RIETY	ΓRIAL					
				12th July		(First line)		SPACIN	G 8580	- Plants/Ad	cre	
		Second T	ransplanti	ng 21st J	uly 2016	(Second li	ne)		(	Crates / Acr	е	
					•							
VARIETY	PLOT		TTING PER		DAYS	Class	Class	Unmarket	Facepac			Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Clause	120	15.3	20.3	25.3	10	91	0	9	879	0	0	0
CLX 33518	170	16.3	19.3	25.3	9	92	4	4	987	0	21	0
Average		15.3	19.3	25.3	10	91	2	7	933	0	10	0
Dark green le	eaf. Uniform. Well	covered. Slightly	wide base	to curds. Ve	ry good Fa	cepack mate	erial.			,		
												<u> </u>
												<u> </u>
Elsoms	121	20.2	23.2	27.2	7	92	6	2	987	0	32	0
Madiot	171	23.2	1.3	6.3	11	92	2	6	988	0	11	0
Average		22.2	26.2	3.3	9	92	4	4	988	0	21	0
Very uniform	. Clean leaf. Mediu	m depth curds. C	ood curd p	rotection. V	ery good Fa	acepack ma	terial. Easy	to cut and b	ag. Very fe	w defects.		
Elsoms	122	20.2	27.2	6.3	14	92	3	5	947	48	18	0
Mystique	172	2.3	6.3	8.3	6	90	0	10	965	0	0	0
Average		25.2	3.2	7.3	10	91	2	7	956	24	9	0
Very uniform	. Slightly paler inne	er leaf. Very good	Facepack	material. C	lean leaf. V	ery few defe	cts. Looks	very good in	trays.			
Elsoms	123	23.2	27.2	2.3	7	78	18	4	837	0	97	0
Isadora	173	27.2	6.3	8.3	9	84	0	16	901	0	0	0
Average		25.2	3.2	5.3	8	81	9	10	869	0	49	0
	Some off white cu	rds. Paler inner l	eaf. Uniforr	n in size. So	me good F	acepack ma	terial. A fe	w small plan	ts/curds.			
,					<u> </u>	T .						
Hazera	124	6.3	13.3	13.3	7	92	4	4	987	0	21	0
AE 9307	174	6.3	13.3	13.3	7	94	2	4	1008	0	11	0
Average		6.3	13.3	13.3	7	93	3	4	998	0	16	0
	. Good curd cover.	0.0			cut and nad			S. Clean leaf		dood in tra		

	2016-17	WINTE	R CAU	LIFLOV	VER VA	RIETY	ΓRIAL					
	ETY PLOT	First Tran	splanting	12th July	2016	(First line)	)	SPACIN	G 8580 F	Plants/A	cre	
		Second T	ransplanti	ng 21stJ	uly 2016	(Second lin	ne)		C	rates / Acı	re	
VARIETY	PLOT	CL	ITTING PERI	IOD	DAYS	Class	Class	Unmarket	Facepack	<b>(</b>		Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Elsoms	125	2.3	8.3	13.3	11	90	0	10	965	0	0	0
Tempest	175	6.3	13.3	16.3	10	86	2	12	924	0	11	0
Average		4.3	11.3	15.3	11	88	1	11	945	0	5	0
Very uniform	n. Excellent Facepa	ck material. Tigh	t wrapper l	eaf. Slightly	wide base	to curds. Da	rk green k	eaf. A few sn	nall curds.			

	2016-17	LATE	WINTER	RCAUL	IFLOW	ER VAR	RIETY 1	RIAL				
								SPACIN	G 8580	Plants/A	cre	
									Crates / Acre			
VARIETY	PLOT	CL	CUTTING PERIOD		DAYS	Class	Class	Unmarket	Facepack			Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Elsoms	1	14.4	22.4	26.4	12	73	18	9	739	64	95	0
Cartagena	5	14.4	17.4	26.4	12	71	20	9	763	0	107	0
Average		14.4	20.4	26.4	12	72	19	9	751	32	101	0
	leaf. Strong hab	it. Uniform. Lo	ose wrapp	er leaves.	Will turn	off white if	not cut e	very 3 days	. Good Fa	cepack m	aterial.	
	pth. Slightly oper									•		
Hazera	2	15.3	15.3	20.3	5	82	8	10	879	0	43	0
VT 4013	6	15.3	16.3	25.3	10	74	13	13	794	0	70	0
Average		15.3	16.3	23.3	8	78	10	12	837	0	57	0
	af. Slightly wide	base to curds.	Slightly tv	visted wra	pper leaf o	ver curds.	Uniform.	Very good	Facepack	material.	Soft leaf.	
	y good in trays.											
Elsoms	3	7.4	14.4	17.4	10	98	0	2	1051	0	0	0
Fletcher	7	14.4	14.4	21.4	7	91	5	4	977	0	24	0
Average		10.4	14.4	19.4	9	94	3	3	1014	0	12	0
	m. Excellent Fac	epack materia	I. Deep so	lid curds.	Dark greer	leaf. Ver	y good cu	rd cover. C	ean leaf.	Looks ver	y good in t	rays.
01	4	0.4	7.4	444	44	00		4.4	004	0	00	
Clause	4	3.4	7.4	14.4	11	83	3	14	884	0	36	0
CLX33612	8	7.4	7.4 <b>7.4</b>	17.4 <b>15.4</b>	10 10	94 <b>89</b>	3	2 <b>8</b>	1008	0	21	0
Average Wide base	l l to curds. Not eas	y to pack Goo		_					946 curds Mo		28 b	0
	pack material. So					Very Onlio	liii. Siigiii	IIY KIIODDIY	curus. IVIE	l lan dept	 	
Good race	back material. Sc	Jila Caras. A le	w siliali p	lants/curus	s.							
Elsoms	9	25.3	29.3	3.4	9	74	2	24	772	29	11	0
Douglas	13	25.3	29.3	3.4	9	60	16	24	601	57	64	29
Average		25.3	29.3	3.4	9	67	9	24	687	43	37	15
Dark green	leaves. Light cur	rds. Some sma	II curds. L	arge leafy	frame. So	me good	Facepack	material. S	ome frost	ed curds i	n early Apr	il.
A few multi	headed stumps.	Some small p	lants/curd	S.								

	2016-17	ILAIL	WINIE	R CAUL	<b>IFLOW</b>	ER VAR	RIETY T	RIAL				
		Transplar	Transplanted 12th July					SPACING 8580 Plants/Acre				
									Crates / Acre		е	
VARIETY	PLOT		CUTTING PERIOD		DAYS	Class	Class	Unmarket	Facepac			Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Monsanto	10	13.3	15.3	15.3	2	91	2	7	787	276	12	0
Tenfold	14	15.3	15.3	18.3	3	92	4	4	407	826	21	0
Average		14.3	15.3	17.3	3	91	3	6	597	551	16	0
Dark green I	eaf. Solid curds.	High stump.	Good curd	cover. Wi	II turn off v	vhite if no	t every 2 c	or 3 days. G	ood Face	pack mate	rial.	
Syngenta	11	3.4	7.4	14.4	11	91	4	5	930	64	24	0
Delon	15	3.4	7.4	17.4	14	96	0	4	965	86	0	0
Average		3.4	7.4	16.4	13	93	2	5	947	75	12	0
	n. Strong vein to	leaves. Slight	twist to le	eaf over cu	rds. Sligh	tly wide b	ase to cur	ds. Very go	od Facep	ack mater	ial.	
	fects. Easy to cut											
Elsoms	12	15.3	15.3	25.3	10	92	0	8	987	0	0	0
Jerome	16	13.3	15.3	20.3	7	86	2	12	922	0	11	0
Average		14.3	15.3	23.3	9	89	1	10	955	0	6	0
Solid curds.	Medium depth. [	Dark green sh	ort upright	t leaves. V	ery uniforn	n. Very go	od Facep	ack materia	al. Some r	ingspot or	outer lea	ves.
	over. Slightly cri											
Seminis	16B	25.3	25.3	3.4	9	84	2	14	763	191	12	0
SV0890AC	17B	25.3	29.3	3.4	9	82	5	13	858	32	24	0
Average		25.3	27.3	3.4	9	83	4	13	811	111	18	0
	. Very pale yello	w inner leave	s. Solid cı	urds. Sligh	tly off whit	e. Fairly u	niform. S	lightly poin	ted curd c	over. Wel	covered.	
	ion in plant/curd											
	•											
Syngenta	17	13.3	13.3	15.3	2	100	0	0	815	172	0	0
C5033	21	15.3	15.3	16.3	2	86	2	12	86	1115	11	0
Average		14.3	14.3	16.3	2	93	1	6	451	644	5	0
	. Pale green inn				e large cu		m. Verv a	uick headi				
		Facepack ma					<u> </u>					

	2016-17	LATE	WINTE	R CAUL	IFLOW	ER VAF	RIETY 1	ΓRIAL				
		Transpla	nted 12th .				SPACIN	SPACING 8580		cre		
										Crates / Acr	е	
VARIETY	PLOT	С	CUTTING PER		DAYS	Class	Class	Unmarket	Facepack			Class 2
		10%	50%	90%		1	2	%	8	6	16	12
Elsoms	18	20.3	29.3	29.3	9	87	5	8	930	0	24	0
Vogue	22	25.3	25.3	29.3	4	92	8	0	922	86	42	0
Average		23.3	27.3	29.3	6	90	6	4	926	43	34	0
Very unifor	m. Good curd	d protection. Very	good Face	pack mate	erial. Solic	smooth c	urds. A fe	w off white	curds.			
Syngenta	19	6.3	8.3	13.3	7	94	2	4	236	1029	11	0
C5044	24	6.3	8.3	13.3	7	92	0	8	236	1001	0	0
Average		6.3	8.3	13.3	7	93	1	6	236	1015	5	0
Very pale le	eaf. Slightly	wide base. Large	wrapper le	eaves over	curds. Ve	ry large he	avy curd	s. Very large	frame. V	ery unifor	m.	
Clause	20	29.3	3.4	14.4	16	92	8	0	987	0	43	0
CLX33116	23	25.3	3.4	7.4	13	92	6	2	897	116	33	0
Avelmore												
Average		27.3	3.4	11.4	15	92	7	1	942	58	38	0
Very Unifor	m. Medium s	sized frame. Very	good curd	cover. Me	dium dept	h solid cur	ds. Very g	ood Facepa	ck mater	ial.		
Upright dar	k green leav	es. Slightly paler	green inn	er leaf.								