



HORTICULTURE RESEARCH INTERNATIONAL
KIRTON

Summary Report (1993)

HDC Project BOF17

**NARCISSUS VARIETY ASSESSMENT:
TRIALS AT ROSEWARNE 1955-1989**

Gordon R Hanks

Horticulture Research International
Kirton
Boston
Lines PE20 1EJ

Project Co-ordinator: Johnny Walkers
Project Commenced: October 1990
Project Completed: October 1993
Keywords: Narcissus, Daffodil, Bulbs

724957

723477

WILLINGTON ROAD · KIRTON · BOSTON · LINCOLNSHIRE PE20 1EJ
TELEPHONE BOSTON (0205) ~~722391~~ · FACSIMILE BOSTON (0205) ~~722922~~

CHAIRMAN G.T. PRYCE · CHIEF EXECUTIVE: C.C. PAYNE · COMPANY SECRETARY: T.G. HELLER

AUTHOR OF REPORT

G R Hanks BSc, MPhil, MIHort, CBiol, MIBiol

AUTHENTICATION

I declare that this report represents a true and accurate record of the data available.

.....
(signature)

G R Hanks
Horticulture Research International
Willington Road
Kirton
Boston
Lincs PE20 1EJ

Date 15/3/94

Report authorised by:

.....
(signature)

M B Wood
Head of Station
Horticulture Research International
Willington Road
Kirton
Boston
Lincs PE20 1EJ

Date

**NARCISSUS VARIETY ASSESSMENT: SUMMARY OF TRIALS
AT ROSEWARNE 1955-1989**

CONTENTS

	Page
SUMMARY	1
INTRODUCTION	1
CULTIVATION AND RECORDING	2
THE SUMMARY REPORT	3
NAMES AND CLASSIFICATION	4
QUANTITATIVE DATA	4
FLOWER DESCRIPTIONS	93
ACKNOWLEDGEMENTS	207
BIBLIOGRAPHY	207
FIGURES 1 AND 2	209

COMMERCIAL IN CONFIDENCE

NARCISSUS VARIETY ASSESSMENT: SUMMARY OF TRIALS AT ROSEWARNE 1955-1989

SUMMARY

Narcissus varieties were assessed in a trial at Rosewarne Experimental Horticulture Station over the period 1955 to 1989. The data collected included emergence and flowering dates, flower and bulb yield, flower size and flower description. The data from this work is brought together in the present report, comprising, in all, information on 2186 cultivars and other taxa.

INTRODUCTION

The narcissus collection at Rosewarne Experimental Horticulture Station (EHS), Cornwall was started when a nucleus of cultivars was transferred from the Gulval demonstration centre of the Cornwall County Council on its closure in 1952. Additional cultivars with commercial potential were added periodically, once the price of bulbs had fallen to such a level that commercial quantities might be available within the next five to ten years. The varieties were more than just a collection: the bulbs were grown under standardized conditions, and information on season, flower yield and quality (including vase-life) and bulb yields were recorded. The reports of these assessments form a valuable database for bulb growers.

In 1989, government cuts in the funding of horticultural R&D led to the closure of Rosewarne EHS. The narcissus variety trial was moved to Kirton EHS (now Horticulture Research International - Kirton), Lincolnshire. From 1990, continuing work has been funded under contract from the Horticultural Development Council. The contract provided, first, for completing the reporting of data collected at Rosewarne, and, secondly, for funding an on-going trial at Kirton. The present trial is run along lines similar to those at Rosewarne, although, in the interests of economy, the amount of data collected for each cultivar has been reduced. It is also the intention to reflect the changing needs of the bulb industry closely in the choice of cultivars now represented. Results from the trial at Kirton will be summarised annually, beginning in 1994.

The results of the trial at Rosewarne were presented initially as a series of three reports issued free by NAAS (later ADAS), covering the periods 1955-1963, 1964-1967 and 1968-1971. These reports covered 652, 620 and 197 taxa respectively, and, in addition, the third report listed brief details of 310 varieties which were discarded before being fully recorded, as they were considered unsuitable for flower production in the south-west. Occasional reports were given in the annual reports of Rosewarne EHS. No more detailed reports were issued by ADAS, but some further data covering the period from 1971 to 1989 were summarised in reports issued by HRI Kirton to the HDC in 1991 and 1992, covering an additional 195 taxa. Although assessments were related primarily to potential for bulb and outdoor flower production in the south-west, the data are, of course, also more generally applicable. The spread of flowering season is greater in the south-west than in eastern England; in Cornwall, early cultivars may flower up to six weeks earlier than in the east, while later cultivars flower about the same time in all districts.

The present report brings together in one volume all previously reported material, representing in all 1667 separate taxa, plus information on file which had not previously been reported, and covers 2186 taxa in all. The information presented comprises quantitative data on bulb and flower production and flower descriptions. The table of quantitative data is also available on disk as a Lotus 1-2-3 file.

CULTIVATION AND RECORDING

The Rosewarne soil is of the brown earth type, with a large proportion of shale and quartz fragments from the underlying rocks. It is well drained and of a reasonable depth. The land was cropped in rotation with the usual vegetable and flower crops of the area, which maintained a good structure. A light dressing of a high potash compound fertiliser was worked in before planting. Weeds were controlled by standard herbicides.

The site is very exposed, and strong, often salt-laden, gales occurred, particularly during the winter and early spring. Areas for the trial were protected by 8 ft (2.44 m) high hedges 80 ft to 140 ft (24.4 m to 42.7 m) apart, but additional removable 4 ft (1.22 m) high lath screens were erected across the plots before the flowering season. In spite of this protection, gales still caused serious damage to the more brittle-stemmed cultivars. The winter climate is normally mild, prolonged or hard frost being unusual. The average rainfall, about 1072 mm per annum, is fairly evenly distributed throughout the year.

Before 1962, bulbs were planted in beds and lifted after three years. After 1962 all the cultivars were planted in ridges at 76 cm centres and lifted after two years. Growing in beds will give slightly greater yields than growing in ridges. In the table, flower and bulb yields for the pre-1962 data have been adjusted to give approximate figures equivalent to two-year-down growing. Each cultivar plot consisted of 24 bulbs, comprising where possible eight double-nosed bulbs, eight single-nosed bulbs and eight offsets. Planting was about 13 cm deep to the base of the bulb, and 10 cm apart in two rows in the base of the furrows. The plots were 1.22 m long with 0.9 m cross paths and a 3.1 m path between every ten beds for the erection of lath windbreaks. This wide path spacing, by increasing the root run for the crop, may have tended to give slightly higher flower and bulb yield than a normal field planting.

After 1965 the bulbs were planted in nets. The bulbs were spaced out and fastened 10 cm apart in lengths of tubular plastic netting 2.44 m long. When planted in the plots the net was doubled round to produce the double rows. By using this method, planting and lifting could be done much more quickly with less error, and leaving no bulbs in the ground, although growth might be slightly restricted compared with growing loose. Metal labels attached to the nets prevented mixing.

When necessary, fungicidal sprays were applied to control leaf diseases. All bulbs were hot-water treated, bulbs being pre-warmed at 30°C prior to treatment to prevent damage. Bulbs were treated at planting with aldrin against large narcissus fly (the use of aldrin was not revoked until May 1989). All pesticide treatments and other cultural treatments followed current ADAS recommendations.

New cultivars were obtained from many sources, including Great Britain, Australia, Ireland, The Netherlands, New Zealand, the United States of America and other parts of the world where narcissus are grown. All were planted in isolation for the first year, after which they joined the main trial. After three years, when fully acclimatised, the first assessment was made. In this year many were discarded as unsuitable for commercial cut flower production. The most common faults at this stage were short scapes (stems) and, to a lesser degree, flowers which either hung their heads when young, were small, or which flowered well below the foliage. Cultivars which appeared suitable for cut flowers were grown on for seven to eight years, by which time flower and bulb production could be fully assessed over a period of at least four two-year-down cycles (except in the later years of the trial where recording was curtailed). Standard cultivars, like Carlton, Fortune and Golden Harvest, were kept in the trial throughout, for reference. All records were taken in the year of lifting. For full details of the records taken, see 'Quantitative Data' (below). Flower descriptions were compiled for most of the varieties, in addition to the quantitative data.

THE SUMMARY REPORT

The present report has gathered data from five sources: the three earlier NAAS/ADAS reports, additional information described in the 1991/1992 reports, and data which were on file but which had not yet been reported, the latter comprising not only additional years' data for previously reported varieties, but also data on previously unreported varieties. In compiling this fifth data-set, original files of raw data have been checked to ensure that no variety has been missed. A problem arose with some sets of data which were incomplete, and the policy adopted was to exclude these incomplete years of data where sufficient data from other years was available for a variety, but to include them in cases where this was the only information available for a variety.

The only trial varieties which have not been included in the present report are ones where stocks were lost before assessment could take place, either due to severe basal rot (Windjammer (1Y-Y)) or severe virus infestation (Ballymaster (2W-OOR), Bridesmaid (2W-WPP), Coolgreany (3W-GYY), Drumawillan (2Y-W), Golden Sovereign (1Y-Y), Kathleen (1W-Y), Lime Regent (2Y-Y), Northern Sceptre (2W-YYR), Orion (2W-O), Queenscourt (1W-W), Rio Rouge (2Y-R), Strangford (3W-GYR) and Torch Bearer (2Y-Y)). Other varieties may have developed severe virus or basal rot later, once assessments had begun: these appear in the table without data, and a reason is given in the 'Flower Description' section. Other stocks which were evidently mixed or wrongly named when brought in were also discarded. Duplicate accessions of a few varieties, a few varieties found to be synonymous, and two un-named varieties, have also been excluded.

The method of presenting results varied in the earlier reports. In the 1955-1963 report, dates and flower and bulb yields were quoted for one representative year, while in the 1964-1967 and 1968-1971 reports, dates were quoted for one representative year but yields were averaged across each four years' data; for data presented in the 1992 report and the additional data compiled for the present report, information has been averaged across those years for which it was available, as the number of years data varied considerably in different cases. Records such as flower measurements and vase-life were not recorded each year, and means of the available years were used. Where a variety occurred in more than one of the five data-

sets, the summary data given in the present report are means of the values given by those data-sets.

As in any assessment carried out over many years by many staff, it is likely that some inconsistencies in subjective assessments will occur. Compilation of large amounts of data, much from the pre-computer age, inevitably carries the risk of transcription errors, despite careful checking.

NAMES AND CLASSIFICATION

The name and classification of each cultivar have been checked against 'The International Daffodil Checklist (1989)' of the Royal Horticultural Society, and its supplements (ie, the 15th to 19th supplements (1989 to 1993) to 'The International Daffodil Register (1969)'). For wild species, etc (Division 10), nomenclature has been checked against Fernandes' classification outlined in 'The International Daffodil Checklist (1989)', with colour codes added where appropriate from 'Daffodils to Show and Grow', the abridged classified list of daffodil names of the American Daffodil Society (1989). Notes in parenthesis ('15', '16', 'ADS', etc.) by the variety names in the table, indicate where the supplements to the Checklist, or the American list, should be consulted for additions to or deviations from the 1989 RHS Checklist names and classifications.

Synonyms are included only where alternative names have been used in earlier reports of the trial. Only two cultivars in the list, Welcome (raiser, Oscar Ronalds) and Silver Sand (received from T Harper & Sons) were not traced in these publications, and these are indicated by 'NL' (not listed) in the table. As there is doubt about the synonymy of cultivars California and Pentewan, and as two separate stocks were included in the trial, data for both are given.

Although in a few cases flower colour descriptions differ from classifications given in the RHS Checklist, it has generally been assumed that cultivars in the trial were correctly named; two exceptions were Silvester and Sylvester, which appeared to have been reversed in the trial records and have now been corrected, and a 2Y-O variety given in the trial records as Lothario which may well be Lotherio. In one case, Hayle, two entries appear in the Checklist under this name (2Y and 2Y-YOO), and the identity of the stock in the trial was unclear.

QUANTITATIVE DATA

Where no data appear in the table for a cultivar, it was regarded as poor or unsuitable for flower production in the south-west and discarded from the trial: see 'Flower Descriptions' for details. Where the entry 'nd' appears, this records was not taken.

Years in trial. The years given in the table refer to the second year of the two-year growing cycle, when recording and bulb lifting took place. The earliest and latest years the variety was included in the trial are given.

First shoot emergence date. This is expressed relative to Golden Harvest, eg a value of -10 means emerging 10 days before Golden Harvest, 10 means 10 days after Golden Harvest.

Date of emergence is important in the planning of weed control and disease control spray programmes. Cultivars nearly always emerge in the same order, but the actual date of emergence will vary from year to year according to the weather. The order in which cultivars emerge does not correspond with the order of flowering, and often a late cultivar emerges early and *vice versa*.

First flower date. This is also expressed as days relative to Golden Harvest. It gives the date of the first flower to open. Cultivars normally follow each other in the same order although the date at which flowering starts will vary widely, being influenced both by the previous year's growing temperature and the weather in the current season. The spread of flowering season of narcissus varies from year to year according to the weather. Flowering will often be delayed at the start of the season, but the latest cultivars flower at about their usual date.

There are considerable variations in flowering dates within each division, but the usual order of flowering is:-

1. Yellow trumpets and some coloured large cups;
2. Bicolors and white trumpets and large red cups with yellow perianths;
3. Large cupped cultivars with white perianths (pink and red cupped later than yellow or orange cupped) and small cupped cultivars with yellow perianths;
4. Small cupped cultivars and poeticus cultivars; and
5. Narcissus Plenus (synonym Poeticus Flore Pleno).

Duration of flowering (number of days from first to last flower open) is not included: figures can be misleading as they depend to a certain extent on weather conditions prevailing at the time of flowering.

Percentage bulb weight increase. This is calculated as:

$$\frac{(\text{Weight lifted after 2 years} - \text{Weight planted}) \times 100}{\text{Weight planted}}$$

Thus, a percentage weight increase of 100% means that the lifted weight was double the planted weight, or a 200% increase means three times the weight of bulbs was lifted as was planted. Basal rot can seriously reduce yields in some varieties, and where it was seen this is noted in the table.

In the case of data from the period 1955-1963, only three-year-down data were available: these figures have been multiplied by a factor (0.595) based on research at Rosewarne (see Rees, Wallis & Tompsett, 1973), to arrive at yield increases equivalent to two-year-growing.

Number of flowers. This is expressed as the number of flowers per planted bulb obtained in the second year. As in the case of bulb yield, three-year yields for 1955-1962 data have been converted to two-year data (by multiplying by 0.533, see NAAS, 1963, p.9).

The greatest yields of flowers are found among large-cup cultivars, but trumpets tend to give lower yields. Within each division there is no marked difference between crops from coloured and white cultivars.

Leaf poise. This is described as erect (E), spreading (S), or bent (B). See Figure 2.

To the flower producer the appearance of the leaves is not important since these are left on the plant and not picked unless the stock is to be discarded, but the height of the leaves in relation to the flowers can be critical. If the leaves are level with, or higher than, the flowers it is possible for the leaves to chafe against the buds during storms causing such severe mechanical damage that the flowers become unmarketable. Cultivars with leaves shorter than the flower stem will escape damage, as will those with longer foliage which is spreading or bent away from the flower buds.

Stem length. This was measured from ground level to the bend of the neck at the 'gooseneck' stage.

Apart from the need for the flowers to stand well above the foliage, a good length of stem is needed for market, especially for flowers picked in bud. Stems lengthen considerably in the last two or three days before the flower opens, and this extra stem length is lost when picked in 'pencil' or 'tight gooseneck' stage. Stems of the earliest flowering cultivars are frequently rather short, but at present few longer stemmed earlies are available. The bottom 5 cm of the stem may be lost in picking.

Perianth diameter and corona (trumpet) length and diameter. Recorded across the widest extent of the fully open flower.

Flower position. Recorded as above foliage (A), or below foliage (B).

Flower substance. This is a measure of the thickness of the perianth and corona: poor (P), medium (M), good (G), or very good (VG).

The substance of the flower has little relation to the length of life of the flower in water: the longest lived flowers are often found among the smaller, lighter types.

Vase-life. Flowers were picked at the gooseneck stage with colour showing, with the exception of doubles which were more fully developed. They were placed in standard conditions similar to those expected in a heated living room. The temperature was controlled at 16°C with 65% relative humidity, with fluorescent lighting for 12 hours per day. The flowers usually opened within 24 hours and the measurements of vase-life started from the date of picking. It finished when the edges of the perianth segments started to turn brown. In most narcissus the corona persists for some time after this, but in a few varieties such as Glynver and Spry, the corona edge dies first. Browning of the perianth is usually preceded by increasing transparency of the petals.

A defect which may occur during the vase life of the flower is excessive growth of the necks, a fault which may even render a cultivar unsuitable for sale. This character occurs most frequently in Divisions 2, 3 and 9, but it is less common in white cultivars of Divisions 2 and 3. Often cultivars with naturally long necks tend to grow even longer in water. If the necks grow stiffly in water, the flowers are acceptable, since they still face at right angles to the stem.

Occasionally some cultivars under certain conditions of temperature and humidity may be affected by *Botrytis* (smoulder) on the flowers, which could reduce the vase-life by 3 to 4 days. It should be noted that flowers picked as green buds would be less likely to be affected.

A vase life of 7 to 8 days from the date of picking is considered to be satisfactory.

Disease notes. Where basal rot or virus were noted in field plots, or smoulder became evident during vase-life testing, this is noted in the final column of the table (B, V or S, respectively). No attempt has been made to record the severity of disease.

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
ABABA	2W-O	1972 - 1987	4	10	186	2.82	S	39	10.1	1.6	3.6	A	M	9.1	
ABADAN	2W-YRR	1955 - 1967	12	12	239	2.08	SB	42	9.0	1.8	4.1	A	G	8.1	
ABALONE	2W-YYP	1976 - 1981	6	28	190	2.04	S	43	11.4	2.6	4.1	B	VG	10.3	S
ABEL TASMAN	11W-WYO	1976 - 1981	21	35	157	2.03	S	36	10.0	3.0	6.7	B	VG	9.7	
ABELARD	2Y-YOO	1955 - 1963	4	32	408	1.87	SB	49	11.0	1.8	3.8	A	G	nd	
ACCENT	2W-P	1976 - 1982	39	31	181	2.05	S	35	10.9	3.0	4.1	A	G	9.3	
ACCOLADE	3W-R	1973 - 1973	0	38	202	2.12	S	48	10.0	1.1	2.5	A	VG	11.8	
ACROPOLIS	4W-R	1977 - 1984	10	42	180	1.47	SB	49	10.7	1.1	nd	A	G	8.4	S
ACTAEA (16)	9W-YJR	1955 - 1987	6	31	276	2.04	SB	52	9.4	0.6	1.9	A	G	9.1	
ACTRICE	2Y-O	1973 - 1976	28	26	148	2.83	S	37	11.5	2.5	5.0	A	G	8.5	V
ADA FINCH	1W-W	1968 - 1982	19	5	138	2.08	S	36	11.5	5.0	5.8	B	G	9.0	VS
ADAM	3Y-O	1955 - 1973	23	27	239	1.95	SB	50	10.0	1.2	3.2	A	VG	8.4	
ADAMANT	2Y-YYO	1955 - 1963	7	8	361	1.87	SB	38	10.0	2.8	3.0	B	M	nd	
ADELA	2W-Y	1955 - 1963	-7	17	199	1.87	S	47	11.0	2.9	3.8	A	G	nd	
ADMIRATION	8Y-ORR	1955 - 1963	-11	23	52	1.60	SB	38	6.0	0.7	1.7	A	M	nd	
ADORNMENT	2Y-O	1972 - 1972	-11	16	244	1.22	S	43	11.0	3.1	4.7	A	G	10.0	
ADVANCE GUARD	1Y-Y	1955 - 1963	32	12	173	1.33	S	44	12.0	5.2	6.5	A	G	nd	
ADVENTURE	2Y-Y	1955 - 1963	-7	12	252	1.87	S	42	11.0	4.6	4.1	A	G	6.8	
AEROLITE	2Y-Y	1955 - 1963	21	9	303	2.13	S	37	12.0	3.9	5.5	A	G	nd	
AFLAME	3W-YOO	1955 - 1984	-3	16	224	2.38	SB	54	10.0	1.7	3.2	A	G	9.6	
AFTER ALL	3W-YJR	1973 - 1980	32	64	91	2.06	S	35	10.7	1.5	3.9	B	VG	9.8	
AGATHON	2Y-Y	1955 - 1972	1	14	216	2.05	SB	44	12.0	4.1	4.4	A	G	7.7	
AGNES ANDREA	2W-O	1964 - 1967	47	24	nd	nd	SB	34	10.0	1.9	4.9	A	G	6.6	
AGNES MONTEFIORE	2W-WWY	1955 - 1980	10	14	175	2.46	S	42	12.0	4.2	5.5	A	G	9.3	V
AGORA	2W-O	1975 - 1987	36	26	136	2.56	B	43	11.0	1.7	4.7	B	M	7.6	
AHOY	11W-Y	1964 - 1967	41	38	208	1.50	S	29	10.0	3.0	7.3	A	M	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
AIR MARSHAL	2Y-O	1972 - 1972	-9	29	103	1.38	S	40	11.5	2.1	2.6	nd	VG	8.6	
AIRCASTLE	3W-Y	1976 - 1981	22	38	131	2.08	S	43	11.1	1.4	4.1	B	G	10.2	
ALABASTER	4W-W	1976 - 1984	68	65	141	1.35	S	43	7.7	0.5	1.2	A	G	11.8	
ALAMEIN	2Y-O	1955 - 1982	3	20	272	2.80	S	46	10.0	2.2	3.1	A	G	9.6	
ALARM	2Y-O														
ALAYNE	2W-YYO	1968 - 1980	23	29	191	2.32	SB	44	11.5	2.6	4.2	A	G	8.2	B
ALBANY	8W-YOO	1955 - 1963	-28	30	238	2.93	SB	62	9.0	0.9	2.7	A	G	nd	
ALBATROSS	3W-YYO	1955 - 1967	19	32	351	2.76	SB	60	10.0	1.1	2.2	A	P	8.0	
ALBERT SCHWEIZER	1Y-Y														
ALCATRAZ	1Y-Y	1979 - 1981	18	20	162	1.83	S	39	13.5	5.5	7.0	A	VG	11.1	S
ALDERGROVE	2W-Y														
ALEPPO	3W-R	1955 - 1979	10	34	212	2.07	SB	48	10.0	1.3	3.2	A	VG	8.9	
ALEX FLEMING	2W-O	1967 - 1969	4	22	219	1.41	S	31	9.5	1.4	3.5	A	G	7.7	
ALGECIRAS	3W-R	1964 - 1967	17	46	345	1.80	SB	44	8.5	0.9	2.9	A	VG	7.2	
ALGIERS	3W-O	1968 - 1972	43	33	191	1.48	SB	42	9.8	1.6	2.8	A	G	9.1	
ALICANTE	2W-O	1968 - 1973	38	46	140	1.66	SB	43	12.2	2.2	3.7	A	VG	10.4	
ALIGHT	3Y-O	1955 - 1963	14	12	376	1.87	S	47	10.0	1.7	2.8	B	M	nd	
ALL GLORY	1W-Y	1968 - 1976	49	30	138	2.17	S	37	11.8	4.8	5.0	A	G	8.8	BS
ALLUREMENT	2W-P	1976 - 1987	21	22	165	2.59	S	36	10.1	2.5	4.4	A	M	7.8	
ALPHONS DIEPENBROCK	1Y-Y	1972 - 1972	37	19	110	1.00	S	35	11.5	5.5	5.7	A	G	nd	
ALPINE EAGLE	1W-Y	1964 - 1967	51	29	nd	nd	S	37	11.5	4.8	4.9	A	VG	8.8	
ALROI	2Y-Y	1955 - 1963	32	15	797	1.87	S	44	10.0	3.8	4.3	A	G	nd	
ALTYRE	3W-W	1964 - 1967	34	32	206	1.90	SB	38	11.5	1.7	3.3	B	G	7.6	
ALVERADO	2W-Y	1968 - 1968	71	26	267	2.60	S	34	11.0	3.3	4.7	A	G	8.0	
ALYCIDON	1W-W	1968 - 1972	50	33	138	1.51	S	31	11.0	4.6	4.6	A	G	8.5	BS
AMATEUR	3W-O	1955 - 1963	27	24	169	1.33	S	44	10.0	1.7	5.0	A	M	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
AMBASSADOR	1W-W	1964 - 1967	53	34	nd	nd	S	30	10.5	4.8	4.8	A	G	9.4	
AMBER FLASH	2W-O														
AMBERGATE (17)	2O-O	1976 - 1982	5	38	164	1.64	B	46	9.6	1.5	3.4	A	G	10.7	
AMBERLEY	2Y-Y	1955 - 1973	2	24	295	1.84	S	40	11.0	2.9	3.8	B	VG	11.0	
AMBOYNA	1W-Y	1955 - 1963	4	18	299	1.87	S	34	10.0	4.5	4.7	B	M	nd	
AMIABLE	3Y-O	1955 - 1963	7	35	198	1.87	SB	56	10.0	1.3	3.5	A	M	nd	
AMOR	3W-YYO	1980 - 1984	9	25	171	2.04	SB	41	13.3	2.0	5.0	A	VG	8.3	
ANACAPRI	3W-YYR	1976 - 1985	21	44	109	1.71	S	50	10.3	1.0	2.7	A	G	8.7	
ANAKIM	1W-Y														
ANDALUSIA	6Y-O														
ANDES	1W-W														
ANDROCLES	4W-W	1973 - 1973	32	3	178	1.50	S	21	9.5	4.2	4.0	A	M	7.8	V
ANGELINE	3W-WWY	1964 - 1967	19	16	275	2.10	SB	36	9.0	1.5	2.8	A	M	7.0	
ANGEL'S WINGS	2W-W	1955 - 1982	10	10	202	1.77	SB	40	10.0	2.5	4.5	B	G	8.2	
ANN ABBOTT	2W-P	1968 - 1980	38	31	137	2.22	SB	38	10.5	2.7	3.5	A	G	8.4	S
ANNAPURNA	1W-W	1964 - 1967	11	30	136	1.50	S	29	11.5	5.2	5.6	B	G	8.8	
ANNE FRANK	4W-O	1977 - 1979	28	35	186	1.64	SB	44	10.0	1.2	nd	A	G	8.7	V
ANTRON	2Y	1972 - 1978	13	24	139	2.50	SB	43	9.5	2.0	2.6	A	G	9.3	S
ANTWERP	1Y-Y	1964 - 1967	40	28	nd	nd	S	33	10.5	5.1	4.2	A	G	6.2	
ANZENA	2W-Y														
APPROVAL	2Y-OOR	1968 - 1972	24	24	120	2.26	SB	40	10.8	2.0	3.8	A	G	8.4	
APRICOT BEAUTY	2W-O	1964 - 1967	-21	27	nd	nd	S	29	12.0	3.2	6.0	B	G	7.0	
APRICOT DISTINCTION	3Y-R	1955 - 1963	-14	24	400	1.87	SB	54	10.0	1.4	3.4	A	G	nd	
APRICOT SENSATION	2W-Y	1972 - 1972	43	34	36	0.50	SB	24	10.0	2.3	4.9	B	G	9.4	
APRIL HARVEST	1W-Y	1977 - 1979	22	20	209	1.94	S	38	10.5	4.6	4.8	A	G	8.7	
APRIL MERIT	2W-W	1977 - 1978	48	25	268	1.60	S	33	12.5	4.8	4.0	B	G	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Petianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
APRIL MESSAGE	1W-W	1977 - 1981	34	22	141	1.76	S	40	11.2	4.6	4.3	A	G	10.7	BS
APRIL ROSE	2W-P	1979 - 1981	76	47	228	1.82	SB	32	10.0	3.4	3.4	B	M	9.8	
APRIL TEARS	5Y-Y	1964 - 1967	40	52	288	2.90	S	27	4.5	0.9	1.4	B	M	8.9	
ARANJUEZ	2Y-YYO	1955 - 1981	13	20	229	2.66	S	49	9.1	1.8	3.4	A	VG	9.0	
ARBAR	2W-O	1972 - 1973	-2	22	92	1.67	S	43	10.8	1.7	3.6	A	VG	10.4	S
ARCADIA	2W-O	1970 - 1987	31	29	120	2.09	B	39	10.7	1.8	4.0	A	G	9.1	B
ARCTIC DORIC	2W-WW	1972 - 1980	18	17	137	1.90	S	38	10.7	3.6	3.2	A	G	10.4	BS
ARCTIC GOLD	1Y-Y	1964 - 1985	17	23	193	2.25	E	40	10.2	4.2	3.7	A	VG	9.3	
ARCTIC MOON	2W-W	1968 - 1973	27	17	169	1.74	S	36	11.0	4.0	5.3	A	G	7.2	B
ARCTIC PRINCE	1W-W	1968 - 1972	55	33	113	1.75	SB	32	10.8	4.8	5.0	A	G	8.2	BS
ARDARA	2W-W	1968 - 1972	58	51	115	1.34	SB	38	10.8	3.1	4.1	A	G	7.8	B
ARDBANE	2W-W	1972 - 1979	36	32	94	1.76	S	37	10.8	3.8	4.0	A	G	10.1	S
ARDELJNIS	1W-W	1955 - 1963	36	20	130	1.33	S	34	11.0	4.6	4.2	B	VG	8.0	
ARDOUR	3Y-R	1972 - 1980	13	22	178	2.55	SB	46	9.8	1.5	2.6	B	G	8.8	
ARGENT	4W-Y	1964 - 1967	5	16	nd	nd	SB	40	9.5	1.9	nd	A	P	7.4	
ARGOSY	1Y-Y	1972 - 1973	23	29	126	1.98	S	30	11.0	4.8	4.4	A	G	8.4	S
ARGUROS	3W-GOR	1964 - 1967	-5	46	nd	nd	SB	52	10.5	1.0	2.7	A	G	10.0	
ARGYLL	1Y-Y	1968 - 1973	39	24	108	1.36	S	35	11.4	4.6	4.8	A	G	8.2	BS
ARK ROYAL	1Y-Y	1968 - 1982	40	24	145	2.02	S	36	1.1	4.5	4.8	A	VG	7.8	S
ARLINGTON	2W-YOY	1955 - 1963	17	20	210	1.33	S	50	10.0	1.8	4.0	A	G	nd	
ARMADA	2Y-O	1955 - 1984	-23	8	182	2.34	S	46	11.0	2.6	3.6	A	VG	8.7	
ARMAGH	1Y-Y	1972 - 1975	40	27	230	2.47	S	31	11.0	4.9	4.9	A	G	8.3	
ARNHEM	3Y-O	1968 - 1982	66	48	170	2.46	SB	40	10.5	1.2	3.4	A	G	8.3	B
ARRANMORE	1Y-Y														
ARRIBA	2Y-O														
ARTILLERY	3W-R	1972 - 1987	17	34	173	1.98	B	44	10.7	1.5	3.3	A	G	9.6	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
ARTOIS	1Y-Y														
ARUBA	2Y														
ASCOT	4Y-Y	1977 - 1984	9	33	154	1.76	S	42	10.8	2.3	nd	A	G	10.3	S
ASKELON	1W-W	1964 - 1967	13	24	nd	nd	S	32	10.5	4.9	4.4	A	G	8.8	
ASPASIA	8W-Y	1955 - 1981	7	32	301	3.47	SB	52	6.0	0.5	1.5	A	M	9.2	
ATATURK	2Y-O														
ATHFORD	1Y-Y	1955 - 1963	36	38	112	1.87	E	42	10.0	4.7	4.0	B	G	nd	
ATLAS	1Y-Y	1972 - 1972	31	29	181	1.05	S	35	11.5	4.8	5.6	A	VG	9.0	
ATTRACTIVE	1Y-Y	1982 - 1982	45	-12	294	1.64	nd	nd	nd	nd	nd	nd	nd	nd	
AUDA	4Y-Y	1976 - 1976	43	31	243	0.96	SB	39	10.0	2.0	nd	A	M	8.6	V
AUDUBON	3W-WWR	1976 - 1981	3	32	211	2.00	SB	41	11.1	1.7	3.5	B	G	9.8	
AUGUST GREEN	1W-Y	1976 - 1978	28	16	306	0.90	S	35	12.0	4.9	4.8	B	VG	10.4	
AUGUST PINK	2W-P	1972 - 1981	26	15	168	2.33	S	35	9.6	3.4	3.6	A	G	9.0	
AURELIA	7Y-Y	1955 - 1967	-10	36	220	1.07	SB	38	8.0	2.6	2.1	B	G	9.0	
AVALANCHE	8W-Y														
AVE	2W-W	1955 - 1963	7	24	458	0.80	SB	42	12.0	3.8	3.9	A	G	9.4	
AVENGER	2W-O	1973 - 1987	5	31	147	1.99	B	46	9.7	1.5	3.6	A	G	10.0	
AVIEMORE	2Y-YOO	1955 - 1963	17	26	273	1.87	SB	10	2.9	1.9	1.9	G	nd	nd	
AYALA	2Y-O	1981 - 1983	39	48	278	1.50	B	40	10.5	1.8	3.4	A	VG	nd	
BABY DOLL	6Y-Y	1955 - 1967	42	20	158	3.25	S	26	9.0	3.9	3.0	A	G	6.8	
BABY MOON	7Y-Y	1964 - 1967	-23	57	nd	nd	SB	25	3.5	0.4	1.1	B	M	14.4	
BACCARAT	11Y-Y	1964 - 1981	35	27	194	2.04	S	48	9.5	3.2	8.2	A	G	10.6	B
BACH	1Y-Y	1955 - 1967	24	9	220	2.20	E	34	10.0	4.2	4.7	A	M	7.1	
BACKHOUSE'S GIANT	2Y-O	1964 - 1967	-5	18	133	2.20	S	34	12.0	4.1	3.9	B	G	9.0	
BADGER	2Y-O	1968 - 1971	75	46	160	1.50	SB	39	11.0	2.6	3.3	B	VG	9.8	
BAHRAM	2Y-O	1968 - 1982	22	14	196	2.56	S	40	10.8	2.0	3.7	A	G	9.5	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
BALALAIKA	2Y-YR	1968 - 1973	35	27	212	2.82	SB	40	9.9	1.8	3.2	A	G	9.2	S
BALAMARA	2Y-Y	1976 - 1987	14	-14	186	1.81	E	44	10.2	3.9	3.9	A	M	9.5	
BALI HAI	4W-R	1973 - 1981	-4	33	159	2.08	SB	43	9.6	1.4	nd	A	G	9.8	
BALLINTOY	2Y-R	1964 - 1967	8	43	238	2.80	SB	47	10.0	2.0	2.6	A	G	10.6	
BALLOO	2W-W														
BALLYCASTLE	3W-WWO	1964 - 1967	40	37	297	2.70	SB	41	10.0	1.2	3.4	A	G	8.8	
BALLYFERIS	1W-Y	1955 - 1963	-4	26	141	1.87	E	40	11.0	4.4	4.2	A	M	nd	
BALLYGARVEY	1W-Y	1973 - 1981	40	27	140	2.24	SB	32	10.2	4.5	4.2	A	G	9.4	BS
BALLYMARLOW	2Y-R	1964 - 1981	11	25	271	2.89	SB	37	10.0	2.0	2.5	A	G	9.0	
BALLYMARTIN	2W-GYY	1972 - 1975	27	49	203	1.36	SB	30	9.5	1.6	4.3	B	G	9.2	V
BALLYSILLAN	3Y-R	1964 - 1967	-32	24	nd	nd	SB	41	11.0	1.4	2.5	A	G	8.8	
BALLYWALTER	1W-Y	1972 - 1973	32	31	155	1.31	S	30	11.2	4.9	4.8	A	G	10.2	
BAMBI	1W-Y	1964 - 1967	20	-19	280	0.90	S	14	6.5	2.4	2.7	A	P	6.0	
BAMBINO	2Y-O	1955 - 1981	-5	6	252	2.32	S	43	10.0	2.8	3.2	A	VG	9.9	
BANBRIDGE	1Y-Y	1964 - 1967	17	10	nd	nd	S	33	11.0	4.8	4.1	A	VG	9.2	
BANDSMAN	2Y-Y	1955 - 1963	-7	15	309	2.40	S	46	9.0	3.0	3.0	A	M	nd	
BANN	2W-P	1955 - 1963	-4	17	211	1.87	S	34	10.0	3.2	3.8	B	G	nd	
BANTAM	2Y-OOR	1964 - 1967	40	41	nd	nd	SB	30	7.0	1.2	3.0	B	G	9.4	
BANTRY	3W-YR	1955 - 1963	17	32	350	1.87	SB	62	10.0	1.3	3.5	A	VG	9.0	
BARBARA ANN SCOTT	3W-YYO	1964 - 1967	40	17	219	1.90	SB	43	10.5	1.5	3.9	A	G	8.6	
BARDIA	2Y-O	1955 - 1963	14	17	325	1.87	SB	41	10.0	2.1	3.4	B	M	nd	
BARLOW	6Y-YWW	1973 - 1974	23	-6	342	1.60	SB	30	8.8	2.9	2.4	A	M	nd	V
BARRETT BROWNING	3W-O	1964 - 1967	-2	12	236	2.46	S	40	9.6	1.4	3.6	A	M	8.3	S
BARTIZAN	2Y-O	1955 - 1963	-14	12	295	2.40	SB	44	11.0	2.4	3.4	B	M	nd	
BARTLEY	6Y-Y														
BASHFUL	2W-W	1955 - 1967	2	19	311	2.38	S	42	10.0	2.1	3.1	A	G	10.0	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
BASTION	1Y-Y	1955 - 1979	37	32	154	1.72	S	37	10.6	4.4	4.0	B	G	9.9	V
BATH'S FLAME	3Y-O	1955 - 1967	-10	12	496	3.12	SB	44	10.5	2.3	3.3	A	M	8.7	
BATTLEAXE	2Y-O	1955 - 1963	21	21	292	1.87	SB	46	10.0	2.4	3.1	A	M	nd	
BAWNBOY	1Y-Y														
BAYARD	1Y-Y														
BAYEUX	1Y-Y	1955 - 1963	14	8	255	1.87	E	42	10.0	4.6	4.6	A	G	nd	
BEALITA	3Y-O	1973 - 1980	5	29	201	2.29	SB	44	9.9	1.4	3.5	B	G	9.6	B
BEATEN GOLD	1Y-Y	1968 - 1981	34	21	160	2.00	S	42	10.8	4.5	4.8	A	G	8.8	B
BEAU MONDE	2W-O	1972 - 1972	66	31	188	2.00	S	24	10.0	2.5	5.2	B	G	7.0	
BEDOUIN	3W-O	1964 - 1967	13	33	nd	nd	SB	39	9.0	1.3	2.6	A	M	6.6	
BEECHER STOWE	3W	1955 - 1963	-4	17	273	1.87	SB	41	10.0	2.0	4.1	A	G	nd	
BEEFEATER	2Y-O	1972 - 1972	-28	19	198	2.38	S	37	11.5	2.6	4.2	A	G	8.4	
BEERSHEBA	1W-W	1955 - 1967	18	15	159	1.66	SB	38	12.0	4.9	3.9	B	G	9.8	
BEGUILDY	3Y-GGR														V
BEIRUT	2W-YR	1955 - 1963	-11	15	367	1.87	SB	46	10.0	1.4	3.5	A	VG	nd	
BELISANA	2W-O	1968 - 1977	30	30	178	2.00	SB	46	11.8	2.4	4.9	A	VG	7.6	S
BELLA VISTA	2W-ROO	1968 - 1982	49	46	180	2.02	SB	51	10.0	1.2	3.0	A	VG	9.0	
BELLE CHINOISE	3W-O	1955 - 1963	32	37	493	2.40	SB	46	8.0	0.6	1.9	A	P	nd	
BELLEEK	2W-P	1964 - 1967	72	56	nd	nd	S	22	10.0	3.1	4.6	B	G	9.6	
BELLEVUE	2W-O	1964 - 1967	37	25	nd	nd	SB	33	11.0	2.6	3.8	A	G	9.2	
BELTANY	1Y-Y														
BELTRIM	2W-GPP	1982 - 1982	85	41	367	2.00	nd	nd	nd	nd	nd	nd	nd	nd	
BELYTA	2W														
BENE	1Y-Y	1974 - 1976	23	22	185	0.50	S	34	9.0	3.6	3.8	A	G	nd	V
BENEDICITION	3W-W	1973 - 1981	52	44	149	2.05	SB	45	11.1	1.2	2.7	A	G	9.5	S
BENGHAZI	2Y-YOO	1964 - 1967	-15	18	238	2.50	S	40	11.0	2.4	2.8	A	G	9.6	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
BERGEN	2W-W	1955 - 1963	36	22	162	1.87	S	32	10.0	3.9	3.0	A	G	nd	
BERIMA	2Y-Y	1974 - 1975	28	10	352	1.59	S	34	10.5	3.8	3.5	A	G	9.6	V
BERMUDA	2Y-O	1955 - 1973	4	16	211	1.92	SB	44	12.0	2.5	3.6	A	G	8.2	
BERNARDINO	2W-YYO	1955 - 1967	12	22	253	1.98	S	44	10.0	2.2	3.8	A	G	7.8	
BERTHA ATEN	2W-O	1955 - 1963	52	40	236	1.33	SB	40	9.0	1.3	3.8	A	G	nd	
BERYL (17)	6W-YYO	1955 - 1967	18	11	274	3.56	S	27	8.0	1.2	1.7	A	M	10.0	
BESSBROOK	2W-W	1972 - 1973	31	37	158	1.56	S	37	10.8	3.2	4.8	A	G	12.7	S
BETTER TIMES	2Y-YYO	1955 - 1963	0	10	204	2.13	SB	42	12.0	3.0	4.8	B	G	nd	
BIANCA	1W-W														
BIBKIE	2W-Y	1972 - 1972	31	30	273	3.50	S	32	9.0	2.5	3.6	B	G	11.0	
BIG CHIEF	1Y-Y	1979 - 1981	8	18	170	2.06	S	32	14.5	5.5	6.7	B	VG	8.9	S
BIG KEITH	1Y-Y	1976 - 1976	44	13	187	0.67	S	42	11.0	4.5	4.3	A	G	nd	V
BIG WJG	4W-Y	1976 - 1981	5	35	138	2.07	SB	44	11.2	2.8	nd	B	G	8.8	
BILEE	1Y-Y	1978 - 1980	0	-4	285	1.83	S	36	12.5	5.0	5.7	A	G	12.0	
BILL WEBBER	2W-O	1976 - 1981	-1	14	270	2.30	SB	40	11.4	3.7	5.0	A	G	9.5	
BINDA	1W-P	1976 - 1980	41	17	158	0.87	S	29	10.8	4.3	3.4	B	VG	nd	
BINKIE	2Y-W	1955 - 1967	-7	21	322	3.72	S	40	10.0	2.6	3.7	B	M	11.0	
BIRD OF DAWNING	1Y-Y	1971 - 1987	13	5	153	2.32	S	36	11.9	5.5	5.3	A	G	10.0	
BIRMA	3	1955 - 1973	7	12	279	2.07	SB	43	10.3	1.9	2.4	A	M	9.0	
BISMARCK	1W-W	1955 - 1963	21	38	206	1.87	S	45	11.0	4.6	3.4	A	G	nd	
BITHYNIA	3W-WWO	1973 - 1973	-5	43	198	2.75	S	46	11.0	1.7	4.0	A	G	12.0	S
BIZERTA	2W-Y	1955 - 1963	29	20	319	2.40	S	38	10.0	2.6	3.7	B	VG	nd	
BLACK PRINCE	9W-GYR	1955 - 1963	-7	52	227	1.87	SB	58	8.0	0.5	1.7	A	M	nd	
BLAIR HOUSE	3W-O	1968 - 1982	55	43	176	2.34	SB	44	10.5	1.4	4.3	B	G	7.8	
BLANQUET	4W-Y	1977 - 1980	88	68	209	1.69	S	38	8.0	0.4	nd	A	G	9.8	
BLARIS	2W-P	1972 - 1982	19	31	158	2.51	S	36	10.8	1.9	2.9	A	G	11.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
BLARNEY	3W-OOY	1955 - 1975	16	35	201	2.20	SB	56	10.0	1.2	3.4	A	G	10.0	
BLARNEY'S DAUGHTER	2W-OOY	1968 - 1980	24	32	135	2.16	SB	42	11.5	1.8	3.8	A	G	9.0	B
BLEASBY GORSE	1Y-Y	1976 - 1987	4	0	123	1.86	S	45	10.8	5.1	5.3	A	M	8.9	B
BLenheim	1Y-Y	1971 - 1971	65	33	145	1.10	S	36	10.0	4.7	4.2	A	G	6.2	S
BLISS	1W-W	1968 - 1977	52	14	188	1.91	S	36	11.5	4.3	4.6	A	G	6.4	B
BLOEMFONTEIN	2Y-YYO	1973 - 1982	-12	22	188	2.61	SB	50	10.5	2.3	3.7	A	G	10.5	
BLUSH	2W-P	1964 - 1967	28	33	200	2.40	S	37	10.0	4.4	4.2	A	G	9.0	
BLUSTER (18)	2Y-R	1968 - 1971	47	36	142	2.00	SB	37	10.0	2.2	3.6	A	G	10.8	B
BO PEEP	2W-P	1967 - 1967	52	55	384	1.80	SB	38	11.5	2.5	3.2	A	G	8.6	
BOBOLINK	2W-YYO	1973 - 1987	-2	3	173	2.52	S	41	10.4	2.3	5.0	A	G	9.6	
BODILLY	2W-Y	1955 - 1967	8	14	212	2.14	SB	40	10.0	3.4	4.1	A	G	7.8	
BOKHARA	2Y-O	1955 - 1963	0	12	209	2.40	SB	43	10.0	2.1	3.0	B	G	nd	
BOLTON	7Y-Y	1955 - 1967	-10	27	264	2.50	E	48	8.0	1.3	1.7	B	M	8.0	
BOMBAY	2Y-YYR	1968 - 1982	33	46	130	1.76	SB	44	10.8	2.0	3.0	A	VG	9.1	
BONAPARTE	2Y-Y	1955 - 1963	10	4	133	2.40	S	42	10.0	2.6	3.6	A	P	nd	
BONFIRE	3Y-O	1955 - 1963	7	31	302	2.40	EB	50	10.0	0.9	2.3	A	M	nd	
BONNEVILLE	2W-W														
BONYTHON	1W-Y	1955 - 1963	29	3	156	1.87	S	40	12.0	4.6	5.0	A	G	nd	
BORDER CHIEF	2Y-O	1968 - 1982	26	28	204	2.51	SB	40	10.6	2.2	3.4	A	G	8.8	B
BORDER LEGEND	2Y-O	1973 - 1973	22	21	169	2.00	S	25	12.0	2.6	5.4	A	G	10.8	
BOREAS	2W														
BOSWELL	3W-YYR	1964 - 1973	36	38	172	2.14	SB	49	10.5	1.4	3.6	A	G	8.0	
BOSWIN	1W-Y	1955 - 1981	1	8	243	2.59	E	43	10.0	4.1	4.9	A	M	7.1	B
BOTANY BAY	2Y	1964 - 1967	40	21	216	3.50	SB	44	11.0	2.3	3.3	A	G	7.8	
BOVAGH	2W-YYO	1982 - 1984	52	28	nd	2.57	B	42	10.3	1.9	3.1	A	G	9.0	
BRABAZON	1Y-Y	1975 - 1987	18	-2	164	2.34	S	40	10.0	4.4	4.8	A	M	7.9	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
BRAEMAR	1Y-Y	1964 - 1967	40	31	nd	nd	S	26	11.0	4.4	4.3	A	G	nd	
BRAHMS	2W-O	1973 - 1987	40	38	146	2.20	B	41	10.8	1.8	4.1	A	G	9.9	B
BRAIDWOOD	2Y-Y	1976 - 1981	26	22	200	2.15	S	42	10.2	3.0	4.2	A	G	8.1	
BRANDON	1Y-Y	1955 - 1967	-4	1	222	2.45	S	42	12.0	4.5	4.5	A	M	6.0	
BRANIEL	2Y-O	1955 - 1963	-20	8	216	1.87	SB	45	10.0	2.4	3.6	A	M	nd	
BRAVADO	2Y-O	1968 - 1981	48	38	185	2.14	S	42	10.8	1.8	4.0	A	G	8.0	
BRAVOURE	1W-Y	1980 - 1985	20	20	160	1.74	S	44	12.6	4.8	3.8	A	VG	9.9	
BRAVURA	3W-O	1968 - 1972	43	51	126	0.92	SB	36	10.5	1.1	3.2	A	G	9.2	
BREAD AND CHEESE	2W-Y	1955 - 1963	3	22	309	1.87	S	47	11.0	3.8	4.4	A	G	nd	
BREAGE	2Y-Y														
BREAK OF DAY	2Y-R	1979 - 1979	46	25	502	1.80	S	36	10.0	2.0	3.0	A	G	8.6	
BRIDAL CROWN	4W-Y														
BRIDGET HILL	2Y-O	1964 - 1967	20	29	180	4.20	SB	38	10.0	2.0	3.3	A	G	9.4	
BRIGHT BEAUTY	3W-O	1955 - 1963	-11	23	441	2.40	SB	49	10.0	1.6	3.0	A	M	9.0	
BRIGHT CHARM	3W-O	1964 - 1967	51	40	nd	nd	SB	33	10.5	1.6	4.5	B	G	8.2	
BRIGHT CIRCLE	3W-O	1955 - 1963	-11	26	334	2.93	SB	60	10.0	1.2	3.7	A	G	nd	
BRIGHT LAD	2Y-O	1976 - 1982	7	-6	144	1.88	SB	49	10.8	2.7	3.3	A	G	8.6	
BRIGHT LASS	2Y-YYR	1972 - 1981	7	10	164	2.09	SB	48	9.6	2.0	4.0	A	G	10.5	
BRIGHTLING	2Y-OOR	1955 - 1963	36	31	400	1.33	SB	52	10.0	1.2	3.5	A	M	nd	
BRILLIANCY	3Y-YYO	1955 - 1967	-28	9	304	3.00	SB	52	11.0	1.5	2.6	A	P	9.4	
BRILLIANT STAR	1Y-OOY	1973 - 1973	32	28	181	1.36	SB	37	10.0	2.1	3.6	B	G	9.6	
BRIMSTONE	1Y-Y	1955 - 1963	66	32	187	1.87	S	34	11.0	5.1	5.4	A	G	nd	
BRITISH CHARM	2Y-O	1955 - 1963	-14	5	193	1.87	SB	48	10.0	2.4	3.2	A	M	10.0	
BROOKFIELD	2W-W	1972 - 1972	66	31	100	0.83	S	30	11.0	4.7	3.9	A	G	8.4	S
BROOKVILLE	2W-Y	1964 - 1967	-2	26	nd	nd	SB	35	12.0	2.5	5.1	B	G	8.2	
BROUGHSHANE	1W-W	1955 - 1984	14	34	172	2.00	S	48	12.2	5.2	5.4	A	VG	9.6	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
BRUNSWICK	2W-Y	1955 - 1984	25	11	181	2.04	S	39	10.8	3.4	4.2	A	G	9.3	
BRUSSELS	1W-W	1968 - 1981	54	25	156	1.95	S	38	11.6	4.8	4.6	A	G	8.1	VS
BRUTUS	2Y	1955 - 1963	-11	17	305	2.40	E	44	9.0	2.6	3.5	A	M	nd	
BRYHER	3W-W	1976 - 1981	-1	46	123	2.03	SB	56	10.2	0.9	2.0	A	G	10.2	S
BUCKLER	2W-ORR	1955 - 1963	14	15	234	1.87	SB	58	10.0	1.5	4.1	A	M	nd	
BULLA	1W-Y	1976 - 1977	4	5	122	1.73	S	34	10.2	4.2	4.2	B	VG	9.5	V
BULLION	1Y-Y	1964 - 1977	12	4	128	1.76	S	36	10.0	4.5	5.0	A	G	8.0	
BUNCO	2Y-YR														
BUNCRANA	2W-O	1955 - 1963	29	29	264	1.87	SB	43	10.0	2.8	2.9	A	G	7.8	
BUNDORAN	2W-P														
BURGEMESTER GOUVERNEUR	1Y-Y	1955 - 1984	20	18	161	2.06	S	43	11.9	4.9	5.2	A	G	9.6	B
BURNING HEART	11Y-OOW	1964 - 1967	-16	31	nd	nd	SB	37	10.5	2.7	6.0	A	G	7.8	
BURRATOR	3W-YR	1977 - 1979	-16	36	106	1.69	S	44	11.2	1.3	2.8	A	G	7.7	
BUSHMILLS	3W-YO	1971 - 1987	2	47	151	2.06	B	59	10.1	1.1	2.7	A	G	9.8	
BUSHTIT	6Y-Y														
BUSTER	2Y	1972 - 1973	27	22	152	1.90	S	42	9.2	2.2	3.3	A	G	8.9	
BUTTER AND EGGS	4Y-Y	1968 - 1972	16	28	180	1.94	SB	44	9.3	1.6	nd	A	M	6.9	S
BUTTER KING	1Y-Y	1977 - 1981	13	9	173	1.83	S	35	9.5	3.5	3.6	A	G	11.6	
BUTTERCUP	7Y-Y	1955 - 1963	-7	31	329	2.40	S	47	9.0	3.0	2.4	A	G	11.8	
BUTTERSCOTCH	2Y-Y	1976 - 1987	23	15	153	1.79	S	43	11.5	4.3	4.5	A	M	8.3	
BUZZARD	2W	1964 - 1967	49	37	269	2.10	S	36	9.5	3.4	3.3	A	G	6.0	
C. J. BACKHOUSE	2Y-O	1955 - 1967	24	20	247	2.14	S	47	9.0	1.9	2.0	A	P	8.0	
CAEDMON	9W-YR	1955 - 1982	2	45	188	2.06	SB	48	8.0	0.6	1.8	A	G	9.8	
CAERLEON	2Y-O	1955 - 1963	0	20	291	2.40	SB	58	10.0	2.1	3.4	A	G	nd	
CAIRNDHU	2W-GPP	1984 - 1984	60	45	nd	0.50	nd	nd	nd	nd	nd	nd	nd	nd	
CAIRO	2W-R	1964 - 1967	7	51	nd	nd	SB	33	9.0	1.3	3.2	A	VG	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
CALABAR	2Y-Y	1972 - 1972	60	26	120	1.75	S	35	11.5	3.5	4.6	A	G	9.4	
CALEDONIA	2W-YYO	1955 - 1963	36	30	259	2.40	SB	48	11.0	1.5	4.5	A	M	nd	
CALIFORNIA (15)	2Y-Y	1955 - 1984	-5	5	263	2.86	S	48	10.2	3.0	3.1	A	G	10.2	
CALIFORNIA GOLD	2Y-O	1964 - 1967	-16	15	257	2.20	SB	45	11.0	2.6	3.8	A	G	8.8	
CAMBER	1Y-Y	1968 - 1971	23	27	215	1.40	S	38	11.0	5.2	5.3	A	G	5.1	S
CAMBRIA	1Y-Y														
CAMELLIA	4Y-Y	1955 - 1963	32	29	76	1.33	SB	42	10.0	nd	nd	A	M	nd	
CAMELOT	2Y-Y	1967 - 1967	52	54	245	1.00	SB	39	12.0	2.7	3.3	A	G	nd	
CAMERONIAN	1W-W														
CANARYBIRD	8Y-GOO	1955 - 1967	-45	25	193	2.50	SB	41	5.0	0.6	1.6	A	M	9.4	
CANASTA	11W-Y	1973 - 1981	-23	21	169	1.74	SB	40	10.6	3.0	8.1	A	G	8.3	S
CANDIDA	4W-Y	1973 - 1984	16	26	128	1.66	SB	42	10.2	2.3	nd	A	G	8.3	BS
CANDOUR	2W-W	1964 - 1967	-39	17	174	2.00	S	36	11.5	4.4	4.4	B	G	7.4	
CANOPIUS	1W-W	1964 - 1967	69	34	nd	nd	S	28	10.5	4.6	5.0	A	G	8.2	
CANTABILE	9W-GYR	1955 - 1982	43	57	224	2.24	SB	48	8.0	0.5	1.9	A	G	9.3	
CANTATRICE	1W-W	1955 - 1980	16	19	173	1.74	S	36	12.0	4.8	3.6	A	G	7.6	
CAPE HORN	1W-Y	1972 - 1972	17	31	108	2.62	S	38	11.5	5.2	6.1	A	VG	8.6	
CAPE KENNEDY	11W-O	1973 - 1982	35	43	115	1.51	SB	40	10.6	3.1	5.9	A	G	8.5	
CAPISCO	3W-GYO	1977 - 1987	25	34	221	3.04	S	52	8.9	0.9	1.9	A	M	9.1	
CAPPAROE	3W-R	1973 - 1980	2	37	156	2.15	SB	44	10.2	1.2	3.1	A	VG	8.5	
CARACAS	2O-O	1984 - 1984	35	52	nd	1.80	S	43	9.0	1.1	nd	nd	G	nd	
CARAMEL	9W-O	1977 - 1981	27	5	175	1.80	S	42	10.6	4.4	4.9	A	G	10.4	
CARAVELLE	1Y-Y	1972 - 1973	-12	24	159	1.33	S	36	10.8	4.7	4.0	A	VG	10.1	S
CARBINEER	2Y-O	1955 - 1967	2	20	260	2.28	S	50	10.0	2.0	3.3	B	VG	9.5	
CARDIGAN	2W-Y	1968 - 1973	58	29	167	2.01	S	34	10.5	3.6	3.6	B	G	10.0	
CAREFREE	1W	1968 - 1973	48	34	116	1.59	S	38	11.9	5.0	5.0	A	VG	8.4	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
CAREYSVILLE	2W-Y	1972 - 1981	11	23	117	1.63	S	46	12.2	4.2	4.9	A	VG	9.1	
CARGAN	2Y-Y	1964 - 1967	11	29	nd	nd	S	35	10.5	4.2	4.0	B	VG	7.0	
CARLTON	2Y-Y	1955 - 1987	3	15	243	2.25	S	40	11.8	3.5	5.0	A	G	7.8	BS
CARMINOWE	3W-R	1955 - 1963	-4	40	288	2.40	SB	50	10.0	0.9	2.6	B	M	nd	
CARMOA	2W-YYP	1964 - 1967	11	27	215	2.20	S	31	10.0	2.9	4.2	A	G	9.6	
CARNALEA	2W-WWY	1964 - 1967	10	26	234	2.30	S	36	9.5	1.7	4.1	B	G	10.0	
CARNATION	4Y-O	1964 - 1967	20	25	nd	nd	SB	29	10.0	1.6	nd	A	M	6.6	
CARNBEG	2Y-OOR	1973 - 1982	17	18	187	2.85	SB	36	10.6	2.7	4.0	A	G	11.0	
CARNLOUGH	2W-WWP	1955 - 1963	14	12	215	1.87	S	38	10.0	3.5	3.0	B	VG	nd	
CARNMONEY	2W-W	1964 - 1967	-24	26	280	1.90	S	38	11.0	2.9	3.9	A	G	9.6	
CARNMOON	3W-WWY	1968 - 1971	75	46	202	1.70	SB	41	9.5	1.6	3.4	A	VG	8.6	
CARNSORE	3W-GR	1955 - 1963	-11	37	284	2.40	SB	53	10.0	1.3	2.8	A	G	nd	
CARO NOME	2W-WPP	1977 - 1980	13	24	254	2.05	SB	37	11.1	2.2	4.0	B	VG	9.4	S
CAROLINE FOX	1W-Y	1976 - 1978	26	26	169	2.24	S	34	10.2	4.0	3.5	A	G	10.5	S
CARPATICA	3W-O	1964 - 1967	17	26	166	1.60	SB	42	8.5	1.1	3.1	A	G	9.4	
CARRICK	2W-P	1977 - 1978	34	47	286	1.43	S	31	10.2	3.6	3.4	B	G	10.4	
CARRICKBEG	1Y-Y	1977 - 1982	34	24	204	2.22	S	36	11.3	4.6	3.7	A	G	9.4	B
CARRICKMANNON	2W-GPP	1977 - 1977	75	41	272	2.24	S	26	11.5	3.0	3.0	B	G	11.0	
CARRIGART	3Y-O	1964 - 1967	-21	18	275	2.70	SB	40	10.0	1.4	2.5	A	G	8.0	
CASA D'ORO	1Y-Y	1972 - 1972	17	15	174	1.85	S	32	12.0	5.4	6.5	A	M	6.0	
CASABIANCA	2W-Y	1972 - 1972	8	19	173	3.00	S	41	12.5	3.6	5.8	A	G	8.2	
CASABLANCA	2W-Y	1964 - 1979	-35	28	209	1.32	SB	39	10.5	2.5	3.9	A	G	6.8	
CASSANDRA	9W-YR	1964 - 1967	-32	33	nd	nd	SB	33	9.0	0.4	1.6	B	G	7.6	
CASSATA	11W-W	1971 - 1982	18	10	226	2.07	S	35	10.4	3.2	8.9	A	G	8.7	S
CASTELLA	2W-W	1955 - 1963	-4	19	280	2.40	S	48	10.0	2.3	5.2	A	M	nd	
CASTLE COOLE	2W-WWO														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
CASILE KENNEDY	2W-WWY	1964 - 1967	-50	30	nd	nd	SB	33	10.0	1.8	4.0	B	G	7.6	
CASILE OF MEY	2W-W	1972 - 1972	17	30	122	1.83	S	30	11.0	4.1	4.3	A	G	8.6	S
CASILEBAR	1Y-Y	1964 - 1967	-9	20	nd	nd	S	33	10.5	4.1	4.3	A	G	8.0	
CASILEAREAGH	1Y-Y	1964 - 1973	49	29	172	1.89	S	36	11.5	5.2	4.6	A	VG	8.2	V
CASILEROCK	2Y-R	1968 - 1982	4	21	194	2.60	S	36	10.0	2.1	3.3	A	G	8.5	
CASILEWELLAN	1Y-Y	1972 - 1981	11	40	150	1.87	S	39	10.9	4.5	5.0	A	G	9.0	
CATHAY	2Y-R	1979 - 1985	-4	38	173	2.50	S	40	10.3	2.3	3.3	B	G	9.1	
CATRIONA	2W-YYW	1972 - 1973	38	28	91	1.35	S	31	10.0	2.2	4.2	A	G	8.2	
CAVALIER	3W-YOO	1967 - 1967	56	54	138	0.80	SB	35	8.0	1.1	2.2	A	G	nd	
CAVALIERO	2Y-YOO	1955 - 1973	-16	5	191	2.54	SB	39	10.3	2.4	3.3	A	M	9.5	
CAVODA (16)	1W-GPP	1982 - 1984	67	31	nd	1.88	S	41	nd	nd	nd	A	nd	nd	
CELEBRITY	1W-Y	1955 - 1963	-7	8	178	1.87	S	35	10.0	4.7	4.6	A	M	nd	
CELESTIAL	5Y-Y														
CERVANTES	1Y-Y	1964 - 1967	-5	-6	304	1.60	S	27	10.5	3.9	4.0	A	P	7.8	
CEYLON	2Y-O	1968 - 1981	-2	20	190	2.23	S	34	11.0	2.4	3.0	B	G	10.9	
CHAMPAGNE	2W-P	1955 - 1963	29	22	259	1.60	SB	38	12.0	3.5	5.5	B	VG	nd	
CHANTERELLE	11Y-Y														
CHANTILLY	2W-O	1972 - 1972	53	34	86	1.29	S	31	9.0	2.2	4.8	A	G	7.0	
CHARIOTEER	1Y-Y	1966 - 1968	37	37	180	1.70	S	44	12.5	5.6	4.9	A	VG	6.8	S
CHARIS	2W-P	1964 - 1967	27	27	201	2.70	SB	42	10.5	2.4	3.5	A	G	7.0	
CHARITY MAY	6Y-Y	1964 - 1967	25	15	184	3.80	S	25	9.0	2.6	2.3	A	G	9.8	
CHARLES	3W-YOO	1955 - 1963	7	22	340	1.87	E	46	9.0	0.9	2.6	A	M	nd	
CHARTER	2Y-W	1977 - 1987	22	19	205	3.30	E	40	10.8	3.4	3.3	A	G	14.0	
CHATSWORTH	1W-Y	1955 - 1963	10	15	252	1.87	S	38	10.0	3.6	4.0	A	G	nd	
CHEERFULNESS	4W-Y	1955 - 1987	26	40	262	2.36	E	45	5.8	0.5	1.5	A	G	10.2	
CHEERIO	2Y-O	1955 - 1963	-4	15	187	1.87	S	48	9.0	1.8	3.7	B	VG	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
CHELSEA CHINA	2W-GPP	1964 - 1967	55	33	nd	nd	SB	37	9.5	2.2	3.3	B	G	9.4	
CHELSEA DERBY	2W-P	1976 - 1981	2	37	134	2.21	S	38	9.2	2.6	3.0	B	G	10.1	B
CHEMAWA	2Y-OOY	1976 - 1981	-3	16	149	2.18	S	46	8.5	2.2	2.8	A	G	10.8	
CHERIE	7W-P	1955 - 1967	-4	32	248	2.50	E	47	8.0	2.0	2.1	B	M	7.8	
CHEVALIER	1Y-Y	1977 - 1977	50	18	433	1.17	S	34	10.5	4.1	3.2	B	VG	10.2	V
CHIEF JUSTICE	1W-Y	1964 - 1967	56	40	nd	nd	S	30	10.5	4.9	5.3	A	VG	8.0	
CHIFFON	2W-P	1972 - 1981	63	38	139	2.27	S	40	9.6	2.1	3.5	A	G	8.0	B
CHILLAGOE	2W-W	1976 - 1987	-1	7	99	1.40	S	39	12.7	4.8	5.3	A	M	11.1	B
CHILLION	1Y-Y	1974 - 1974	50	13	309	1.75	S	39	11.5	4.9	4.5	A	VG	9.4	V
CHILTERN	1W-Y	1972 - 1973	28	8	102	1.19	S	38	11.5	4.7	4.8	A	G	9.9	
CHINA PINK	2W-P														
CHINESE WHITE	3W-W	1968 - 1971	60	46	192	1.90	SB	44	10.5	1.4	3.7	B	G	8.1	
CHINITA	8Y-YYR	1955 - 1963	-20	29	265	1.87	SB	58	7.0	0.6	2.1	A	M	nd	
CHINOOK	2W-Y	1968 - 1987	36	24	185	2.22	S	42	11.0	3.1	5.4	A	G	8.0	BS
CHITRA	2Y-R	1974 - 1974	37	26	340	0.71	S	42	9.5	2.0	2.5	A	G	9.0	V
CHMOY	2W-Y	1972 - 1975	10	16	135	2.04	S	39	11.0	5.0	4.4	A	G	7.8	
CHOIR BOY	2W-O	1977 - 1980	5	13	323	1.96	SB	40	9.8	2.4	3.5	A	G	9.1	
CHUNGKING	3Y-O	1955 - 1973	20	23	301	1.86	SB	46	10.0	1.4	2.9	A	VG	7.6	
CHYNANCE	2W	1972 - 1975	34	32	130	2.35	S	35	10.5	2.0	3.3	A	G	8.6	
CIBOLA	2Y-Y	1968 - 1987	3	-6	175	2.01	S	37	11.1	3.7	5.1	A	G	8.3	B
CICELY	2W-W	1964 - 1967	-51	9	nd	nd	S	31	9.5	3.3	2.8	B	G	7.8	
CIMAROSA	1W-Y														B
CINEL	2Y-R	1976 - 1980	4	20	193	2.00	SB	50	9.5	1.6	3.8	A	G	9.9	V
CINZANO	2W-O														
CITRIX	1Y-Y	1964 - 1967	5	14	nd	nd	S	29	11.5	4.8	5.5	B	M	6.0	
CLACKAMAS	2W-YYO														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
CLACKRATTLE (17)	2Y-O	1955 - 1963	-21	22	480	1.87	E	46	9.0	1.8	3.0	A	G	nd	
CLAMOR	2W-YYR	1955 - 1963	43	20	401	2.40	SB	42	9.0	1.5	3.0	A	G	nd	
CLAVA	2W-Y	1955 - 1963	-7	23	210	1.87	SB	53	12.0	2.0	4.3	A	G	nd	
CLEENA	2W-OOY	1964 - 1967	33	12	191	2.40	S	33	9.5	2.9	3.9	A	G	7.0	
CLEMENTINE	2W-Y	1973 - 1982	27	17	145	2.58	S	38	10.2	3.1	4.0	A	VG	8.8	S
CLOCKFACE	3W-YYO	1968 - 1973	28	50	113	2.01	SB	46	11.0	1.0	3.2	A	G	9.9	
CLOGHEEN	3W-Y	1976 - 1977	5	45	115	2.21	SB	34	11.0	1.3	3.6	B	G	8.2	
CLONCARRIG	1Y-Y	1968 - 1981	26	34	180	2.06	S	41	10.5	4.1	5.0	A	VG	8.2	VS
CLONEEN	2W-W	1973 - 1977	38	39	126	2.58	SB	48	10.7	2.3	3.2	B	G	7.9	S
CLONMORE	1Y-Y	1964 - 1967	47	30	nd	nd	S	30	11.0	4.8	4.9	A	G	6.6	
CLOUDCAP	2W-WWP	1976 - 1981	1	36	153	1.91	SB	49	10.8	2.5	4.7	B	G	10.6	S
COLORANGE	1Y-O	1976 - 1980	38	28	144	1.78	SB	35	10.2	2.6	7.1	B	G	10.3	
COLORATURA	3W-WWO	1976 - 1987	18	34	149	1.98	S	40	10.2	1.6	4.1	A	M	9.4	
COLUMBINE	3W-WWO	1964 - 1967	51	57	290	1.60	S	45	8.0	0.5	1.9	A	M	10.0	
COMMODORE	2W-O														B
COMPRESSUS (16)	8W-Y	1955 - 1967	-27	22	174	2.15	S	34	4.0	0.6	1.6	B	M	10.8	
COMPTON MACKENZIE	2W-P														
CONBEG	2W-R														
CONFUOCO	2Y-R														
CONFUSION	2W-P	1976 - 1978	24	10	291	1.64	S	39	8.8	2.7	3.0	B	G	8.5	
CONGO	2Y-O	1972 - 1981	26	13	164	2.59	S	39	10.4	1.7	3.7	A	G	8.8	
CONSPICUOUS (16)	3Y-YYO	1955 - 1967	-3	28	319	2.65	E	49	10.0	1.3	2.2	B	M	10.0	
CONTENT	1W-WWY	1955 - 1967	-14	10	194	2.18	SB	43	11.0	4.9	3.8	A	G	8.8	B
CONTRAPUNT	2Y-O	1973 - 1981	11	19	147	2.37	S	41	10.0	2.1	2.6	A	G	10.3	B
CONVAIR	1Y-Y	1976 - 1977	36	2	112	1.42	S	35	10.8	4.8	3.8	A	G	8.7	
COOL CONTRAST	1W-Y	1977 - 1977	43	26	240	1.74	S	32	11.0	4.9	4.0	A	VG	10.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
COOLATTIN	2W-Y	1976 - 1982	30	35	146	2.17	S	39	10.2	2.6	2.9	A	G	10.9	S
COPPELIA	2W-P	1981 - 1981	78	41	275	1.00	SB	31	10.0	2.2	3.1	nd	G	nd	B
COPPER BOWL	2Y-O	1955 - 1963	-7	23	244	1.87	SB	49	11.0	2.4	3.7	A	M	nd	
COPPERSMITH	2Y-O	1968 - 1971	54	32	221	2.00	SB	44	11.0	2.5	4.0	B	G	7.4	
CORAL DAWN	2W-Y														
CORAL ISLAND	2W-GPP	1968 - 1980	37	22	129	2.50	SB	45	11.0	2.3	3.7	A	G	7.8	B
CORBY	2W-W	1964 - 1967	56	43	190	2.90	S	33	10.0	3.1	3.3	B	G	8.2	
CORDIAL	2W-P	1976 - 1980	30	32	144	2.44	S	36	10.4	3.2	4.0	B	G	10.4	B
CORINTHIAN	1Y-Y	1955 - 1973	4	16	209	1.51	S	42	10.0	4.8	5.1	B	G	7.6	
CORNCRAKE	3W-O	1968 - 1973	58	58	164	1.69	SB	42	9.2	1.0	2.8	A	VG	10.5	
CORNET	6Y-Y	1967 - 1970	5	-11	241	3.64	S	25	9.2	3.9	3.0	A	G	8.2	
CORNISH FIRE	2Y-O	1955 - 1967	-5	15	274	2.24	S	42	10.0	1.8	2.9	A	P	8.6	B
COROFIN	3W-YR	1968 - 1972	67	38	147	1.22	SB	40	8.8	1.1	3.0	A	G	8.9	
CORONACH	3W-R	1968 - 1972	4	44	118	1.08	S	44	9.0	0.9	2.5	A	VG	9.0	
CORONADO	2Y-P	1970 - 1970	40	34	126	0.90	S	27	10.5	3.1	4.6	B	G	nd	
CORREGIO	2W-O	1964 - 1967	5	41	nd	nd	SB	40	10.5	2.1	3.1	B	G	10.0	
CORRYMBELA	2W-Y														
CORTINA	1W-W	1964 - 1967	34	26	nd	nd	S	28	11.0	5.0	6.0	B	G	8.4	
COTOPAXI	2Y-O	1955 - 1963	-14	12	445	1.87	S	47	10.0	1.6	3.3	A	G	nd	
COTTERTON	2W-W														B
COTTESMORE	2Y-YOO	1955 - 1963	-7	29	343	1.87	SB	52	10.0	1.7	3.2	A	M	nd	
COUNSELLOR	1Y-Y	1955 - 1973	14	35	167	0.94	S	38	12.1	5.2	4.6	A	G	9.1	V
COUNTRESS OF STAIR	1W-Y	1964 - 1967	13	29	nd	nd	S	32	11.0	4.8	4.7	A	G	nd	
COURAGE	2W-W														B
COURT MARTIAL	2Y-O	1968 - 1984	22	6	196	2.73	S	40	10.4	2.5	3.3	A	VG	10.5	
COVE	2Y-Y	1968 - 1987	-7	-8	167	2.15	E	44	12.3	3.4	5.2	nd	P	8.3	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
COVENT GARDEN	1Y-Y	1955 - 1967	11	8	212	1.76	E	41	10.0	4.1	4.4	A	M	7.3	
COVERACK GLORY	2Y-Y	1955 - 1967	-1	14	343	2.60	S	48	10.0	3.0	3.6	A	M	9.7	
COVERACK PERFECTION (19)	2W-YYO	1955 - 1963	7	15	223	1.87	SB	47	11.0	2.3	5.0	A	G	nd	
COVERACK PRIDE	1Y-Y	1968 - 1979	39	4	176	2.21	S	40	11.0	5.1	5.9	A	G	8.9	
CRAFORD	8W-O	1955 - 1963	-14	15	201	1.33	S	34	5.0	0.6	1.5	A	M	5.4	
CRAIGYWARREN	2Y-R	1955 - 1982	2	28	248	2.53	SB	54	11.0	1.9	3.3	A	VG	9.8	
CRASKIN	1Y-Y	1955 - 1963	0	6	238	2.40	S	42	11.0	4.9	5.0	A	G	nd	
CREAM CUP	2W-Y														
CREAMED HONEY	1Y-W	1977 - 1978	43	10	302	2.07	S	40	10.8	4.6	4.6	A	M	10.1	
CREBILLY	2W-WWY	1964 - 1967	-2	26	185	2.30	S	32	10.5	2.0	3.7	B	G	7.0	
CRENVER	3W-YYO	1955 - 1980	27	42	244	2.10	S	57	9.0	0.7	2.6	A	VG	8.0	
CREPELLO	3W-GWY	1972 - 1975	18	44	180	1.62	S	41	10.1	1.0	2.2	A	VG	9.6	
CRESCENDO	2Y-YYO	1955 - 1980	-11	14	308	2.69	SB	58	10.0	2.4	2.7	A	VG	9.5	
CRESCENT	2Y-R	1955 - 1963	-14	15	441	1.87	E	50	10.0	2.4	3.6	A	M	nd	
CRINOLINE	2Y														
CRITERION	2W-OOY	1964 - 1973	4	33	202	1.92	SB	38	11.5	2.0	4.6	A	G	9.8	
CROCK OF GOLD	1Y-Y	1972 - 1973	23	10	172	2.23	S	34	11.2	4.6	5.2	B	VG	6.0	
CROCUS	2Y-Y	1955 - 1963	-7	18	195	1.87	E	47	11.0	4.0	5.0	A	VG	9.6	
CROESUS	2Y-YYO	1955 - 1967	22	24	335	2.78	SB	46	9.0	1.2	3.0	A	P	9.6	
CROMARTY	1Y-Y	1955 - 1976	22	25	169	2.15	E	38	11.0	4.5	4.3	A	G	6.7	
CROWN DERBY	3W-YYO	1964 - 1967	51	39	185	1.60	SB	38	11.0	1.7	3.4	A	G	9.0	
CRUMLIN	1Y-Y	1973 - 1978	52	32	84	2.13	S	31	10.5	4.8	4.3	A	VG	10.0	S
CRUSADER	2W-O	1972 - 1975	36	26	209	2.03	SB	36	11.5	2.1	4.8	A	G	9.0	
CRYSTAL QUEEN	2W-W	1955 - 1963	0	17	369	2.40	S	48	11.0	3.5	3.8	B	P	nd	
CUSHENDALL	3W-GWW	1955 - 1967	35	54	166	2.04	SB	46	8.0	0.6	1.9	A	G	8.0	
CUTTY SARK	7Y-Y	1955 - 1967	-7	21	204	1.45	SB	44	8.0	1.1	1.9	A	P	9.4	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
CYCLADES	6Y-Y														
CYCLONE	6Y-Y														
CYCLOPE	1Y-Y	1984 - 1985	14	5	170	2.50	S	31	12.5	5.1	6.8	A	VG	7.3	
CYNOSURE	2W-O	1955 - 1963	21	24	336	1.60	SB	48	10.0	1.7	2.2	A	P	nd	
DACTYL	9W-GYR	1955 - 1967	24	56	161	1.84	SB	52	8.0	0.5	1.9	A	VG	7.2	
DAISY SCHAFER	2W-WYY	1955 - 1963	7	17	124	1.87	S	42	10.0	3.9	4.7	A	M	nd	
DALBOYNE	2Y-O	1964 - 1980	-1	10	246	2.80	S	36	10.0	2.4	2.4	B	G	8.2	
DALLAS (16)	3W-GWW	1968 - 1982	62	58	166	2.32	SB	48	9.0	0.7	2.1	A	G	10.6	
DAMASCUS	3W-R	1955 - 1963	-7	20	325	1.87	SB	48	10.0	1.0	2.7	B	M	nd	
DAMSON	2W-O	1955 - 1967	-16	16	300	2.22	SB	57	10.0	2.2	3.1	A	G	8.4	
DANAE	2Y-O	1984 - 1985	10	19	nd	1.10	S	37	10.8	2.8	3.7	A	G	9.8	
DANCING PARTNER	2W-P	1977 - 1978	58	34	298	2.27	S	32	10.8	3.6	3.6	A	G	11.5	
DAPHNE	4W-W	1955 - 1967	27	35	238	2.02	E	48	7.0	nd	nd	A	G	9.6	
DARLINGTON	2Y-O	1972 - 1972	17	19	296	1.15	S	38	11.0	2.8	3.9	A	G	10.2	
DAVID SPRY	1Y-Y	1976 - 1982	12	4	176	1.98	S	43	12.0	4.8	5.0	A	VG	9.3	
DAVIOT	2W-OOY	1964 - 1967	7	26	nd	nd	S	35	10.0	2.3	2.5	A	G	10.0	
DAWLEY	2Y-Y	1972 - 1977	15	18	174	1.84	S	37	10.0	3.3	3.8	A	G	11.0	
DAWN	5W-Y	1964 - 1967	18	31	192	2.20	SB	33	8.0	1.0	2.3	A	M	8.7	
DAWSON CITY	1Y-Y	1955 - 1967	19	22	211	1.94	S	38	10.0	4.3	3.9	A	M	6.9	
DAYDREAM	2Y-W	1977 - 1987	30	27	193	3.13	E	36	9.5	3.5	3.8	A	M	11.4	B
DEAN	2W-YOO	1955 - 1980	9	13	222	2.42	S	44	11.0	2.4	3.7	A	VG	8.0	B
DEANNA DURBIN	2W-O	1955 - 1967	28	26	282	2.24	SB	52	10.0	2.1	3.5	A	M	7.2	
DEBUTANTE	2W-P	1977 - 1978	72	33	400	1.06	S	24	10.8	2.8	4.1	B	VG	9.2	
DECENCY	1Y-Y	1955 - 1973	10	28	187	1.79	S	41	10.0	4.2	4.6	A	G	7.7	
DECOR	2Y-YOO	1955 - 1963	14	8	243	1.87	S	36	12.0	3.1	4.6	A	G	nd	
DEDERANG	1W-Y	1973 - 1975	-2	10	230	1.18	S	34	11.2	4.0	4.2	A	M	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
DEERFIN	2Y-Y	1977 - 1978	40	24	258	1.91	S	36	10.8	3.5	3.2	B	M	10.6	
DEFIANCE	2Y-O	1973 - 1981	27	34	178	2.42	SB	47	10.1	1.6	3.1	A	G	9.7	
DEIRDRE	3W	1968 - 1982	24	49	260	2.55	SB	46	11.0	1.4	3.9	B	VG	9.4	
DELIBES	2Y-O	1964 - 1967	34	16	192	2.00	SB	36	10.5	2.2	5.1	A	G	9.0	
DENHAM	2Y-O	1972 - 1981	9	17	91	1.66	SB	37	9.9	2.5	3.8	A	G	10.2	
DEODORA	2W-WWY	1970 - 1987	16	26	132	1.93	B	44	11.4	2.1	4.6	A	G	9.8	BS
DERVISH	2Y-O	1964 - 1967	24	23	184	2.20	SB	45	10.0	2.0	2.9	A	G	9.6	
DEDEMONA	2W-W	1976 - 1980	9	33	131	2.49	S	38	11.5	4.4	3.9	A	G	10.8	S
DEVILRY	2Y-O														
DEVON LOCH	1W-W														
DEW-POND	2W-W														
DICK	1W	1964 - 1967	23	16	nd	nd	S	33	12.0	5.1	4.4	B	G	7.6	
DICK WELLBAND	2W-O	1955 - 1967	14	32	247	2.04	SB	57	11.0	2.2	4.7	A	M	9.4	
DILIGENCE	1Y-Y	1972 - 1972	33	21	124	0.69	nd	nd	nd	nd	nd	nd	nd	nd	
DILLENBURG	2Y-R	1964 - 1967	20	29	nd	nd	SB	28	10.5	2.6	3.9	B	G	8.8	
DINKIE (17)	3Y-GYR	1964 - 1967	51	45	180	2.20	S	44	7.5	1.2	2.6	A	G	9.2	
DINTON GIANT	1W-Y	1968 - 1972	70	33	112	1.47	S	32	10.8	4.7	5.4	A	G	9.0	BS
DIOLITE	2Y-YYO	1964 - 1967	20	18	181	2.40	SB	42	11.5	2.0	3.2	A	G	9.0	
DIOTIMA	1Y-Y	1955 - 1963	14	35	282	2.13	E	48	7.0	nd	nd	A	G	nd	
DIPLOMAT	2W-Y	1955 - 1967	20	6	265	1.88	S	44	10.2	3.3	3.8	A	G	8.5	
DIVOT	2W-Y	1955 - 1963	14	5	281	1.87	S	46	11.0	3.4	4.5	A	G	10.0	
DOCTOR JOHANNES BEIJER	2W-O	1964 - 1967	-21	26	nd	nd	SB	38	10.5	1.2	3.5	A	G	8.8	
DOCTOR PERON	2W-GOO	1955 - 1963	14	23	321	1.87	S	50	10.0	1.7	3.6	A	G	nd	
DOE ROSS	1Y-Y	1976 - 1987	18	0	143	2.38	S	43	11.4	4.9	5.0	A	M	8.5	B
DOLLY MOLLINGER	11W-OWO														
DOMINATOR	1Y-Y	1955 - 1982	1	-1	141	1.72	E	37	12.0	5.1	5.3	A	G	7.4	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
DONIZETTI	2W-Y														
DONNA BELLA	11Y-OOY	1973 - 1977	38	49	122	1.92	SB	38	10.2	1.7	4.0	A	G	8.5	
DONORE	1Y-Y	1968 - 1973	95	38	186	1.94	S	30	11.0	4.7	5.1	A	G	8.4	
DOUBLE EAGLE	4W-O	1976 - 1980	11	28	139	1.50	SB	38	10.1	1.3	nd	B	G	8.1	
DOUBLE EVENT	4W-Y	1968 - 1982	26	46	150	2.47	S	44	10.0	1.4	nd	A	VG	8.2	S
DOUBLE FORTUNE	4Y-Y														
DOUBLE MING	4W-W	1964 - 1967	3	51	159	2.00	S	49	7.0	nd	nd	A	M	8.2	
DOUBLEDALE	4W-Y	1974 - 1985	26	20	163	2.00	S	35	10.5	3.1	nd	A	G	7.7	VS
DOUBLET	4W-Y	1977 - 1984	40	56	153	2.53	S	52	9.0	0.5	nd	A	M	11.0	
DOUBTFUL	3Y-R	1968 - 1976	27	21	162	1.90	S	46	10.9	1.6	2.7	A	G	9.6	
DOUMER	1Y-Y	1955 - 1963	14	23	187	1.87	S	38	11.0	5.2	5.6	B	G	nd	
DOVE WINGS	6W-Y														
DOWNHILL	3W-W														
DOWNPATRICK	1W-Y	1979 - 1985	15	35	153	2.27	S	40	12.7	5.0	4.9	A	VG	10.6	S
DRAGOMAN	3W-R	1964 - 1967	40	45	nd	nd	SB	39	10.0	1.0	3.2	A	VG	10.0	
DRAGOON	3W-R	1955 - 1963	14	37	369	1.87	SB	56	10.0	1.4	3.1	A	P	nd	
DREAM GIRL	2W-YYO														
DREAMLIGHT	3W-GWR	1955 - 1963	21	46	369	1.87	SB	48	8.0	0.6	2.2	A	G	nd	
DREEN	2W-P	1964 - 1967	51	31	224	3.30	SB	32	10.0	3.2	4.0	A	G	9.0	
DRENAGH	2W-P	1977 - 1981	74	39	252	2.79	S	33	10.5	3.1	3.7	B	VG	13.5	
DROMONA	2W-Y	1984 - 1984	28	40	nd	1.33	S	41	10.5	2.8	3.8	nd	VG	nd	
DROMORE	2W-Y														B
DRUMBOE	2W-WWP	1982 - 1984	63	36	nd	1.60	S	37	10.5	2.9	4.3	A	VG	9.0	
DUKE OF ANJOU	1W-Y	1955 - 1963	29	17	238	1.87	S	43	10.0	4.7	5.0	B	M	nd	
DUKE OF BEDFORD	1W-Y	1955 - 1963	29	12	52	0.80	S	31	11.0	4.8	4.4	A	VG	nd	
DUKE OF WINDSOR (15)	2W-OOY	1968 - 1982	36	32	176	2.90	SB	44	11.3	2.3	4.8	A	G	8.4	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
DULCIMER	9W-GYR	1955 - 1982	8	74	160	1.80	SB	57	8.0	0.4	1.6	A	G	9.8	
DUNCAN	2Y-Y	1968 - 1980	-2	18	222	2.66	S	40	10.8	3.2	3.8	B	G	10.6	S
DUNDARAVE	2Y-R	1982 - 1984	64	35	nd	2.10	B	33	10.0	1.8	3.3	A	G	nd	
DUNDRUM	1Y-Y	1968 - 1977	43	12	142	1.71	S	34	11.5	4.9	5.2	B	VG	8.3	B
DUNEBA	2Y-R	1974 - 1976	20	15	232	0.87	SB	38	10.8	2.2	3.4	A	G	9.1	V
DUNFANE	2W-W	1964 - 1967	-5	17	nd	nd	S	34	11.0	4.1	4.2	A	VG	9.4	
DUNGANNON	1Y-Y	1968 - 1971	69	29	205	2.70	S	38	11.5	4.8	4.8	A	G	6.4	S
DUNGIVEN	1Y-Y	1968 - 1971	15	25	167	2.20	S	42	11.5	4.6	5.0	A	VG	6.7	S
DUNKELD	2Y-O	1955 - 1973	0	23	262	2.12	SB	44	10.0	1.7	4.0	A	G	8.4	
DUNLEWEY	2W-W	1955 - 1963	29	22	198	1.87	S	30	12.0	3.2	3.8	B	G	nd	
DUNLOY	2W-P	1964 - 1973	54	36	134	2.04	SB	39	10.0	2.5	3.3	A	G	8.2	
DUNLUCE	1W-W	1955 - 1963	7	23	187	1.87	SB	43	12.0	4.9	4.0	A	G	nd	
DUNMINNING	2Y-R														
DUNMURRY (15)	1/2W-Y	1972 - 1975	61	32	208	1.85	S	33	11.3	4.2	4.8	A	VG	10.5	
DUNSEVERICK	2W-W														V
DUTCH MASTER	1Y-Y	1955 - 1987	12	-3	198	1.98	E	40	11.9	5.1	5.2	A	G	7.6	
EARLY BIRD	3Y	1971 - 1987	-13	7	160	2.39	B	51	10.1	1.9	3.1	A	M	8.1	
EARLY BRIDE (19)	2W-YOO	1969 - 1987	19	9	156	2.47	E	38	11.0	2.6	4.7	A	M	7.8	S
EARLY GLORY	1Y-Y	1955 - 1963	14	0	270	2.40	E	42	10.0	4.5	4.0	A	M	nd	
EARLY GOLD	1Y-Y	1972 - 1977	8	9	154	1.59	S	43	10.2	3.8	4.6	A	G	7.4	S
EARLY HIGHNESS	1W-Y	1983 - 1983	-7	5	nd	1.50	S	32	10.5	4.0	4.8	nd	G	nd	
EARLY LIGHT	2Y-O	1968 - 1977	22	1	207	1.70	S	37	9.0	2.7	3.3	A	G	7.5	S
EARLY MIST	2W-W	1972 - 1980	18	16	129	1.36	SB	41	11.8	4.5	4.0	A	G	9.8	BS
EARLY PERFECTION	8W-Y	1964 - 1969	-27	4	215	2.08	S	26	4.9	0.6	1.6	A	M	8.8	
EARLY PRINCE	1W-Y	1972 - 1977	11	8	85	1.35	S	36	12.0	4.8	5.2	A	G	10.5	S
EARLY SENSATION	1Y-Y	1964 - 1984	11	5	210	2.13	S	41	11.2	5.0	5.2	A	G	10.2	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
EARLY SPLENDOUR	8W-O	1968 - 1982	-11	28	163	2.95	SB	37	7.0	0.7	1.8	A	G	8.7	B
EARLY SURPRISE	3W-R	1964 - 1967	55	27	nd	nd	SB	25	8.0	1.1	2.0	B	P	7.6	
EASTER BONNET	2W-P														
EASTER BRIDE	1W-W	1964 - 1967	20	40	nd	nd	S	28	10.0	4.2	4.4	B	VG	8.6	
EASTER JOY	1Y-Y	1968 - 1973	72	26	196	1.30	S	32	11.1	4.7	5.2	A	G	7.6	S
EASTER MOON	2W-GWW	1971 - 1987	38	32	147	2.18	B	47	10.4	2.2	2.9	A	G	8.4	S
EASTER MORN	2W-W	1964 - 1967	47	24	171	2.20	S	32	10.5	4.3	4.9	A	G	7.8	
ECHO	2W														
EDDY CANZONY	2W-YYO	1964 - 1982	34	20	214	2.54	SB	33	11.0	2.1	4.7	A	G	8.6	
EDENBANK	3W-R	1976 - 1978	-8	29	134	1.96	SB	42	9.0	1.0	2.7	A	G	9.1	
EDWARD BUXTON	3Y-OOR	1955 - 1987	1	16	225	3.45	SB	48	10.4	1.6	3.0	A	G	8.9	S
EFFECTIVE	1W-Y	1964 - 1977	13	29	161	2.48	S	40	10.2	3.8	3.5	B	G	10.0	B
EGGS AND BACON	4W-O	1964 - 1968	30	32	220	2.00	nd	nd	nd	nd	nd	nd	nd	nd	
EL PASO	2Y-O	1955 - 1963	-7	7	264	1.33	S	45	10.0	2.8	2.9	A	G	nd	
ELABORATE	2Y-O	1977 - 1977	7	2	208	2.12	SB	37	11.0	2.5	3.3	A	G	9.0	
ELEANOR MAY	4W-R	1974 - 1974	24	37	464	0.43	SB	49	nd	nd	nd	A	nd	nd	V
ELF	2Y-WWP														
ELGIN	1Y-Y	1955 - 1963	0	4	126	1.87	E	36	11.0	4.5	4.7	A	G	nd	
ELIZABETH BANDON	2W-R	1955 - 1963	-14	26	264	1.87	SB	51	11.0	2.1	3.2	A	M	nd	
ELLESMERE	2Y-YOO	1955 - 1963	-11	34	399	2.40	SB	48	10.0	1.7	4.1	A	M	nd	
ELMWOOD	2Y-R	1972 - 1978	9	31	157	2.45	SB	43	10.5	2.4	4.4	A	G	9.8	
ELSPETH	3W-YYO	1955 - 1963	-14	21	392	2.40	S	47	10.0	1.3	2.8	B	M	nd	
ELVIRA	8W-YYO	1955 - 1967	4	28	318	2.35	E	46	6.0	0.6	1.6	A	M	9.6	
EMBER	2Y-O	1972 - 1973	-1	26	136	1.96	S	45	12.0	2.8	3.8	A	G	11.4	S
EMPEROR	1Y-Y	1955 - 1967	22	23	191	2.50	S	41	9.0	4.0	3.8	A	P	7.4	
EMPIRE	2W-WYY	1955 - 1963	32	22	348	1.87	S	38	12.0	3.6	4.2	B	M	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
EMPRESS	1W-Y	1955 - 1967	-4	24	173	2.34	SB	38	10.0	4.3	3.4	B	M	7.0	
EMPRESS OF IRELAND	1W-W	1973 - 1981	21	22	110	1.76	S	38	12.1	4.7	5.1	A	VG	12.2	BS
ENCHANTRESS	2W-P	1979 - 1981	89	42	198	1.80	S	30	10.0	3.1	4.2	B	G	9.9	
ENGADINE	3W-W	1964 - 1967	49	41	nd	nd	SB	30	8.5	1.2	3.1	B	G	8.6	
ENNISKILLEN	3W-R	1968 - 1982	26	38	212	2.46	SB	44	10.5	1.1	3.1	A	VG	9.7	
ENTERPRISE	4Y-O	1976 - 1984	18	26	137	2.15	SB	43	11.1	2.8	nd	B	G	8.1	S
ENTRANCEMENT	1Y-W	1973 - 1987	33	8	143	2.47	S	39	10.7	4.5	3.9	A	G	8.4	
EPIC	9W-YOR	1955 - 1963	27	39	252	1.87	SB	40	8.0	0.5	1.9	A	G	nd	
ERIE	2Y-O	1955 - 1963	-7	15	322	2.40	S	50	10.0	1.8	3.5	A	M	nd	
ERLICHEER	4W-Y														
ERNANI	2W-YYO														B
ESKIMO	2W-W	1955 - 1963	21	23	282	1.87	S	44	10.0	3.5	3.6	B	M	nd	
ESTELLA DE MOL	1W-Y														
ETHEL	3W-Y	1976 - 1976	56	23	72	0.90	SB	35	8.5	1.3	2.9	A	G	7.6	V
EVANGELINE	3W-Y	1955 - 1963	-4	22	295	2.40	SB	52	10.0	1.2	2.2	A	P	nd	
EVE DALY	1W-Y	1977 - 1977	21	15	246	1.35	SB	34	11.0	5.2	4.6	A	G	12.6	V
EVENING	2W-W	1964 - 1967	37	33	192	2.40	S	30	10.0	3.3	3.4	B	G	9.6	B
EVERTON	2Y	1972 - 1973	52	23	103	1.90	S	35	10.5	1.8	3.8	A	VG	7.8	
EVITA PERON	2W-O	1972 - 1972	60	34	88	1.50	S	27	9.0	2.5	3.2	B	G	9.2	
EVOLUTION	11Y-Y	1964 - 1967	26	25	232	2.00	S	29	11.5	3.6	9.5	A	M	7.2	
EXPLORER	1Y-Y	1972 - 1972	31	14	163	2.00	S	32	11.0	4.5	5.4	A	G	8.0	
EXPRESSIVE	1Y-Y														
EXTOL	4Y-R	1976 - 1980	25	34	214	1.70	SB	40	9.9	1.7	nd	B	G	9.2	
E. E. MORBEY	2Y-O	1976 - 1981	0	12	145	2.22	S	47	10.4	2.6	4.0	A	G	12.5	S
FAIR COLLEEN	3W-OOY	1968 - 1971	32	46	140	1.30	SB	41	9.5	1.2	3.4	B	G	7.8	B
FAIR GREEN	3W-GYO	1973 - 1979	6	50	138	2.31	SB	59	10.3	0.9	2.3	A	VG	9.6	V

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
FAIR MAIDEN	3W-YYO	1955 - 1963	17	24	714	2.13	S	44	9.0	1.1	2.4	A	P	nd	
FAIR TRIAL	1W-Y														
FAIRLIGHT	1Y-Y	1976 - 1982	12	3	154	2.02	S	40	11.1	4.8	4.4	A	G	10.9	
FAIRMILE	3W-GYO	1982 - 1984	68	49	nd	1.50	B	41	9.5	0.9	2.7	A	VG	nd	
FAIRNESS	4W-O	1955 - 1963	-14	30	60	0.80	SB	52	6.0	0.6	nd	A	M	nd	
FAIRY DREAM	1W-W														
FAIRY KING	2Y-O	1955 - 1963	7	12	315	1.87	SB	54	10.0	2.1	3.4	A	M	8.2	
FAIRY TALE	3W-YYO	1968 - 1976	40	46	177	2.07	S	44	9.5	1.2	2.9	B	G	7.8	
FAITHFUL	2Y-Y	1955 - 1980	-4	13	193	3.18	E	44	10.0	3.8	4.1	A	VG	8.0	
FALAISE	4W-O	1955 - 1967	20	53	262	1.10	SB	55	10.0	1.3	nd	A	G	8.0	
FALSTAFF	2Y-R	1979 - 1981	6	18	430	1.93	S	33	9.5	2.2	2.6	B	G	11.9	
FAMILLE ROSE	2W	1977 - 1977	67	21	413	0.71	SB	28	10.0	3.0	4.8	A	M	9.6	
FANCHEON	2W-YOR	1955 - 1963	0	26	317	1.87	S	44	12.0	2.0	4.4	A	M	nd	
FANCY FREE	2W-YYP														
FANTEE	1Y-Y	1973 - 1973	19	4	255	1.67	S	36	11.5	4.6	5.4	B	G	nd	
FAREWELL (18)	2W-Y	1964 - 1967	47	24	184	2.70	SB	35	12.0	2.9	3.7	B	VG	8.0	
FASHION	11Y-YYO	1979 - 1984	38	26	157	1.46	S	40	10.5	3.5	5.7	nd	G	8.1	
FASTNET	2W-YYR	1964 - 1967	51	32	265	3.00	S	33	10.5	2.1	4.2	B	G	7.8	
FAVOURITE	2W-Y	1977 - 1978	42	45	334	2.53	S	41	10.8	2.3	3.8	A	G	10.1	
FAWNGLO	2Y-Y														
FAY MARK	2W-O	1964 - 1967	13	26	nd	nd	SB	36	9.5	1.7	4.4	A	G	8.4	
FEBRUARY BICOLOR	2W-Y	1977 - 1981	-5	-8	159	1.81	SB	28	11.3	4.6	5.2	B	VG	nd	
FEBRUARY GOLD	6Y-Y	1955 - 1967	7	-10	198	2.18	S	25	8.0	2.9	2.4	B	M	9.0	
FEBRUARY SILVER	6W-W														
FEDORA	2W	1964 - 1967	53	39	137	1.80	S	35	10.0	1.9	4.3	B	G	8.4	
FELINDRE	9W-GYR	1968 - 1982	109	64	156	2.06	SB	36	8.0	0.7	2.6	B	VG	10.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
FERMOY	2W-YOO	1955 - 1980	28	37	267	2.01	SB	42	12.0	2.1	4.2	A	G	8.8	
FERNIE	3W	1968 - 1972	64	28	138	1.82	SB	42	10.0	1.1	3.2	A	G	7.7	
FERNISKY	2W-W														
FESTIVITY	2W-Y	1977 - 1980	21	28	175	2.58	SB	35	12.1	3.3	3.8	B	VG	9.5	S
FEU DE JOIE	4W-O	1955 - 1963	-7	15	229	1.87	SB	42	13.0	1.8	nd	A	M	nd	
FIELD MARSHAL	2Y-O	1967 - 1978	-1	31	159	2.90	S	40	9.8	2.0	2.8	A	G	9.9	
FIERY CHARIOT	2Y-R	1982 - 1984	54	34	nd	1.67	B	36	11.0	1.8	3.5	A	VG	nd	
FIERY KNIGHT	4Y-O	1955 - 1963	0	17	274	1.87	SB	46	10.0	2.1	nd	A	M	nd	
FINCOOL	1W-W	1984 - 1984	35	38	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	V
FINE GOLD	1Y-Y	1964 - 1967	11	14	nd	nd	S	28	9.5	4.5	4.0	A	G	6.6	
FINGLAS	3W-WWY	1968 - 1972	68	39	174	1.52	S	36	10.0	1.6	3.8	B	G	8.9	S
FINLAND	2W-Y	1955 - 1984	-3	3	207	2.06	S	41	11.1	3.8	5.1	A	G	8.1	B
FINMACOOL	2W-W	1982 - 1984	84	37	nd	1.25	S	32	11.5	4.0	5.1	A	VG	nd	
FINOLA	1W-W	1973 - 1973	17	17	189	1.61	S	36	11.5	4.8	4.8	A	G	9.0	S
FINTONA	2W-P	1973 - 1973	62	17	258	2.00	S	30	11.0	1.8	4.2	A	VG	6.4	S
FIRE CHIEF	2Y-O	1977 - 1987	9	11	178	2.55	S	44	10.3	2.4	3.3	A	M	9.4	
FIRE GLEAM	2Y-O														
FIRE GUARD	2Y-YOR														
FIRE ISLAND	2Y-O	1972 - 1972	60	29	187	0.83	S	28	8.5	1.8	2.9	A	G	12.6	
FIREBRAND (19)	3WY-R	1955 - 1967	24	29	453	2.36	SB	50	9.0	1.0	1.7	A	P	9.0	
FIRECRACKER	2Y-O														
FIREGLOW	2Y-O	1964 - 1967	40	24	293	2.00	SB	37	9.5	1.3	3.7	A	M	8.8	
FIREMASTER	2Y-O														
FIRENZE	4W-Y	1974 - 1974	52	32	397	0.33	SB	35	9.0	2.1	nd	A	M	nd	V
FIREPROOF	2Y-O	1968 - 1980	27	27	208	2.84	SB	39	11.5	2.4	2.9	A	G	9.6	S
FIRETAIL (18)	3W-R	1964 - 1967	20	35	185	2.80	S	42	10.0	0.7	2.2	A	M	8.8	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
FIREWATER	3W-YYO	1972 - 1972	-7	34	240	1.38	S	40	10.5	1.2	2.3	A	G	8.4	
FIRST FROST	2W-W	1975 - 1975	55	23	nd	nd	SB	42	10.5	3.7	2.3	A	G	nd	V
FLAMENCO	2W-O	1955 - 1980	5	27	250	2.17	SB	47	10.0	1.4	3.4	A	G	8.3	B
FLAMING METEOR	2Y-R	1976 - 1981	6	19	144	2.49	SB	37	12.0	2.2	3.0	B	G	9.5	
FLAMING MORN	2Y	1976 - 1981	11	8	158	2.13	SB	47	10.8	3.0	3.7	A	G	9.4	B
FLANEUR	11Y-Y	1964 - 1967	46	18	nd	nd	SB	35	10.0	3.6	9.0	B	G	6.8	
FLASH	2Y-O	1972 - 1972	-36	18	306	1.92	S	43	10.5	2.1	3.2	A	G	11.0	
FLEURIMONT	2Y-YYR	1955 - 1981	24	30	210	2.12	SB	38	10.3	2.3	4.5	A	VG	8.7	
FLEVO	1Y-Y	1972 - 1973	4	4	151	1.75	S	34	10.2	4.5	4.8	A	G	9.6	
FLORA TRISTAN	2Y-O	1955 - 1963	7	20	433	1.87	SB	46	11.0	2.2	5.0	A	G	nd	
FLORA'S FAVOURITE	2W-Y														B
FLORIADE	2Y-O														
FLOWER CARPET	1Y-Y	1955 - 1973	-3	7	198	2.05	E	40	11.0	4.3	5.0	A	G	8.6	
FLOWER DRIFT	4W-OYO	1977 - 1980	40	88	165	2.77	S	32	10.4	1.5	nd	B	G	8.4	
FLOWER RECORD	2W-WWO	1955 - 1973	24	22	311	2.83	S	41	10.1	1.7	3.6	A	M	8.8	
FLOWERSONG	2Y-YYO	1970 - 1987	18	8	155	2.55	S	42	10.7	2.6	4.6	A	G	8.8	S
FLUSH	2W-P	1955 - 1963	-4	16	301	1.33	S	41	12.0	3.5	3.7	B	G	7.0	
FLYING SAUCER	2W-Y	1972 - 1980	22	19	171	2.31	S	41	10.5	2.4	5.3	A	G	9.6	B
FOGGY DEW	3W-W	1964 - 1967	40	30	198	1.90	SB	42	9.0	1.3	3.3	B	G	8.0	
FOLLY	2W-O	1972 - 1972	66	38	53	1.46	SB	32	10.0	1.3	2.8	A	G	9.0	
FONDANT	2W-Y	1977 - 1978	50	31	198	2.39	S	34	12.0	4.4	4.2	B	G	8.3	
FORAY (18)	2W-WWP	1972 - 1972	-11	34	223	1.29	S	40	11.0	3.3	3.9	B	G	9.4	
FORERUNNER	1Y-Y	1955 - 1967	2	-12	188	1.84	S	40	11.0	4.1	4.9	A	G	9.0	B
FORESCATE	1Y-Y	1972 - 1973	15	2	211	1.79	S	30	10.8	4.6	5.8	A	G	7.3	
FORESIGHT	1W-Y	1955 - 1981	19	3	173	2.01	S	32	10.0	4.2	4.3	A	G	7.9	
FOREST FIRE	2Y-O	1964 - 1967	51	33	nd	nd	SB	41	10.0	2.1	3.4	A	M	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
FORFAR	3W-O	1955 - 1973	2	25	229	2.02	SB	56	10.0	1.4	3.0	A	G	9.7	
FORTHRIGHT	2Y-O	1968 - 1971	77	34	173	1.60	SB	36	10.5	3.0	4.3	B	VG	10.7	S
FORTISSIMO	2Y-O	1972 - 1972	8	29	172	2.17	SB	28	14.0	3.8	5.2	A	G	8.0	
FORTUNE	2Y-O	1955 - 1987	18	9	174	2.19	SB	48	10.5	3.2	4.0	A	G	8.3	
FORTUNE'S BLAZE	2Y-O	1955 - 1963	17	20	446	2.40	S	50	8.0	2.1	3.1	A	M	nd	
FORTUNE'S CREST	2Y-O	1955 - 1980	7	12	201	2.45	S	46	10.0	2.4	3.4	A	G	8.5	
FORTUNE'S SUN	2Y-O	1955 - 1963	-11	17	359	1.87	SB	46	8.0	1.7	3.4	B	M	nd	
FORTWILLIAM	1Y-Y	1973 - 1981	20	28	129	2.01	S	39	11.1	4.8	4.2	A	G	8.8	BS
FOXHUNTER	2Y-O	1968 - 1971	19	32	138	1.70	SB	43	12.0	2.6	3.8	A	VG	8.3	S
FOYLE	2W-P	1964 - 1967	-5	18	nd	nd	S	32	11.0	3.4	3.9	B	G	8.4	
FRA DIAVOLO	1Y-Y														
FRANCISCUS DRAKE	2W-YYO	1955 - 1963	-7	22	322	1.87	SB	48	12.0	2.3	3.2	A	M	nd	
FRANKLIN	2Y-Y	1964 - 1980	16	12	232	2.92	S	38	11.0	3.8	5.2	A	G	8.2	
FRENCH SOL	8Y-Y	1964 - 1964	-70	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
FRIENDLY	2Y	1968 - 1980	14	20	188	2.76	S	38	10.8	3.5	5.3	A	G	7.3	S
FRIGID (17)	3W-GGW	1968 - 1972	85	34	116	1.31	S	44	9.0	0.7	1.8	A	G	10.4	
FRILEUSE	11W-Y														
FRILLED BEAUTY	2Y-Y	1955 - 1963	32	18	233	1.87	S	42	10.0	2.8	3.3	B	M	nd	
FROLIC	1W-Y														
FRONTIER	1Y-Y	1972 - 1972	29	13	89	1.12	S	37	11.0	4.8	5.0	A	G	9.0	
FROST AND FLAME	3W-R	1973 - 1973	61	32	270	1.27	S	34	10.0	1.5	3.8	B	G	9.8	S
FROZEN	2W-W	1964 - 1967	49	31	nd	nd	S	29	10.0	2.7	4.4	B	G	6.0	
FURBELOW	4Y-O	1973 - 1984	11	36	148	2.14	S	44	9.0	1.9	nd	A	G	9.9	
FURY	2Y-O	1968 - 1980	40	29	154	2.70	SB	40	10.8	2.5	3.5	A	G	9.0	BS
GADDESBY	2W	1964 - 1967	8	38	250	2.10	S	36	11.5	2.4	3.8	A	G	9.6	
GAIL O'KAHN	2W-Y														

Cultivar	Division	Years in trial	Emerg- ence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower sub- stance	Vase- life (days)	Disease notes
GALA	2W-O														
GALASHIELS	1Y-Y	1976 - 1987	7	-7	172	2.43	E	38	12.1	4.7	5.9	A	M	8.0	
GALAXY	2W-P														
GALCADOR	2Y-O	1955 - 1963	49	23	610	1.33	SB	36	9.0	1.6	3.4	A	G	nd	
GALILEE	3W-GWO	1968 - 1972	28	32	159	1.75	SB	38	10.8	1.4	3.4	A	G	8.6	
GALWAY	2Y-Y	1955 - 1963	17	12	280	1.33	E	38	11.0	4.4	4.6	A	VG	7.0	
GARDEN GIANT	2Y	1964 - 1967	-2	23	267	1.70	SB	37	10.5	2.5	4.9	A	G	8.2	
GARDEN PRINCESS	6Y-Y	1955 - 1963	7	4	324	2.93	S	35	10.0	3.5	4.0	B	M	nd	
GARDENVALE	2W-OOY	1977 - 1987	23	35	222	2.11	B	43	10.0	2.3	4.1	A	M	9.6	BS
GARLAND	2Y-YOO	1955 - 1973	32	34	238	1.62	S	38	9.9	1.5	3.8	B	G	8.6	
GARRON	1Y-Y	1955 - 1982	23	18	198	2.19	S	38	12.0	4.9	4.5	B	G	7.4	B
GARTAN	2Y-O	1964 - 1967	55	57	nd	nd	S	38	8.5	1.7	2.7	B	G	nd	
GATH-A-BAWN	2W-W	1967 - 1967	70	45	153	0.70	S	28	10.5	3.3	3.2	B	G	nd	
GAUCHO	2Y-O	1964 - 1980	-6	12	211	2.76	S	35	10.0	2.5	3.8	A	G	7.0	
GAY BOY (18)	2Y-R	1964 - 1967	-50	15	nd	nd	S	44	9.0	1.9	3.1	A	G	10.0	
GAY COLOURS	2W-R	1982 - 1984	87	50	nd	1.25	B	27	10.0	1.4	3.3	B	G	nd	
GAY RECORD	4W-O	1977 - 1984	-7	43	146	1.76	SB	49	9.1	1.0	nd	A	VG	9.6	B
GAY TIME	4W-R	1968 - 1982	36	35	137	2.00	SB	49	11.0	1.4	nd	A	VG	10.9	
GAYLORD	2Y-YOO	1981 - 1985	-15	6	291	3.40	S	45	10.2	2.7	3.0	A	G	8.8	
GEHEMA	2Y-Y	1955 - 1963	-11	31	268	1.87	SB	49	11.0	2.8	4.2	A	M	nd	
GEISHA	2W-P	1972 - 1972	77	37	114	1.58	S	32	12.5	4.0	5.3	A	VG	11.2	
GEM OF ANTRIM	2W-P	1977 - 1987	32	38	159	2.44	E	39	11.0	2.2	3.1	A	M	10.0	
GEM OF ULSTER	2W-P	1977 - 1981	28	48	220	1.99	S	46	10.7	2.3	3.1	B	VG	10.3	
GENERAL PATTON	1W-Y	1964 - 1967	-9	22	nd	nd	S	32	10.5	4.8	6.1	B	G	nd	
GENERAL SMUTS	2Y-O	1955 - 1963	-14	12	409	1.33	SB	44	11.0	1.9	4.3	B	G	nd	
GENTILITY	2W-W	1968 - 1982	94	24	148	2.54	S	31	10.8	2.8	3.8	A	G	10.2	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
GEORGE ARMSTRONG	2Y-Y														
GEORGE LEAK	2W-O	1971 - 1987	3	34	150	2.27	B	45	11.4	1.9	4.4	A	G	9.4	
GEORGE SAND	2W-O														
GEORGIA MOON	2Y-Y	1973 - 1981	-4	9	168	2.38	S	38	11.6	3.8	5.5	A	G	12.5	S
GERANIUM	8W-O	1955 - 1980	-8	33	187	2.36	SB	39	6.9	0.7	1.8	A	G	8.4	
GERTIE MILLAR	2W-Y	1955 - 1963	0	22	242	1.87	SB	38	10.0	4.0	4.7	A	M	nd	
GERVO	2W-W	1964 - 1967	-12	26	nd	nd	SB	28	12.0	3.4	6.4	B	G	6.0	
GHANA	2Y-Y	1976 - 1976	59	21	138	1.25	S	36	10.5	3.8	2.3	A	VG	nd	V
GIANT MUTICUS	1Y-Y														
GIBALTAR	2Y-O	1964 - 1967	20	24	166	1.90	S	43	8.5	1.6	2.9	B	G	9.2	
GILFORD	3W-O	1968 - 1973	36	45	137	1.22	S	40	10.2	1.0	2.8	A	G	9.2	
GIOVANNI	2W-O	1976 - 1977	30	22	150	2.15	SB	44	10.2	1.6	3.6	A	VG	9.0	
GIpsy Moth	2W-W	1977 - 1978	48	34	200	2.05	S	34	9.8	1.9	2.9	A	G	9.0	
GLACIER	1W-W														
GLANDORE	2W-W	1984 - 1984	77	50	nd	1.50	nd	nd	nd	nd	nd	nd	nd	nd	V
GLEAMING	2Y-O	1955 - 1963	-7	17	374	2.40	SB	49	10.0	2.6	3.2	A	G	8.0	
GLEN FIRE	2W-O	1955 - 1963	-4	37	327	2.40	SB	47	12.0	2.3	3.5	A	G	nd	
GLEN SPRINGS	2W-YYO	1955 - 1963	14	24	290	1.87	S	42	10.0	2.2	5.0	A	M	nd	
GLENBANE	1W-W	1964 - 1967	56	40	nd	nd	S	28	10.5	4.5	4.4	A	G	nd	
GLENBURVIE	2W-OOR	1976 - 1977	38	16	148	1.15	SB	38	9.2	1.6	3.2	A	G	8.7	V
GLENBUSH	2W-W														
GLENCAIRN	1W-Y	1964 - 1967	8	28	nd	nd	S	36	11.5	5.2	5.1	A	G	9.0	
GLENDALOUGH	2W-W	1964 - 1967	19	23	210	2.00	S	31	11.5	3.9	4.0	A	G	8.6	
GLENDERMOTT	2W-W	1973 - 1973	32	31	214	1.33	S	26	12.0	4.2	4.8	B	VG	10.4	S
GLENGARRIFF	1W-YYW	1964 - 1967	20	15	172	1.90	S	25	11.0	4.6	5.0	B	G	8.4	
GLENGORMLEY	2W-OOY														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
GLENJESLJE	2W-W	1964 - 1967	5	32	187	2.80	S	33	10.5	2.7	3.3	B	G	9.0	
GLENMANUS	2W-W	1968 - 1977	51	46	143	1.92	S	38	10.9	3.4	3.8	B	G	9.6	B
GLENOCOM	2W-W	1964 - 1970	7	32	185	2.36	SB	38	11.8	4.2	4.8	B	G	8.7	S
GLENSHANE	2W-W	1972 - 1976	20	34	128	1.90	S	37	10.8	2.0	4.0	A	G	9.0	VS
GLENSHESK	1W-W	1972 - 1972	-1	19	52	0.71	S	27	11.0	4.9	4.9	B	G	8.0	
GLENTIES	2W-O	1972 - 1973	54	46	130	1.65	S	38	10.2	1.5	3.0	A	G	10.9	V
GLENVAR	3W-R	1968 - 1972	56	44	84	1.82	SB	40	9.0	1.3	2.8	A	G	9.9	
GLENWHERRY	3W-R	1964 - 1967	11	37	271	2.60	SB	53	10.0	0.9	2.8	A	VG	10.8	
GLORIA	1W-W	1955 - 1963	21	12	198	1.87	S	34	11.0	4.7	4.6	A	G	nd	
GLORIA MUNDI	2Y-YOR	1955 - 1967	12	32	162	1.36	S	42	9.0	1.7	2.5	A	P	7.0	
GLORIFICATION	2W-YOO	1964 - 1967	41	16	nd	nd	SB	35	10.5	2.0	4.6	A	G	7.4	
GLORIOSUS	8W-Y														
GLORIOUS	8W-O	1955 - 1967	-30	17	265	3.38	SB	49	7.0	0.8	1.9	A	M	8.3	
GLORY OF LISSE	9W-YYR	1964 - 1967	40	50	270	3.00	S	32	8.5	0.4	1.5	B	M	9.8	
GLORY OF SASSENHEIM	1W-Y	1955 - 1967	32	14	133	1.56	S	32	12.0	5.0	4.2	B	M	7.8	
GLOWING PHOENIX	4Y-O	1955 - 1963	-4	11	163	1.87	SB	37	10.0	2.3	nd	A	P	nd	
GLOWING RED	4W-R	1976 - 1984	9	30	192	1.70	SB	36	9.2	1.5	nd	A	G	8.0	B
GLYNVER	3W-O	1968 - 1980	18	53	166	2.42	SB	46	9.5	0.9	2.8	A	VG	10.8	
GOBLET	1W	1973 - 1973	9	14	326	1.79	S	29	10.5	4.8	5.8	A	G	8.0	
GODOLPHIN	1Y-Y	1955 - 1973	10	11	344	1.68	E	38	12.0	5.0	4.5	A	G	9.7	
GODREY	1Y-Y	1955 - 1963	-9	13	236	1.87	E	36	10.0	4.7	5.6	B	G	nd	
GOG	1Y-Y	1976 - 1976	9	14	104	2.25	S	50	10.5	4.0	5.4	A	G	7.0	V
GOLD BULLION	1Y-Y	1975 - 1975	27	9	nd	nd	S	40	10.0	4.5	3.8	A	G	8.0	V
GOLD COLLAR	11	1964 - 1967	47	28	287	1.30	S	29	11.0	3.7	9.9	B	G	7.0	
GOLD CROWN	2W-Y														
GOLD MEDAL	1Y-Y	1955 - 1963	60	18	232	1.87	S	29	11.0	4.9	5.0	A	VG	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf pose	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
GOLD RIM	2	1972 - 1972	31	29	164	2.00	S	23	10.5	2.7	4.9	B	G	9.0	
GOLD RUSH	11Y-YOY	1971 - 1977	-2	43	128	1.23	SB	38	10.5	2.8	5.9	A	G	8.6	
GOLDBEATER	1Y-Y	1955 - 1963	-9	1	309	2.40	E	44	10.0	4.9	4.6	B	G	nd	
GOLDCOURT	1Y-Y	1955 - 1967	26	23	194	1.62	E	35	10.0	4.0	3.5	A	VG	8.8	
GOLDEN ACE	2Y-Y	1976 - 1980	22	13	164	2.00	S	42	10.0	3.8	3.2	A	G	10.8	
GOLDEN ADVANCE	1Y-Y	1964 - 1967	10	2	nd	nd	S	31	12.5	5.5	6.3	A	VG	8.0	
GOLDEN AGE	1Y-Y	1972 - 1980	24	23	130	2.01	S	38	9.8	4.3	3.5	A	G	7.4	
GOLDEN AMBER	2Y-ORR	1982 - 1984	33	22	nd	1.30	B	38	10.5	2.5	3.6	A	VG	nd	
GOLDEN BRACELET	2Y-O	1968 - 1972	50	14	152	1.88	SB	30	10.5	2.5	4.8	A	G	6.6	
GOLDEN CASTLE	4Y-O	1955 - 1963	7	3	152	0.80	S	39	11.0	4.2	nd	A	G	nd	
GOLDEN CHEER	2Y	1976 - 1977	-12	11	102	1.98	S	45	13.0	3.8	4.4	A	G	10.5	
GOLDEN CLARION	1Y-Y	1976 - 1981	3	26	106	1.71	S	38	10.3	4.4	4.5	A	G	9.4	
GOLDEN CLOUD	1Y-Y	1976 - 1979	8	30	66	1.44	S	35	10.4	4.7	4.4	B	VG	8.2	B
GOLDEN COCKEREL	1Y-Y	1968 - 1977	44	30	168	1.80	S	39	10.4	4.3	4.3	A	VG	7.8	S
GOLDEN CYCLE	6Y-Y														
GOLDEN DAWN	8Y-O	1970 - 1982	-45	2	240	4.73	S	31	4.8	0.7	1.6	B	M	10.6	
GOLDEN DAY	1Y-Y	1983 - 1987	29	20	146	2.92	S	41	12.8	6.1	5.3	A	G	5.6	
GOLDEN DOLLAR	1Y-Y	1964 - 1967	36	24	nd	nd	S	38	9.5	4.6	4.3	A	G	8.2	
GOLDEN DUCAT	4Y-Y	1955 - 1984	2	18	172	1.73	E	54	11.3	2.7	nd	A	G	9.2	B
GOLDEN EMPEROR	1Y-Y	1955 - 1963	14	31	112	1.33	S	38	9.0	4.3	4.2	A	P	6.0	
GOLDEN EMPIRE	1Y-Y	1970 - 1987	13	13	140	2.27	S	40	11.1	4.7	4.5	A	G	10.9	
GOLDEN GOBLET	7Y-Y	1955 - 1963	-11	17	121	0.80	E	38	11.0	3.9	4.8	B	G	nd	
GOLDEN GODDESS	1Y-Y	1972 - 1972	29	17	116	1.29	S	34	12.0	5.0	5.1	A	G	11.6	V
GOLDEN HARBINGER	1Y-Y	1955 - 1967	0	1	194	1.98	E	42	10.0	3.9	4.0	A	M	8.2	
GOLDEN HARVEST	1Y-Y	1955 - 1987	0	0	194	2.27	S	41	12.1	5.0	5.0	A	G	9.4	B
GOLDEN HORN	1Y-Y	1972 - 1980	26	26	132	2.20	S	38	10.0	4.6	3.6	A	VG	8.9	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
GOLDEN INCENCE	7Y-Y														
GOLDEN LACQUER	6Y-Y	1955 - 1967	22	6	188	2.45	S	36	10.0	4.4	4.1	A	G	7.0	
GOLDEN MARVEL	1Y-Y	1955 - 1982	1	5	188	1.98	E	38	11.0	4.8	3.9	A	G	10.6	
GOLDEN MELODY	1Y-Y	1964 - 1967	20	16	186	2.30	S	42	10.0	4.8	4.3	A	G	8.0	
GOLDEN PERFECTION	7Y-Y	1955 - 1979	-13	26	205	2.38	SB	50	10.0	2.3	3.3	B	G	9.7	V
GOLDEN RALLY	1Y-Y	1977 - 1979	16	22	48	1.26	S	31	12.5	4.8	6.4	B	VG	8.2	
GOLDEN RAPTURE	1Y-Y	1972 - 1981	18	27	143	2.02	S	39	11.6	4.9	4.1	A	VG	9.3	
GOLDEN RAY	1Y-Y														
GOLDEN RIOT	1Y-Y	1964 - 1982	36	32	165	2.15	S	31	11.0	5.0	4.7	A	G	7.6	
GOLDEN ROBIN	1Y-Y	1976 - 1987	3	9	170	1.94	E	46	11.7	4.8	4.7	A	M	10.7	
GOLDEN SCEPTRE	7Y-Y	1955 - 1963	-45	15	278	2.67	E	44	8.0	2.9	3.2	B	P	nd	
GOLDEN SMILE	1Y-Y														
GOLDEN SPRAY	2Y-Y														
GOLDEN SPUR	1Y-Y	1955 - 1967	17	3	215	2.04	E	27	10.0	4.3	4.1	B	M	7.0	
GOLDEN STARLIGHT	1Y-Y	1977 - 1987	39	11	116	0.98	E	39	11.6	4.8	4.5	nd	M	8.8	
GOLDEN TOP	1Y-Y	1972 - 1972	17	16	124	2.04	S	35	10.5	4.8	5.3	A	G	8.0	
GOLDEN TORCH	2Y-Y	1955 - 1980	4	16	215	1.95	S	49	11.0	3.9	3.9	A	VG	7.0	
GOLDEN TRIUMPHATOR	2Y-Y	1955 - 1963	-7	12	281	1.87	SB	46	11.0	3.8	4.9	B	G	nd	
GOLDEN VINTAGE	2Y-Y	1977 - 1981	-7	3	164	1.66	SB	50	10.0	3.2	4.3	A	G	9.6	
GOLDEN WEALTH	1Y-Y	1964 - 1967	20	30	nd	nd	S	34	11.0	5.4	5.3	A	G	8.4	
GOLDFLAKE	2Y-Y	1972 - 1972	1	18	148	2.21	S	45	9.5	2.6	3.4	A	G	13.2	
GOLDLOCKS	7Y-Y	1955 - 1967	-2	28	200	1.62	E	47	8.0	1.3	2.1	B	G	11.2	
GOLDSBOROUGH	1Y-Y	1968 - 1972	2	27	156	1.51	S	43	10.8	4.4	4.7	A	G	10.4	S
GOLYER	1Y-Y	1968 - 1971	70	25	176	2.00	S	36	11.5	4.9	4.6	A	G	6.5	
GOORNONG	1W-Y	1977 - 1977	57	16	255	1.00	S	36	11.0	5.1	3.8	A	M	9.2	
GOVERT FLINK	3W-GYO	1955 - 1963	0	24	432	2.40	SB	56	10.0	1.3	3.5	A	VG	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
GOYSCAS	3Y-YYO	1964 - 1973	27	21	260	1.98	SB	42	10.0	1.4	3.3	B	G	7.0	
GRACE MOFFAT	2Y-O	1955 - 1967	-1	17	286	2.04	S	39	11.0	2.1	3.7	A	G	9.1	
GRACKLE	2Y-O	1955 - 1963	14	15	203	1.87	S	44	10.0	2.2	3.7	B	VG	nd	
GRANADA	2Y-O	1955 - 1963	-7	4	302	2.40	SB	46	10.0	3.2	3.5	A	M	9.0	
GRAND GALA	1Y-Y	1977 - 1978	19	12	165	1.62	S	33	12.8	5.2	5.0	B	G	9.4	
GRAND MONARQUE	8W-Y														
GRAND NATIONAL (18)	2Y-R	1955 - 1963	-14	15	375	1.87	S	46	11.0	2.5	4.1	A	G	nd	
GRAND PRIMO CITRONIERE	8W-Y														
GRAND SOLEIL D'OR	8Y-O														
GRANDIS	1W-Y	1955 - 1967	6	22	258	1.98	S	37	10.0	4.2	4.6	B	M	6.4	
GRANVELLE	2W-O	1964 - 1967	47	13	nd	nd	E	31	10.5	2.4	5.2	A	G	7.6	
GRAPE FRUIT	1Y-Y	1955 - 1963	29	27	258	1.07	S	30	11.0	4.5	4.8	B	G	nd	
GRAYLING	2W-YYW	1955 - 1963	-20	17	324	1.87	E	44	11.0	3.7	3.4	A	G	nd	
GREAT LEAP	4Y-Y	1973 - 1980	48	44	155	2.34	S	46	10.2	2.8	nd	A	M	8.2	S
GREAT WARLEY	2W-Y	1955 - 1967	14	23	190	1.98	S	37	12.0	3.0	4.2	B	M	8.0	
GREEN HOWARD	3W-GYY	1964 - 1980	24	30	220	2.81	SB	42	9.5	1.4	3.3	A	G	8.2	
GREEN ICE	2W-GWW	1967 - 1968	45	52	181	2.00	S	36	10.0	2.4	4.0	A	G	10.1	
GREEN ISLAND	2W-GWY	1955 - 1982	11	31	268	1.95	S	52	10.0	2.0	4.1	B	VG	9.4	
GREEN MINT	3W-W														
GREENCASTLE	3W-GYO	1968 - 1972	20	42	150	1.23	SB	38	9.8	1.0	3.0	A	G	9.6	
GREENLAND	2W-W														
GREENMOUNT	3W-Y	1977 - 1980	-1	47	280	2.26	SB	42	9.9	1.2	3.0	B	G	9.0	
GREENORE	2W-WWY	1964 - 1967	8	18	144	2.20	S	39	11.0	2.8	4.3	B	VG	6.6	
GREETING	2W-Y	1955 - 1963	-7	24	267	1.87	S	48	12.0	2.2	3.3	B	VG	nd	
GREGALACH	1W-Y														
GRENOBLE	2Y-Y	1968 - 1987	9	-8	147	2.07	S	46	10.8	3.2	4.0	A	G	7.9	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
GRETA GARBO	2Y-O														
GREY LADY	3W-WWP	1964 - 1967	-5	51	nd	nd	S	38	6.5	0.5	2.1	B	G	7.0	
GRIANAN	1W-Y	1982 - 1984	70	36	nd	1.75	S	36	12.3	5.4	4.4	A	VG	nd	
GRULLEMANS' GIANT	2W-YYO														
GUARDIAN	2W-Y	1955 - 1973	30	26	187	2.12	E	44	11.0	4.0	3.7	A	VG	10.1	
GUIDING STAR	2W-O	1970 - 1970	14	40	181	2.19	SB	36	11.5	1.8	4.1	A	VG	8.2	
GULLIVER	3Y-YYO	1955 - 1967	4	36	290	2.90	EB	49	10.0	1.3	3.1	A	G	9.6	
GUSTAV MAHLER	2W-Y	1972 - 1972	58	31	91	0.79	S	31	9.5	3.4	5.6	A	M	7.0	
GUVEZNE	2W-O	1964 - 1967	51	49	nd	nd	SB	36	11.5	1.8	4.4	A	G	9.8	
GWENTER	2W-W	1955 - 1963	0	1	288	2.40	S	34	10.0	3.9	3.7	A	M	7.4	B
GWILLS	3W-GWO	1955 - 1963	-7	38	298	1.87	S	48	8.0	1.0	2.9	A	G	nd	
GYRFALCON	2W-Y	1964 - 1967	-15	22	nd	nd	S	29	10.0	3.9	3.7	A	G	6.8	
HADES	2Y-O	1955 - 1967	2	28	272	2.14	SB	56	11.0	1.3	3.2	A	G	10.0	
HAKA	2Y	1976 - 1977	28	15	207	2.13	S	36	8.2	3.5	3.4	A	G	7.9	V
HALF MOON	5Y-Y														
HALSKRAAG	11W-W	1964 - 1967	13	30	200	1.20	S	38	10.5	2.6	7.1	B	M	6.4	
HALVOSE	8Y-R	1955 - 1963	-45	15	166	1.33	S	35	6.0	0.6	1.4	A	G	nd	
HALWYN	1Y-Y	1955 - 1963	14	20	286	1.87	S	41	10.0	4.8	5.0	A	G	nd	
HAMZALI (18)	3W-WYR	1968 - 1973	30	34	162	2.06	SB	42	9.8	0.9	2.6	A	VG	10.2	
HAN VAN MEERGEREN	2Y-R	1955 - 1963	-11	17	449	2.40	SB	40	10.0	2.4	3.5	B	M	nd	
HANDCROSS	2Y-Y	1972 - 1980	45	24	163	2.37	S	34	10.9	3.7	4.3	A	VG	9.0	VS
HANNIBAL	11Y-O	1973 - 1982	16	47	122	1.77	SB	41	9.5	2.2	5.3	B	G	8.8	
HARBINGER OF SPRING	1Y-Y	1982 - 1985	8	-7	240	1.36	S	35	10.8	4.5	6.6	B	G	8.7	
HARDY	3W-O	1955 - 1963	14	24	192	2.40	SB	50	12.0	1.6	3.1	A	G	nd	
HARRIER	2Y-O	1968 - 1979	4	22	182	2.34	SB	40	10.8	2.3	3.7	A	G	8.6	VB
HARROGATE	2Y-YYR														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
HARTSDOWN	1W-W	1977 - 1977	50	22	300	1.40	SB	34	12.5	5.2	5.3	A	VG	10.8	
HARVEST MOON	5Y-Y	1964 - 1967	20	26	nd	nd	S	27	9.0	4.0	3.8	B	M	6.2	
HARVESTER	1Y-Y	1955 - 1963	4	10	248	1.87	S	41	10.0	4.6	4.4	A	M	nd	
HATHOR	7Y-Y														
HAVELOCK	2Y-Y	1955 - 1967	11	14	290	2.25	S	46	10.0	2.9	3.9	A	G	8.7	
HAVEMEIJER	3W	1976 - 1980	8	30	162	2.63	SB	39	10.2	1.3	2.9	B	G	9.2	
HAWAII	4Y-O	1977 - 1981	-13	35	263	1.38	SB	46	10.0	1.6	nd	A	G	9.9	BS
HAYLE	2Y/2Y-YOO	1970 - 1970	28	46	199	2.14	SB	41	11.5	2.7	3.4	A	V	9.0	
HAZEL BRILLIANT	2Y-O	1983 - 1983	29	20	nd	1.29	B	32	13.0	2.3	5.2	nd	G	nd	
HAZELWOOD	3W-WWY														
HEART'S DELIGHT	2Y-Y	1955 - 1963	-20	18	277	2.40	SB	52	11.0	2.9	5.1	A	M	nd	
HEAVEN	2W-OOY	1972 - 1972	31	62	51	1.75	S	42	11.0	1.9	3.9	A	G	10.2	
HELEN F. OHMS	2Y-W	1964 - 1967	0	15	nd	nd	SB	46	11.5	2.8	4.2	A	G	8.4	
HELFDORD	1Y-Y	1955 - 1963	10	5	278	1.87	S	41	9.0	3.9	3.6	A	M	nd	
HELIOS	8Y-Y	1955 - 1963	-14	17	201	1.33	SB	34	6.0	0.7	1.9	A	M	nd	
(syn. ABUNDANCE)															
HELIOS (16)	2Y-O	1955 - 1967	3	4	262	2.60	SB	48	10.0	3.0	3.3	A	M	9.3	
HENRY DETERDING	1W-Y	1955 - 1963	-7	15	282	2.40	S	33	10.0	4.6	4.4	B	G	nd	
HENRY IRVING	1Y-Y	1955 - 1967	40	7	357	2.24	S	28	8.0	3.4	3.2	A	M	7.0	
HERBERT BARR	1Y-Y	1955 - 1963	0	13	286	1.87	S	42	11.0	4.4	4.7	A	M	nd	
HESLA	7Y-Y	1955 - 1963	-20	17	204	2.67	S	51	8.0	1.0	2.1	B	P	nd	
HEXAGON	2W-W	1955 - 1963	4	17	241	1.87	S	42	10.0	3.8	4.8	A	M	nd	
HEXAMETER	9W-GYR	1955 - 1982	26	51	277	2.10	S	55	8.0	0.6	2.0	A	G	11.7	
HI KING	1Y-Y	1974 - 1974	49	15	474	1.00	S	37	10.5	4.1	3.6	A	G	11.2	
HIWASSEE	8W-W	1976 - 1982	-30	-20	225	1.84	S	23	5.0	0.4	1.1	B	M	11.4	
HIGH LIFE	2W-OOY	1955 - 1982	4	27	262	2.37	SB	45	11.0	2.1	5.0	B	M	8.3	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
HIGHFIELD BEAUTY	8Y-GYO	1976 - 1986	0	22	220	4.60	B	41	8.2	1.3	2.9	A	G	9.2	
HIGHLAND CASTLE	1W-Y	1973 - 1973	64	32	331	1.46	S	27	11.0	5.0	5.3	B	VG	10.2	S
HILLMOUNT	2W-R	1964 - 1967	10	34	nd	nd	SB	41	10.5	1.3	3.3	A	VG	9.2	
HILLSBOROUGH	1W-Y	1964 - 1967	51	29	nd	nd	S	27	10.0	4.4	4.1	B	G	6.6	
HILLSTON	1Y-Y	1972 - 1979	-8	6	112	1.83	S	43	10.8	4.6	5.0	B	G	8.2	VS
HIMALAYA	1W-W	1964 - 1967	-5	18	nd	nd	S	29	12.0	5.4	6.3	A	VG	8.8	
HINDENBURG	1Y-Y	1955 - 1963	43	15	140	1.33	S	28	11.0	4.7	5.7	B	VG	nd	
HINDUSTAN	1W-W	1964 - 1967	34	21	nd	nd	S	31	11.0	5.1	5.8	A	VG	7.8	
HIS EXCELLENCY	1Y-Y														B
HOLLAND	2Y-O	1955 - 1973	5	15	230	1.89	S	48	10.0	2.1	4.1	A	VG	7.8	
HOLLANDIA	4Y-O	1955 - 1963	-4	8	237	1.87	SB	44	10.0	2.5	nd	B	M	nd	
HOLLY BERRY	2Y-R	1968 - 1982	48	28	224	2.39	SB	42	9.8	2.1	3.7	A	VG	8.7	S
HOLLYWOOD	2Y-O	1964 - 1987	21	0	194	2.40	S	39	10.8	2.9	4.0	A	G	9.5	B
HOME FIRES	2Y-O	1968 - 1980	-6	18	194	2.20	S	43	10.5	2.6	3.2	B	G	10.6	
HOMER	9W-OOR	1964 - 1967	36	33	nd	nd	SB	34	8.5	0.5	1.8	B	M	9.0	
HONEY GOLD	2Y-Y	1977 - 1981	36	21	108	1.69	S	42	11.4	4.0	4.9	A	G	10.0	S
HONEYBIRD	1Y-W	1976 - 1982	33	19	130	1.99	S	38	10.8	4.5	4.0	A	G	11.0	B
HONG KONG	2Y-O	1964 - 1972	34	18	147	1.93	SB	45	11.5	2.5	3.4	A	G	10.0	
HOPESAY	2W-YYO	1964 - 1967	-30	30	nd	nd	SB	37	12.0	1.9	4.8	B	G	8.0	
HORACE	9W-GOR	1955 - 1967	8	35	294	2.92	SB	54	9.0	0.5	1.7	A	M	9.2	
HORIZON	1Y-Y	1977 - 1977	64	19	112	1.40	S	28	10.5	4.6	4.1	B	VG	9.6	V
HORN OF PLENTY	5W-W														
HORSFIELDII	1W-Y	1955 - 1967	32	26	184	1.46	S	34	11.0	4.5	3.8	A	M	7.8	B
HOSPODAR	2Y-O	1955 - 1967	-6	6	211	2.14	E	44	10.0	2.3	3.0	A	M	8.4	
HUNTERLEA	3W-WWR	1977 - 1978	32	54	320	1.68	SB	42	10.2	3.8	2.4	B	G	10.0	
HUNTER'S MOON	1Y-Y	1955 - 1963	14	23	166	2.40	E	42	11.0	4.9	4.9	B	G	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
HUON	9W-YYO	1955 - 1963	60	42	294	1.87	SB	47	8.0	0.4	1.9	A	G	nd	
HUSKY	3Y-YOR	1955 - 1967	0	18	282	2.95	SB	43	11.0	1.9	2.8	A	P	7.0	
HUSSAR	2Y-YYO														
HYPERION	2Y-Y	1972 - 1972	60	38	183	1.08	S	23	11.0	2.8	5.8	A	VG	10.0	
H. CAMP	30-O	1980 - 1980	7	26	715	1.33	SB	30	10.0	1.5	2.4	B	M	9.8	
ICE CRYSTAL	11W-W														
ICE FOLLIES	2W-W	1964 - 1987	22	16	264	2.46	E	35	12.3	2.7	6.4	A	P	7.7	
IDEAL	8W-O	1955 - 1981	-44	16	198	2.29	SB	42	5.0	0.6	1.9	A	M	9.2	
ILLUMINATE	2Y-O	1977 - 1981	-5	5	180	1.97	SB	41	11.1	2.5	3.8	A	G	11.5	
IMOGEN	2W-P														
IMPERATOR	1W-W	1955 - 1981	43	27	142	2.09	S	40	11.0	4.9	4.8	A	G	9.3	B
IMPROMPTU	2W-GYY	1972 - 1972	63	31	172	1.31	S	30	13.5	4.4	5.7	B	G	8.4	
INA	3W-O	1955 - 1980	1	35	226	2.52	SB	54	10.0	1.6	3.8	A	VG	9.8	
INCA GOLD	1Y-Y	1976 - 1978	38	18	160	1.95	nd	nd	nd	nd	nd	nd	nd	nd	
INDIAN CHIEF	4Y-O	1955 - 1963	-4	40	261	1.87	SB	38	11.0	1.7	nd	A	M	nd	
INDIAN SUMMER	2Y-O	1955 - 1963	0	37	255	1.87	EB	54	10.0	1.7	2.5	A	G	9.4	
INDOLINE	1Y-Y	1968 - 1982	18	28	148	2.10	S	44	11.3	4.4	4.6	A	G	8.4	BS
INDONESIA	2Y-O	1955 - 1963	0	24	302	1.87	SB	44	10.0	1.6	4.6	A	VG	nd	
INFATUATION	2W-YYP	1973 - 1973	10	39	127	2.00	S	35	10.5	2.0	3.0	A	G	9.6	
INFERNO	2Y-O	1972 - 1972	60	23	144	1.75	S	30	9.0	1.9	3.2	A	VG	9.6	
INFLAMMABLE	2Y-O	1977 - 1981	24	32	200	2.38	S	38	9.4	2.2	3.1	B	G	8.4	V
INGLESOMBE	4Y-Y	1955 - 1983	16	24	213	2.16	SB	43	9.1	2.4	nd	A	P	8.0	
INGRID VALBERG	11Y-O														
INISHKEEN	1W-Y														
INNISFALLEN	2W-Y	1955 - 1980	30	38	187	2.18	SB	46	10.0	2.7	3.4	A	VG	8.6	
INSULINDE	4W-O	1955 - 1963	17	20	206	1.87	SB	49	11.0	2.1	nd	A	M	9.2	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
INTERIM	2W-YYP	1964 - 1982	26	42	188	2.78	S	41	11.5	2.2	3.2	A	G	8.8	
INTERLUDE	2W-P														
INVER	1Y-Y														
INVERELL	2W-W	1970 - 1970	40	22	209	1.50	SB	31	11.0	3.8	5.4	B	G	nd	
INVERGORDON	2Y-O	1955 - 1963	29	26	371	2.40	SB	43	10.0	2.0	4.2	A	M	nd	
INVERNESS	2Y-YOO	1976 - 1981	7	28	120	2.40	SB	48	11.2	2.0	3.9	A	VG	10.0	
INVESTMENT	1Y-Y	1974 - 1987	11	-4	152	2.32	E	40	11.9	4.9	5.5	A	M	7.9	B
IRANI	4W-Y	1982 - 1984	62	44	nd	1.16	S	33	9.3	1.5	nd	B	G	8.0	
IRENE COPELAND	4W-Y	1955 - 1981	20	24	187	1.72	SB	36	9.9	2.3	nd	A	M	8.4	
IRISH CHARM	2W-O														
IRISH LEGEND	2W-O	1979 - 1985	21	37	270	1.86	B	48	10.0	1.7	3.6	A	VG	10.6	
IRISH LUCK	1Y-Y	1955 - 1982	29	29	182	2.11	S	40	12.0	4.4	4.8	A	VG	9.0	
IRISH MINSTREL	2W-Y	1972 - 1980	45	34	187	2.07	S	42	11.1	2.4	4.0	A	VG	9.3	
IRISH PRINCE	1Y-Y	1976 - 1979	17	14	134	2.00	S	41	11.8	4.8	5.6	A	G	9.2	
IRISH RANGER	3W-O	1984 - 1984	28	42	-20	1.50	S	44	11.5	1.5	3.5	nd	G	nd	B
IRISH ROSE	2W-P	1979 - 1981	71	36	160	0.80	S	29	10.0	2.1	3.2	A	G	10.2	
IRISH SPLENDOUR	3W-R	1977 - 1980	-10	42	195	2.14	SB	46	11.5	1.4	3.8	A	G	12.1	
IRMELIN	8W-R	1955 - 1963	-54	32	271	2.40	SB	47	7.0	0.6	1.9	A	M	nd	
ISLANDREAGH	2W-Y	1973 - 1973	24	29	297	1.46	E	32	11.5	3.9	4.5	B	VG	9.8	S
ISOBELINE	2W-P	1973 - 1973	65	28	208	2.46	E	26	12.0	4.2	4.9	A	G	10.0	.S
ISOLA	3W-YO	1964 - 1967	20	39	294	2.90	SB	46	10.5	0.8	2.7	A	G	10.0	
JACINTA MARTO	2W-O	1955 - 1963	14	8	522	2.40	S	46	10.0	2.5	4.5	A	G	nd	
JACK SNIPE	6W-Y	1955 - 1967	22	11	354	3.52	S	29	8.0	2.0	2.1	A	M	7.4	
JACKPOT	2Y-R	1972 - 1987	-8	25	167	2.81	B	44	11.1	2.0	3.3	A	G	8.8	
JAGUAR	2Y-O	1968 - 1982	30	25	204	3.25	SB	38	9.8	2.0	3.9	A	G	8.9	
JALNA	2Y-R	1955 - 1963	-20	3	294	1.87	SB	46	10.0	2.6	3.1	A	M	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
JAMAICA	3W-O	1968 - 1972	50	34	223	1.45	SB	40	10.0	1.5	3.4	A	VG	9.0	
JANA	6Y-Y														B
JANUS	2Y-Y	1976 - 1981	-10	3	169	2.76	S	36	11.1	3.1	4.6	A	G	8.5	B
JANZ	2Y-R	1974 - 1974	37	30	55	0.17	SB	40	8.5	1.7	2.6	A	M	nd	V
JEANNE DESOR	2W-Y														
JENNIFER RUTH	1W-W	1964 - 1967	47	22	nd	nd	S	26	11.0	4.9	5.0	A	G	6.9	
JENNY	6W-W														
JERJOINT	2Y-YYO	1964 - 1967	40	24	163	2.10	SB	43	9.5	2.1	2.8	A	G	10.6	
JERSEY BEAUTY	2Y-O														
JERSEY CREAM	1W-Y	1955 - 1963	14	18	209	1.87	S	41	9.0	4.0	4.0	A	M	nd	
JESSONDA	2W-YYP	1955 - 1963	14	17	359	1.87	E	38	9.0	3.3	3.4	B	M	nd	
JEST	2Y-Y														
JEZEBEL	3Y-R														
JILL	2W-YYO	1964 - 1967	40	21	nd	nd	SB	35	8.5	1.3	3.5	A	M	6.8	
JINDALEE	2W-Y	1976 - 1987	27	7	168	2.52	S	34	11.8	4.0	4.9	A	M	8.1	B
JINGELIC	2W-P	1977 - 1977	50	13	221	2.00	S	28	10.0	3.4	3.9	B	G	10.2	
JOBI	1Y-Y	1977 - 1977	7	14	232	2.08	S	41	9.5	3.5	2.9	A	G	9.4	V
JOHANNESBURG	2W-O	1964 - 1967	51	26	264	1.60	SB	32	10.0	1.6	3.9	B	G	7.0	
JOHN EVELYN	2W-O	1955 - 1963	14	20	303	2.40	SB	46	11.0	2.1	3.7	B	M	nd	
JOHN GREENWAY	2W-WWY	1972 - 1973	-12	27	208	3.42	S	38	11.5	3.4	6.3	A	G	8.5	S
JOHN OF SALISBURY	2W-Y	1977 - 1980	1	32	236	2.06	S	45	9.8	2.8	3.4	A	G	10.4	B
JOKE FULMER	2W-YOO	1968 - 1982	34	24	219	2.40	SB	37	9.8	1.8	5.1	A	G	8.3	
JOLLI-COEUR	2W-YYO														
JOSEPH MACLEOD	1Y-Y	1968 - 1984	25	-6	178	2.15	S	38	12.0	4.9	5.2	A	G	12.0	B
JOYANT	2W-O	1972 - 1972	31	41	10	0.83	S	16	9.5	2.7	5.1	A	G	nd	
JOYOUS	2W-Y	1976 - 1980	38	12	120	2.00	SB	38	10.0	4.0	4.0	A	VG	7.9	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
JUANITA	2Y-O	1964 - 1967	7	7	nd	nd	SB	35	11.5	2.4	3.8	A	G	7.4	
JUBILANT	2Y-Y	1955 - 1963	21	23	303	2.13	S	48	11.0	3.4	4.1	A	G	9.0	
JUBILATION	2W-Y	1972 - 1972	8	22	201	1.88	S	40	11.5	2.5	5.1	B	G	8.2	
JULES VERNE	2W-Y	1955 - 1972	8	6	153	2.24	S	42	12.0	4.2	4.2	A	G	9.1	
JULIET	9W-YR	1968 - 1973	31	60	112	1.30	SB	47	8.0	0.6	2.1	A	G	8.8	
JULIUS CAESAR	1Y-Y														
JUNCO	1W-W	1977 - 1981	8	5	151	1.82	SB	41	11.1	4.6	5.0	A	G	11.6	BS
JUNE ALLYSON	2W	1964 - 1982	18	26	174	2.24	SB	41	10.0	1.7	4.0	A	G	7.0	
JUVENTUS	1W-Y	1973 - 1973	50	14	228	1.16	S	30	11.0	4.9	4.8	A	VG	10.0	S
KALGOORLIE	1Y-Y	1955 - 1963	14	12	400	1.87	S	37	11.0	4.2	5.0	B	VG	nd	
KANCHENJUNGA	1W-W	1955 - 1967	29	20	281	2.14	S	36	11.0	5.0	5.4	A	VG	7.6	
KANGA	1Y-Y	1976 - 1980	28	24	125	1.28	S	36	10.4	4.8	4.0	A	G	7.5	
KANSAS	3W-GYO	1955 - 1963	14	26	355	1.87	SB	63	11.0	1.5	4.0	A	G	9.0	
KARACHI	2Y-YR	1964 - 1982	0	23	172	2.49	SB	43	9.5	2.0	2.9	A	VG	8.0	
KARAMUDLI	1W-Y	1972 - 1981	36	17	130	2.12	S	37	10.6	4.4	3.8	A	G	10.6	BS
KARANIA	1W-P														
KARELIA (17)	1W-Y	1972 - 1972	78	31	38	0.38	nd	nd	nd	nd	nd	nd	nd	nd	
KATHLEEN FERRIER	2W-Y	1972 - 1973	48	32	132	1.77	SB	36	11.0	2.2	4.8	A	G	7.9	
KEN FARMER	2Y-YR	1982 - 1984	28	3	nd	1.43	B	38	10.3	2.6	3.8	A	G	7.0	
KENBANE	1W-W	1955 - 1963	14	39	165	1.87	S	37	12.0	4.6	5.5	A	VG	nd	
KENMARE	2W-P	1964 - 1967	49	26	200	2.70	SB	30	11.0	3.7	3.9	B	G	8.2	
KERRY PIPER	1Y-Y	1964 - 1967	-5	24	nd	nd	S	31	9.5	3.7	3.8	A	G	7.6	
KESTREL	9W-GOR	1955 - 1963	0	44	389	2.93	SB	53	9.0	0.6	1.9	A	M	nd	
KHARTUM	3W-R	1955 - 1963	14	31	318	1.33	SB	54	9.0	1.0	3.1	A	VG	nd	
KIBO	2W-W	1968 - 1982	56	42	154	2.10	S	42	12.2	3.8	6.5	A	G	9.2	BS
KILBRIDE	2Y-Y	1964 - 1967	20	21	203	1.90	SB	46	11.5	2.4	3.3	A	G	9.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
KILCOOL	2W-YYO	1972 - 1972	50	38	126	0.81	SB	38	nd	nd	nd	A	nd	9.8	
KILCORAN	2W-YYO	1964 - 1967	21	34	220	1.10	SB	34	10.0	1.8	4.1	A	G	8.0	
KILDRUM	3W-R	1968 - 1972	8	43	172	1.72	SB	43	10.2	1.1	3.2	A	G	9.8	B
KILJMANJARO	2W-YOO														
KILKENNY	1Y-Y	1964 - 1967	40	31	211	1.60	S	34	10.0	4.3	4.7	A	VG	7.0	
KILLALA	2W-YYO	1964 - 1967	51	55	nd	nd	S	29	10.0	3.1	3.7	B	G	6.8	
KILLALOE	2W-W	1964 - 1967	41	30	189	2.20	S	33	12.0	4.3	5.5	B	VG	8.6	
KILLARA	8W-Y	1980 - 1981	-29	35	108	1.04	SB	25	7.5	1.0	2.4	B	G	9.6	
KILLJFRETH	3W	1968 - 1972	57	36	176	2.32	SB	36	9.7	1.2	3.6	B	G	9.3	
KILLJGREW	2Y-O	1955 - 1967	-5	15	312	2.66	SB	48	10.0	2.3	3.4	B	M	8.2	
KILLYMOON	2W-W	1977 - 1977	71	28	114	0.36	SB	25	9.5	3.4	3.7	A	G	nd	
KILLYNURE	1W-Y														
KILMORACK	2Y-O	1968 - 1972	34	24	218	2.46	S	44	9.2	2.2	2.7	A	VG	10.8	
KILMORE	3W-O	1970 - 1987	8	35	166	2.23	B	44	9.6	1.6	3.2	A	G	10.2	
KILMORNA	2W-Y	1955 - 1982	28	22	214	2.41	S	46	10.0	2.8	3.6	A	G	6.3	
KILREA	2W-W	1968 - 1982	62	31	193	2.56	SB	34	11.5	4.0	4.3	A	G	8.4	BS
KILTER	3W-R	1955 - 1963	56	29	300	1.87	SB	48	10.0	1.0	2.8	A	M	nd	
KILWORTH	2W-YOO	1955 - 1982	27	39	205	2.08	SB	47	10.9	2.2	3.6	A	G	8.7	
KIMANGE	1Y-Y	1976 - 1977	27	17	234	0.88	S	35	9.0	3.6	2.4	A	G	9.6	
KINARD	2W-Y	1972 - 1972	45	39	202	1.13	SB	38	10.0	2.0	4.4	B	VG	10.0	
KINDLED	2Y-R														
KING ALBERT	1Y-Y	1955 - 1973	11	10	219	1.50	E	36	12.0	5.0	5.3	A	VG	9.5	
KING ALFRED	1Y-Y	1955 - 1967	-14	8	220	1.94	S	49	11.0	4.5	4.5	A	G	7.6	
KING CARDINAL	2W-R	1968 - 1980	44	45	133	2.16	SB	44	10.0	1.6	3.6	A	VG	8.8	B
KING KI	1Y-Y	1974 - 1974	24	15	293	0.75	S	36	11.0	4.6	3.7	A	VG	11.2	V
KING LUD	1Y-Y														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
KING OF HEARTS	2Y-O	1955 - 1967	-13	11	308	2.85	S	40	10.0	2.1	3.4	A	G	8.7	
KINGCRAFT	8W-O	1955 - 1980	-24	34	201	3.27	SB	60	7.0	0.6	2.0	A	G	9.6	
KINGDOM	2W-WWY	1955 - 1963	-14	23	203	2.40	S	50	9.0	2.9	3.5	B	G	nd	
KINGFISHER	3W-GYR	1964 - 1967	51	43	nd	nd	SB	42	8.0	0.7	2.4	A	G	nd	
KING'S RANSOM	1Y-Y	1972 - 1977	35	22	139	2.03	S	34	9.7	4.1	4.1	A	VG	8.8	
KINGSCOURT	1Y-Y	1955 - 1982	11	20	209	2.21	S	43	10.0	4.8	4.7	B	VG	6.9	
KINGSLEY FAIRBRIDGE	1W-Y	1964 - 1967	47	31	204	2.30	S	35	11.0	5.7	4.8	B	G	8.2	
KINGSTON	2W-Y														
KLAMATH	2W-Y														
KNIPP	2Y-YOO	1955 - 1963	-14	26	280	2.40	SB	60	10.0	2.1	3.2	A	P	nd	
KNOCKSTACKEN	1Y-Y	1982 - 1984	65	29	142	2.25	S	30	10.3	4.3	3.6	A	G	nd	
KNOWEHEAD	2W-W	1973 - 1973	50	32	182	2.37	S	28	11.5	4.1	4.0	A	VG	11.6	
KRAKATOA	2Y-O	1955 - 1973	15	14	199	2.15	S	40	10.5	2.8	4.3	A	G	8.7	
KWANNON	2W-Y	1964 - 1967	5	26	205	2.90	S	46	11.5	2.8	3.6	A	G	10.8	
LA ARGENTINA	2W-WOR	1964 - 1967	10	24	nd	nd	SB	45	10.5	2.2	3.9	B	G	7.6	
LA BEAUTE	3W-R	1955 - 1963	49	24	418	2.40	SB	40	9.0	1.0	3.2	A	M	8.0	
LA BELLE	7Y-O	1955 - 1963	32	56	240	1.60	SB	26	6.0	0.9	2.7	A	P	nd	
LA FIANCÉE	8W-Y	1955 - 1963	-7	17	146	1.87	S	35	6.0	0.7	1.8	A	G	nd	
LA RIANTE	3W-YOO	1955 - 1967	18	39	197	1.88	SB	44	8.0	1.0	3.0	B	M	9.5	
LABRADOR	1W-W	1955 - 1963	0	22	180	1.87	SB	36	10.0	4.1	3.7	A	G	nd	
LADY BEE	2W-P	1964 - 1967	9	45	220	1.70	S	25	8.5	2.0	2.3	A	M	8.8	
LADY BIRD	2W-O	1964 - 1967	17	40	187	1.90	S	29	10.0	4.1	5.1	B	VG	7.4	
LADY DIANA MANNERS	3W-YR	1955 - 1967	18	30	266	2.80	SB	63	9.0	1.8	3.1	A	G	7.0	
LADY GODIVA	3W-YYO	1955 - 1963	27	31	392	2.40	SB	50	10.0	1.0	2.1	A	M	nd	
LADY IVOR	2W-P	1964 - 1967	47	25	nd	nd	SB	35	9.5	1.9	3.9	A	G	10.0	
LADY JEAN RANKIN	2W-W	1964 - 1967	49	23	nd	nd	S	28	11.5	4.6	4.1	B	G	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
LADY JOWITT	2W-P	1976 - 1980	43	21	168	1.98	S	39	10.6	3.9	4.4	A	G	8.7	B
LADY KESTIVEN	3W-O	1955 - 1963	21	29	253	1.87	SB	52	11.0	1.1	2.6	A	G	nd	
LADY LUCK	2Y-O														
LADY MARGARET BOSCAWEN	2W-Y	1955 - 1967	3	24	270	2.08	S	45	12.0	3.0	3.5	B	P	6.6	
LADY MOORE	3W-OOR	1955 - 1963	71	24	484	1.87	SB	51	10.0	1.0	2.6	A	M	nd	
LADY PENRHYN	2Y-YYO	1955 - 1963	14	17	195	2.40	SB	47	9.0	2.0	3.8	A	G	nd	
LADY SERENA	9W-GYR	1967 - 1985	27	48	142	2.50	E	54	9.9	0.7	2.4	A	G	9.6	
LAETITIA	8Y-YYO	1967 - 1969	-1	31	198	0.98	S	18	6.2	0.7	1.7	A	G	10.8	
LAGONDA	2W-YYW	1955 - 1963	0	20	242	1.87	E	44	10.0	3.9	4.2	A	P	nd	
LAMINGTON	2Y-O	1964 - 1967	11	17	nd	nd	S	28	10.5	2.4	4.4	A	M	7.8	
LANARTH	7Y-O	1955 - 1967	-48	5	270	3.18	E	46	8.0	1.7	2.8	A	M	10.0	
LANCASTER	3W-GYO	1982 - 1984	54	47	nd	2.60	B	39	10.5	1.1	2.3	A	VG	nd	
LANDITHY	2Y-Y	1955 - 1963	-11	17	325	1.87	SB	45	12.0	3.6	5.0	A	G	nd	
LANDULPH	2Y-Y	1955 - 1963	17	15	418	2.40	S	43	11.0	3.9	3.9	A	G	nd	
LANGLO	1Y-Y	1976 - 1987	16	-1	169	2.46	S	38	11.7	5.0	5.1	A	M	8.1	
LANTANA	1W-Y	1968 - 1971	75	32	147	1.70	S	35	11.5	5.2	5.5	B	G	7.1	BS
LANTEGLOS	2	1968 - 1980	5	2	170	2.32	S	44	10.5	2.5	3.6	A	M	10.0	
LAPFORD	1W-Y														
LAPLAND	1W-W	1964 - 1967	49	22	nd	nd	S	29	11.5	5.0	6.4	A	VG	8.0	
LARGO	2Y-YYO	1972 - 1972	53	23	111	1.67	SB	32	10.5	1.9	4.4	B	G	9.0	
LARKELLY	6Y-O	1955 - 1967	24	26	255	2.84	S	30	9.0	1.6	2.6	A	G	7.0	
LATIFA	2W-O	1964 - 1967	26	22	nd	nd	SB	33	9.5	2.1	4.5	A	M	6.0	
LATONIA	3W-R	1955 - 1963	-4	31	220	1.87	SB	56	10.0	1.0	3.1	A	G	nd	
LATROBE	1Y-Y	1972 - 1973	28	10	194	2.10	S	42	11.2	5.0	6.2	A	G	8.7	
LAURENS KOSTER	8W-Y	1955 - 1967	6	20	355	2.85	S	36	4.0	0.6	1.5	A	M	12.0	
LAVINO	2W-P	1964 - 1967	-51	20	nd	nd	SB	35	11.0	2.2	3.9	B	G	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
LAWALI	4W-P	1976 - 1977	6	18	136	0.82	S	32	8.2	2.7	2.3	A	G	7.7	
LE BEAU	6Y-Y	1955 - 1963	27	8	190	1.33	S	28	9.0	3.7	2.8	A	M	nd	
LEANDER	2Y-O	1973 - 1982	-8	9	219	2.76	S	41	10.1	2.1	3.1	A	G	12.6	S
LEBANON	2W-Y	1970 - 1987	17	10	198	3.24	S	42	11.1	2.8	5.2	A	M	7.7	B
LEEUWENHORST	2W-YYO	1955 - 1963	14	24	364	1.87	SB	54	10.0	2.1	3.7	A	G	nd	
LEINSTER	1Y-Y	1955 - 1963	-4	20	302	1.87	SB	42	11.0	4.9	5.0	A	G	nd	
LEITRIM	2Y-Y	1968 - 1971	73	32	175	1.80	S	35	11.0	4.4	5.1	B	G	6.3	S
LEMBERG	1Y-Y	1955 - 1963	-11	17	199	1.87	S	44	12.0	5.4	4.5	A	G	nd	
LEMINOS	2Y-Y	1964 - 1967	43	24	235	2.80	SB	29	9.5	2.3	3.4	B	G	6.6	
LEMON BEAUTY	11W-Y	1968 - 1973	49	34	227	1.36	SB	35	9.8	2.8	5.6	B	G	9.0	
LEMON CUP	2W-Y	1964 - 1967	21	17	nd	nd	SB	34	11.5	2.4	5.4	A	G	6.0	
LEMON DORIC	2Y-W	1973 - 1981	14	14	163	2.12	S	36	9.9	3.4	3.6	A	G	11.4	BS
LEMON FANCY	1Y-Y	1964 - 1967	18	10	nd	nd	S	33	10.5	4.8	4.8	A	G	6.8	
LEMONADE	3Y-Y	1982 - 1984	49	37	nd	2.00	B	46	10.5	1.4	3.1	A	G	nd	
LEMSTAR	1Y-Y	1955 - 1963	-4	11	418	2.40	S	46	10.0	4.7	4.8	A	G	nd	
LENORE NETHERCOTE	2Y-YYO	1976 - 1977	-2	10	122	2.30	S	44	8.8	2.4	3.3	A	M	10.2	
LEONAINÉ	2W-P	1977 - 1980	56	34	236	2.18	S	40	9.6	2.2	3.9	B	G	10.4	
LEPRECHAUN	2Y-R	1964 - 1967	7	29	210	2.30	S	41	8.0	1.5	1.7	A	M	7.0	
LETTERKENNY	1Y-Y	1968 - 1973	-4	18	222	2.02	S	34	12.0	4.2	4.6	B	VG	7.0	S
LEVIATHAN	2Y-Y	1955 - 1963	0	38	307	1.87	SB	47	10.0	3.0	5.6	A	G	nd	
LHASA	1W-W	1967 - 1968	68	28	324	1.70	S	30	11.8	5.0	5.1	A	G	7.2	
LIBBY HOLMAN	1W-W	1972 - 1973	28	28	178	1.83	S	31	13.2	4.9	5.7	A	G	9.3	S
LIBERTY	1Y-Y	1955 - 1963	14	10	220	1.33	S	32	12.0	4.8	4.3	B	M	nd	
LIBERTY LIGHT	2W-YYO	1972 - 1972	60	25	75	3.12	SB	35	10.5	2.1	3.6	A	G	9.0	
LIBYA	2W-O	1976 - 1977	6	44	186	2.03	SB	46	9.5	1.5	3.0	A	VG	9.9	
LIGHTHEART	1W-Y														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
LIKOVAN	1Y-Y	1972 - 1977	23	17	129	1.54	S	40	11.1	4.6	5.1	A	G	10.2	BS
LILY RONALDS	2W-GYO	1975 - 1975	9	3	nd	nd	S	39	9.0	1.9	4.0	A	M	9.2	V
LILYDALE	2W-W	1975 - 1975	36	8	nd	nd	S	36	12.0	4.7	4.9	B	VG	nd	V
LIMELIGHT	1Y-Y	1968 - 1973	18	10	178	1.90	S	38	11.8	4.3	4.7	A	G	7.3	B
LIMERICK	3W-O	1964 - 1982	16	47	206	2.58	SB	45	9.0	1.1	3.3	A	G	9.0	
LIMONE	1Y-Y														
LINGERING LIGHT	2W-P														
LINN	2W-Y														
L'INNOCENCE	8W-Y	1955 - 1963	7	23	137	1.87	S	27	6.0	0.7	1.7	A	G	nd	
LINTIE	7Y-O	1964 - 1967	49	46	310	3.20	S	33	6.5	0.7	2.0	B	M	9.0	
LISANORE	2W-P	1982 - 1984	71	37	nd	2.50	S	32	10.3	3.0	3.6	A	VG	nd	
LISBANE	3W-GYR	1982 - 1984	74	53	nd	1.17	S	37	10.0	1.0	2.5	A	G	nd	
LISBREEN	2W-GOO	1964 - 1967	51	31	nd	nd	S	23	10.5	3.4	3.5	B	G	8.4	
LISCARTON	3Y-O	1955 - 1963	-7	21	274	2.40	SB	53	11.0	1.5	2.8	A	G	nd	
LISDILLON	1W-P														
LITTLE DIAMOND	3Y-O	1964 - 1967	43	24	200	2.00	SB	32	8.5	1.1	2.4	B	M	8.6	
LITTLE WITCH	6Y-Y	1964 - 1967	34	11	274	2.60	S	25	8.5	2.2	1.5	A	G	8.4	
LIVIA	4Y-O	1955 - 1963	14	17	251	1.87	SB	44	10.0	2.0	nd	A	M	nd	
LIZARD LIGHT	2Y-O	1964 - 1980	11	5	170	2.56	S	35	9.5	2.6	4.0	A	G	9.6	
LOCH HOPE	2Y-R	1981 - 1985	-17	17	241	2.42	B	37	10.4	2.6	2.5	B	VG	9.8	
LOCH MAREE	2W-P	1964 - 1967	12	24	205	2.60	S	32	11.0	3.9	4.4	A	G	8.0	
LOCHINCH	2Y-O	1967 - 1971	-10	35	188	1.90	SB	45	11.6	2.1	3.0	A	G	8.4	S
LOCHINVAR	2W-O	1977 - 1979	16	28	160	1.75	SB	42	10.2	1.5	3.6	A	G	9.2	V
LOGAN ROCK	7	1973 - 1977	-20	18	279	4.24	S	43	7.7	2.2	2.2	A	G	10.7	
LOMBARD	2Y-Y	1955 - 1963	-4	17	213	1.87	SB	44	10.0	3.5	3.7	A	G	nd	
LONGFORD	1W-W	1973 - 1973	30	29	267	0.91	S	34	12.5	5.3	3.6	B	VG	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
LORD ANTRIM	1Y-Y	1955 - 1963	-14	17	241	1.87	E	44	10.0	4.7	5.3	A	G	6.4	
LORD CHANCELLOR	1W-Y														
LORD KITCHENER	2W-Y	1955 - 1963	43	20	325	2.93	S	45	10.0	3.3	4.0	A	M	7.0	
LORD NELSON	1Y-Y	1964 - 1967	34	9	nd	nd	S	31	11.0	5.0	5.6	A	G	7.4	
LORD TEDDER	2W-GYY	1968 - 1980	26	51	138	1.92	SB	43	10.9	2.1	4.8	A	G	8.6	B
LORD WELLINGTON	1Y-Y	1955 - 1963	0	23	228	1.60	S	39	11.0	5.5	4.9	A	G	nd	
LORELEI	2W-P	1955 - 1963	0	8	224	1.87	SB	42	9.0	2.3	4.1	A	M	nd	
LORENZO	2W-O	1979 - 1985	nd	32	226	1.78	B	45	10.2	1.5	3.4	A	G	9.0	
LOTHARIO	2W-Y	1972 - 1980	37	51	108	2.49	SB	45	10.4	2.7	3.6	B	G	8.5	
LOTTA SVARD	1W-Y	1955 - 1963	0	22	275	1.33	S	42	11.0	5.0	4.2	A	G	nd	
LOUGH ARBEMA	3W-GWO	1964 - 1967	26	56	178	2.10	SB	47	8.5	0.5	1.9	A	G	10.2	
LOUGH ERNE	3W-Y YR	1968 - 1972	50	48	162	1.54	SB	47	9.8	1.0	3.0	A	G	10.4	
LOUGH NEAGH	3W-Y	1968 - 1977	30	58	148	1.72	SB	46	10.4	1.2	2.6	B	VG	9.6	
LOUGHGALL	2W-P	1973 - 1977	42	39	216	2.66	S	44	9.7	3.0	2.8	A	G	9.0	
LOUIS D'OR	2Y-Y	1972 - 1972	60	34	214	2.09	SB	27	11.5	3.1	4.4	A	VG	9.0	
LOUIS PASTEUR	2W-GYO	1955 - 1963	-4	38	413	2.13	SB	64	11.0	1.6	4.0	A	VG	nd	
LOUISE DE COLIGNY	2W-O	1955 - 1963	36	36	304	1.86	SB	49	10.0	2.5	3.4	A	G	nd	
LOVE DREAM	3W-O	1955 - 1963	36	34	371	2.4	SB	52	9.0	1.2	3.6	A	G		
LOVE SONG	2W-O OY	1971 - 1987	15	33	162	2.31	B	37	10.6	2.3	5.2	A	G	7.8	
LOVENEST	2W-Y PP	1955 - 1963	-7	20	208	2.40	S	50	10.0	3.9	4.3	A	M	nd	
LUCCOMBE	2Y-O	1955 - 1963	-14	3	444	1.33	SB	40	12.0	2.9	3.5	B	G	nd	
LUCIFER	2W-Y O O	1955 - 1967	14	21	249	2.54	S	40	11.0	2.1	2.3	A	P	12.4	
LUDLOW	2W-W	1955 - 1963	29	15	184	1.87	S	42	10.0	3.7	3.9	B	VG	8.4	
LUNA MOTH	1Y-Y	1972 - 1981	13	13	110	2.03	S	36	12.8	5.2	4.4	A	VG	9.4	BS
LUNAR RAINBOW	2W-Y	1964 - 1967	17	27	273	1.30	S	29	12.0	3.0	4.9	A	G	7.4	
LUNAR SEA	1Y-W	1977 - 1981	4	16	108	1.94	S	37	11.2	4.7	4.4	A	G	10.7	BS

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
LUNAR SPELL	2Y-WWY	1977 - 1981	40	30	186	2.44	SB	38	10.4	3.6	4.0	B	G	10.6	B
LUNE DE MIEL	4W-Y	1955 - 1963	27	17	399	2.93	E	50	10.0	1.6	nd	A	P	nd	
LURGAIN	1Y-Y	1977 - 1977	7	14	230	2.29	S	27	10.5	4.5	4.2	A	VG	8.4	
LUXURY	1Y-Y	1955 - 1963	-10	15	335	1.87	S	43	10.0	4.4	5.2	A	M	nd	
LYN PARKER	1Y-Y	1976 - 1980	16	3	252	1.89	S	38	11.3	4.5	5.6	A	G	10.2	
LYON	2W-OOY	1964 - 1967	51	38	nd	nd	S	32	11.0	2.3	4.6	A	G	nd	
LYSANDER	2W-YYO	1972 - 1980	6	45	103	1.70	SB	39	10.4	1.6	4.0	B	VG	9.5	
M. J. BERKELEY	1Y-Y	1964 - 1967	-50	16	268	2.40	S	36	11.5	5.1	5.4	A	M	6.6	
MABEL TAYLOR	2W-WPP	1972 - 1972	53	24	148	0.93	S	35	11.0	3.1	4.6	B	G	9.0	
MACBEARER	2W-YYO	1955 - 1963	21	23	603	2.40	SB	48	9.0	1.7	3.5	B	M	nd	
MADEIRA	2Y-O	1972 - 1972	31	22	144	1.93	S	27	10.0	1.9	4.0	A	VG	11.0	V
MADELON	2Y-O	1964 - 1967	19	43	nd	nd	SB	29	8.5	1.6	4.5	B	G	7.0	
MADRIGAL	2W-WWY														
MAGHERALLY	2Y-O	1955 - 1980	13	24	219	2.29	SB	57	9.0	1.7	2.2	A	G	9.3	
MAGIC CIRCLE	2W-YYR														
MAGIC DAWN	2W-P	1976 - 1980	63	34	179	2.57	S	34	10.3	3.2	4.4	A	G	7.4	
MAGISTRATE	1Y-Y	1972 - 1981	9	-3	141	1.89	S	41	10.9	4.9	4.9	A	G	10.7	S
MAGNET	1W-Y	1955 - 1963	-11	12	211	2.40	S	38	11.0	4.7	5.0	A	M	nd	
MAGNIFICENCE	1Y-Y	1955 - 1982	1	-6	162	1.79	E	36	11.5	4.2	5.2	A	G	8.0	
MAHEE	1Y-Y														
MAHMOUD (16)	3W-R	1964 - 1967	26	41	201	3.60	SB	43	9.0	0.9	2.7	A	G	8.0	
MAID OF ULSTER	2W-YYR	1982 - 1984	57	40	nd	1.25	B	40	9.5	1.5	4.2	A	VG	nd	
MAIDEN'S BLUSH	2W-P	1964 - 1967	11	22	249	3.20	S	31	10.5	4.3	3.7	A	G	7.8	
MAVEROE	1W-W	1964 - 1967	49	24	132	3.10	S	30	9.5	3.9	3.5	A	G	7.4	
MAJARDE	2Y-O	1955 - 1963	0	20	300	2.40	S	48	11.0	2.2	4.5	A	M	nd	
MAJESTIC	1Y-Y	1974 - 1974	54	19	298	0.38	S	30	10.0	4.2	3.9	B	G	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
MAJOLICA	2Y-Y														
MAJORCA	2Y-YYO	1972 - 1973	49	26	132	1.75	S	38	11.0	2.5	3.4	A	VG	10.2	
MAJORETTE	1Y-Y	1977 - 1978	20	22	266	1.76	S	31	11.8	4.9	6.2	B	G	9.9	
MAKASSAR	2Y-Y	1955 - 1963	-7	22	208	1.87	S	46	11.0	3.0	4.2	A	nd	8.2	
MALHEUR	2W-Y	1972 - 1972	37	21	158	1.40	S	35	11.0	2.5	5.7	A	G	7.8	
MALONE	1Y-Y														
MALTA	2Y-O	1955 - 1967	0	25	227	1.88	S	50	10.0	1.7	2.4	A	VG	10.0	
MALVERN CITY	1Y-Y	1968 - 1984	22	-2	176	1.71	S	42	11.0	4.8	5.3	A	G	8.7	B
MALVERN GOLD	2Y-Y	1972 - 1987	-16	-4	206	2.36	S	44	11.2	3.3	4.6	A	M	9.9	B
MAMOUTH	3W-R	1973 - 1973	-5	31	239	2.25	S	41	8.5	0.9	2.8	A	G	10.0	
MANCHU	2W-Y	1955 - 1963	0	10	337	1.87	S	41	9.0	3.2	4.5	A	G	nd	
MANCO	2W-YYO	1964 - 1967	9	18	nd	nd	S	31	10.5	2.2	5.3	B	G	6.6	
MANDATE	1W-W	1964 - 1967	5	24	nd	nd	S	34	11.0	4.5	3.4	B	G	9.4	
MANDO	1Y-Y	1972 - 1979	-2	-9	161	1.86	S	42	10.7	3.9	4.8	A	G	7.7	
MANDOLIN	2W-GPP														
MANGOSTEEN	3Y-R	1955 - 1963	-20	21	280	1.87	SB	44	10.0	1.6	3.4	A	G	8.0	
MARA	9W-YYO	1976 - 1981	25	43	164	2.55	SB	54	9.7	0.8	2.2	A	G	8.0	
MARAVEL	1Y-Y	1964 - 1982	2	13	192	2.16	S	41	11.0	5.3	4.6	A	G	7.6	
MARCH MADNESS	2Y-R	1979 - 1985	5	18	249	2.40	S	49	10.9	2.7	4.2	A	G	7.8	
MARCH SUNSHINE	6Y-Y	1955 - 1967	0	0	204	2.60	S	26	10.0	2.4	1.8	B	M	7.8	
MARCOA	2Y-O	1976 - 1981	11	35	88	2.32	E	42	11.1	2.4	4.1	A	G	11.5	
MARCURA	2Y-R	1976 - 1982	27	14	139	2.09	S	40	10.3	1.9	3.0	B	G	10.9	
MARFORT	2	1976 - 1979	11	9	196	1.45	SB	42	10.6	2.5	3.5	A	G	9.8	
MARGARET MITCHELL	3W-YYR	1955 - 1987	4	33	211	2.16	SB	46	10.0	0.9	2.3	B	G	9.2	
MARGRIET	2W	1968 - 1980	23	20	189	2.24	SB	43	10.6	2.3	3.9	B	G	9.6	
MARIMBA (19)	2Y-YYO	1974 - 1985	-2	-6	137	1.82	S	43	10.9	2.5	3.6	B	G	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
MARION CRAN	2Y-O	1955 - 1963	-28	20	327	2.93	S	44	9.0	1.7	3.0	B	M	nd	
MARKANT	3W-R	1955 - 1963	7	17	358	1.33	SB	46	10.0	1.3	3.0	A	M	nd	
MARKET GEM	2W-Y	1955 - 1963	14	20	315	2.40	S	50	10.0	2.6	3.0	A	M	nd	
MARKET MERRY	3Y-O	1955 - 1982	4	16	247	2.49	S	48	10.0	2.2	3.9	A	G	7.0	
MARKETHILL	1Y-Y	1968 - 1978	40	38	144	1.86	S	35	10.3	4.8	4.1	A	G	6.8	S
MARKSMAN	2Y-O	1955 - 1963	-4	15	247	2.40	S	45	9.0	1.9	3.1	A	G	nd	
MARS	2Y-R	1974 - 1987	26	13	324	2.81	S	41	10.0	2.6	2.5	A	G	8.8	
MARSHAL TSOEKOF	2W-OOR	1964 - 1967	11	17	331	1.90	SB	39	10.0	2.0	5.2	A	M	7.4	
MARTHA WASHINGTON	8W-O	1955 - 1963	-54	26	265	2.93	SB	70	9.0	1.1	3.2	A	G	nd	
MARY ANN	2W-W	1977 - 1978	16	24	238	1.88	S	28	11.5	4.5	3.9	B	G	8.8	
MARY BLONK	2W-YOO														
MARY BOHANNON	2Y-O	1973 - 1980	5	16	142	2.11	S	33	10.2	3.0	4.7	A	VG	10.0	
MARY COPELAND	4W-O	1955 - 1983	12	30	194	2.28	SB	46	9.9	1.4	nd	A	G	7.9	
MARY HOUSLEY	2W-YYO	1955 - 1967	10	12	253	2.90	SB	41	10.0	1.7	3.8	A	G	7.5	
MARY ROOZEN	2Y-O	1955 - 1973	25	10	172	2.03	SB	39	10.0	2.2	3.4	A	G	9.1	
MASAI KING	2Y-O	1972 - 1980	-5	27	95	2.18	SB	40	10.3	1.8	3.0	A	G	9.0	S
MASAKA	3W-R	1968 - 1792	-8	33	150	1.41	SB	44	10.0	1.2	3.0	A	VG	8.9	S
MASKED LIGHT	2Y-O	1968 - 1982	20	20	162	2.53	SB	44	10.8	2.7	4.0	A	VG	10.0	
MASTER FLORIS	1W-Y	1955 - 1963	21	18	193	1.87	S	29	11.0	4.7	4.6	A	G	nd	
MASTER ROBERT	1Y-Y	1955 - 1963	0	17	181	1.87	S	38	11.0	4.8	4.1	A	G	nd	
MASTERPIECE	3W-OOR	1955 - 1963	-11	46	236	2.40	S	50	7.0	0.9	2.2	A	P	nd	
MATA HARI	2W-O	1955 - 1963	-7	8	223	1.87	SB	42	11.0	2.2	4.5	A	M	nd	
MATADOR	8Y-O	1976 - 1984	-30	-1	125	2.26	E	33	5.7	0.7	2.2	A	G	8.6	
MATAPAN (15)	3W-R	1968 - 1972	22	32	157	1.11	SB	40	9.2	1.0	2.9	A	VG	9.5	S
MATCHLESS	2W-O														
MATLOCK	2Y-R	1964 - 1967	7	24	nd	nd	S	40	12.0	2.3	3.4	A	VG	10.2	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
MAVISTON	2Y-Y	1964 - 1967	47	22	nd	nd	SB	30	10.5	3.9	4.5	B	G	9.4	
MAXIMUS SUPERBUS	1Y-Y	1955 - 1967	-30	1	264	2.02	E	46	10.0	4.4	4.0	A	G	9.0	
MAY MOLONY	2W-Y	1964 - 1967	49	36	nd	nd	S	38	9.5	3.1	3.1	B	G	7.6	
MAY MURIEL	2W-Y	1967 - 1987	33	13	175	2.38	S	37	11.2	2.6	5.4	A	M	8.5	
MAY QUEEN	2W-GYP	1982 - 1984	81	50	nd	2.13	S	26	10.5	2.7	3.7	A	G	nd	
MAYMYO	2Y-O	1968 - 1980	8	2	204	2.72	SB	44	10.0	2.3	3.7	A	G	8.4	B
MEDAILLON	2Y-O														
MEDALIST	2W-YPP	1976 - 1980	-5	23	191	2.14	SB	38	10.9	2.4	4.6	B	VG	7.8	
MEDINA	4W-W	1964 - 1967	49	58	nd	nd	SB	39	7.5	0.9	nd	A	M	10.0	
MEDUSA	8W-O	1955 - 1963	-28	21	299	2.40	SB	50	6.0	0.6	1.8	A	M	nd	
MEGA	9W-GYR	1964 - 1967	20	37	350	2.40	SB	40	8.5	0.5	1.9	A	G	10.6	
MELANITE	3W-YOO	1964 - 1967	33	28	nd	nd	SB	39	9.0	0.8	2.3	A	M	6.0	
MELISSA	1Y-Y	1977 - 1977	57	30	175	1.25	SB	29	10.0	4.3	3.0	A	G	nd	V
MELODY LANE	2W-P	1976 - 1980	40	34	224	2.53	S	40	9.9	3.8	3.4	B	G	8.8	
MELROSE	3W	1968 - 1973	31	34	154	2.94	SB	37	10.0	1.4	3.8	B	G	7.2	S
MENTON	2W-P	1955 - 1963	7	23	259	1.87	S	44	11.0	3.5	4.3	B	M	nd	
MERCATO	2W-YYO	1955 - 1980	-8	17	284	2.95	SB	44	10.0	1.7	4.0	A	VG	7.6	
MERIDIAN	2Y-YYO	1968 - 1971	62	36	140	2.40	SB	38	10.5	1.5	3.6	B	G	9.8	
MERIVALE	2W-W	1964 - 1967	26	17	nd	nd	S	33	9.5	3.5	3.9	B	M	5.6	
MERLIN	3W-YYR	1973 - 1980	15	41	182	1.94	SB	49	8.9	1.0	2.9	A	VG	9.6	
MESSENGER	2Y-O	1964 - 1967	5	29	242	2.70	SB	37	10.5	2.8	4.7	B	G	7.0	
METROPOLITAN	2W-O	1964 - 1967	51	46	nd	nd	SB	31	12.5	2.6	6.9	B	VG	7.0	
MEXICO	2Y-O	1955 - 1982	-1	11	267	2.65	S	40	11.0	2.1	2.9	A	G	10.0	
MEXICO CITY	2Y-O	1972 - 1972	0	30	141	1.25	SB	43	10.0	2.1	3.1	A	VG	10.0	
MIKADO	2W-O	1972 - 1975	16	43	279	1.00	SB	41	11.7	1.8	4.6	A	VG	8.8	
MILAN	9W-GYR	1964 - 1982	28	54	166	1.82	SB	44	8.0	0.7	2.6	A	VG	9.8	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
MILANION	1Y-Y	1955 - 1982	33	21	157	2.01	E	40	10.0	4.5	4.3	A	G	7.2	
MILK AND CREAM	2W-Y	1955 - 1963	7	7	305	1.33	E	46	12.0	2.6	5.9	A	VG	nd	
MILK AND HONEY	4W-Y	1955 - 1963	-4	12	369	1.87	S	38	10.0	2.0	nd	B	P	nd	
MILKMAID	2W-W	1955 - 1963	-14	15	293	1.87	S	46	12.0	3.4	3.7	B	G	nd	
MILLISLE	3W-W														
MINISTER TALMA	1Y-Y	1964 - 1967	5	17	nd	nd	S	31	10.5	5.1	5.1	A	M	8.0	
MIRABELLE	3W-YYO	1977 - 1978	14	36	235	1.92	SB	33	9.0	1.0	2.9	A	G	8.9	
MIRALGO	2Y-O	1979 - 1979	-9	26	521	2.86	S	35	9.5	2.6	2.5	B	G	10.4	
MISSOURI	2Y-O	1955 - 1982	-4	32	203	2.43	SB	53	12.0	2.3	4.0	A	G	9.3	
MISTRAL	11W-Y	1973 - 1980	44	10	148	2.33	S	32	10.3	3.3	8.1	A	G	8.2	V
MISTY MOON	3W-WOO	1955 - 1973	21	51	194	1.73	SB	41	8.0	0.7	2.3	A	G	9.1	
MITE	6Y-Y														
MITYLENE	2W-Y	1955 - 1967	18	24	276	2.65	SB	44	11.0	1.8	3.2	A	M	9.0	
MODESTA	11Y-Y														
MOINA	3W-WWO	1976 - 1978	20	44	129	2.10	SB	42	8.8	1.0	3.0	A	G	8.7	
MOIRA O'NEILL	1W-Y	1955 - 1981	17	21	176	2.16	S	38	10.0	4.4	3.7	A	M	7.3	B
MONACO	2W-O	1964 - 1967	-28	16	239	1.80	SB	42	10.5	1.9	3.7	A	G	8.2	
MONARCH	1Y-Y	1964 - 1967	-5	27	179	2.20	S	33	9.5	4.3	4.3	B	M	7.8	
MONEYMORE	2Y-R	1979 - 1981	22	24	220	1.50	SB	45	11.0	2.4	3.0	A	G	9.9	
MONIQUE	2W-OOR	1964 - 1981	32	21	192	2.96	SB	42	11.0	2.1	4.7	A	G	8.4	
MONKSTOWN	2Y	1976 - 1980	34	16	121	1.57	S	47	12.6	4.6	4.1	A	VG	8.1	
MONTE CASSINO	2W-GRR	1955 - 1963	-7	23	306	2.40	S	48	10.0	1.8	4.0	A	VG	nd	
MONTE RRICO	4W-O	1977 - 1980	-8	40	216	1.65	SB	51	11.0	1.3	nd	A	VG	10.5	BS
MOON BOY	2Y-Y	1964 - 1967	40	19	195	2.30	S	35	11.0	4.5	3.5	B	M	9.0	
MOON GODDESS	1Y-Y	1964 - 1979	17	14	153	2.04	S	36	11.5	5.5	5.6	A	G	8.4	V
MOON ORBIT (19)	1/2Y-Y	1977 - 1977	57	15	265	1.84	S	30	11.0	4.2	4.5	A	G	11.6	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
MOONDANCE	3W-W	1968 - 1971	83	35	176	1.70	SB	36	9.5	1.6	3.0	A	G	8.4	S
MOONGATE	6W-W	1976 - 1976	51	-3	-35	0.14	S	27	8.0	3.2	1.6	A	M	nd	
MOONGOLD	1Y-Y	1964 - 1967	18	30	180	1.70	S	38	10.5	4.6	4.4	A	G	7.2	
MOONIE	2W-GWW	1975 - 1975	48	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	V
MOONMIST	1Y-Y	1972 - 1981	13	4	172	2.18	S	38	11.8	4.7	5.0	A	VG	7.2	B
MOONRISE	1Y-Y	1955 - 1963	0	13	242	1.87	S	35	12.0	5.1	5.5	B	VG	nd	
MOONSHINE	5W-W	1955 - 1963	36	40	205	0.80	SB	24	7.0	2.2	2.4	A	P	nd	
MOONSPELL	2Y-W	1977 - 1980	-25	8	124	2.21	S	38	11.8	4.2	4.4	B	VG	10.0	BS
MOONSTRUCK	1Y-Y	1964 - 1976	4	8	145	1.80	S	39	12.5	5.8	5.6	A	G	9.8	B
MORAY	1W-W	1972 - 1972	31	22	192	2.08	S	29	13.5	5.6	4.6	B	G	8.8	
MOTHER CATHERINE	11W														
GRULLEMANS															
MOUNT HOOD	1W-W	1955 - 1987	35	22	204	1.94	S	35	12.0	4.7	5.2	A	VG	7.9	S
MOUNT PLEASANT (15)	2W-Y	1977 - 1987	50	44	186	2.57	S	43	11.5	3.1	4.9	A	G	8.5	
MOUNT TACOMA	2W	1955 - 1963	-7	20	334	1.87	SB	46	11.0	2.7	6.0	B	G	nd	
MOUNT VERNON	2W-P	1976 - 1980	18	33	239	1.64	SB	36	11.0	2.6	4.5	B	VG	9.8	S
MOUNTJOY	7Y-Y														
MOUNTNORRIS	2W-Y	1977 - 1987	6	37	227	2.59	S	41	9.9	2.7	4.0	A	M	9.1	
MOVILLE	2W-W	1964 - 1967	23	33	nd	nd	SB	42	12.5	2.5	4.3	A	G	9.2	
MOYLENA	2W-P	1972 - 1972	45	22	175	1.67	S	36	11.0	2.8	3.6	A	G	9.8	B
MOYLOUGH	3W-YR	1964 - 1967	20	32	nd	nd	SB	39	9.5	1.0	3.0	A	G	8.0	
MR JINKS	3W-YOO	1968 - 1981	18	42	152	2.26	SB	45	10.0	1.1	2.8	A	G	7.8	S
MRS DAVID CALVERT	3W-GRR	1977 - 1984	26	24	211	1.82	SB	44	9.8	1.4	3.7	A	G	9.7	
MRS E. C. MUDGE	1W-Y	1955 - 1963	52	37	86	1.33	S	40	10.0	4.9	4.7	A	G	nd	
MRS ERNST H. KRELAGE	1W-W	1955 - 1973	38	24	255	2.81	S	39	10.0	4.6	3.7	A	M	7.5	
MRS LANGTRY	3W-W	1955 - 1967	44	32	356	2.96	SB	52	9.0	1.7	2.0	A	P	10.5	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
MRS NETTE O'MELVENY	3W-GYY	1955 - 1963	-7	20	369	2.93	SB	48	10.0	1.3	3.1	A	M	nd	
MRS OSCAR RONALDS	2W-P	1968 - 1973	76	35	172	1.76	SB	33	9.5	2.9	3.5	A	G	10.1	
MRS R. O. BACKHOUSE	2W-P	1955 - 1967	24	32	222	2.30	SB	52	10.0	3.6	3.0	A	M	10.0	
MRS WILLIAM COPELAND	4W-W	1955 - 1963	-7	21	236	1.87	S	36	11.0	3.0	nd	A	M	nd	
MULATTO	1Y-WWY	1955 - 1963	0	1	200	1.87	S	36	10.0	3.5	3.8	B	M	nd	
MULRANY	2Y-Y	1964 - 1967	23	18	nd	nd	S	39	11.0	4.7	5.0	A	G	8.8	
MURMANSK	2W-W	1964 - 1967	53	45	nd	nd	S	36	10.5	3.8	4.5	B	VG	nd	
MUSCADET	2W-Y	1968 - 1987	32	14	205	2.56	S	38	10.2	2.5	5.1	A	M	8.2	
MUSIC HALL	1W-Y	1955 - 1963	14	12	193	1.87	S	38	11.0	5.2	5.5	A	G	8.2	
MUSTANG	2Y-YOO	1955 - 1963	7	15	278	1.87	SB	43	10.0	2.6	3.5	A	G	nd	
MY LOVE (17)	2W-Y	1968 - 1973	52	32	183	1.50	S	36	11.5	3.9	4.1	A	VG	9.0	BS
MY WORD	2W-P	1977 - 1987	1	8	159	2.29	S	40	9.4	3.5	3.8	A	M	8.6	V
MYRIANTHA	3W-WWR	1964 - 1967	18	30	nd	nd	SB	41	8.5	1.4	3.4	A	G	8.2	
MYSTERY	2Y-O	1977 - 1978	40	30	482	1.52	S	36	10.2	2.0	3.8	A	G	9.1	
MYSTIC	3W-GWO	1955 - 1967	12	45	176	2.20	SB	50	8.0	0.9	2.6	A	M	9.2	
NAIROBI	2W-O	1968 - 1973	78	32	170	1.36	SB	39	9.8	2.0	3.6	A	G	8.3	
NAMPA	1Y-W	1970 - 1986	18	15	167	1.86	S	40	11.5	5.1	3.9	A	VG	9.8	BS
NAMSOS	2W-W	1964 - 1967	17	9	256	2.30	S	30	10.5	3.7	5.0	B	G	7.6	
NANCEGOLLAN	7W-W														
NANCHISEL	2Y-O	1972 - 1972	-13	23	182	2.17	S	45	10.5	2.3	2.9	A	G	10.8	
NANNY NUNN	3Y-OOR	1955 - 1963	14	39	342	2.40	S	43	8.0	1.1	2.6	A	M	nd	
NANTUCKET	2W-O	1972 - 1973	38	30	194	1.42	S	43	9.5	1.4	4.2	A	G	10.0	S
NANTWICH	2W	1964 - 1967	51	51	nd	nd	S	35	7.5	1.2	2.5	A	VG	8.6	
NAOMI	2W-YYO	1955 - 1963	-7	32	291	1.87	S	50	10.0	1.7	3.1	B	VG	nd	
NARVIK	2Y-O	1964 - 1980	14	21	198	2.77	SB	49	9.5	2.0	2.9	A	G	10.0	
NAZARETH	2Y-W	1970 - 1987	6	13	201	2.84	S	38	9.4	3.3	3.4	A	M	10.5	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
NEARULA	3W-R	1972 - 1972	45	13	157	1.21	S	42	9.5	1.1	2.8	A	G	11.6	
NELLY	3W-Y	1955 - 1967	-7	30	226	2.25	SB	50	10.0	1.3	2.5	A	G	8.0	
NEMO	3W-YYO	1964 - 1967	47	29	218	2.10	SB	40	9.0	1.3	3.2	A	G	6.4	
NEVANA	2W-P	1976 - 1977	54	19	119	0.50	S	28	10.0	3.4	3.9	A	M	7.8	V
NEW GOLD	2Y-Y	1955 - 1982	-7	23	208	2.27	S	39	11.0	2.8	3.9	A	VG	9.5	
NEW HORIZON	1Y-Y	1964 - 1967	10	22	nd	nd	S	31	11.5	4.9	3.9	B	G	9.6	
NEWCASTLE	1W-Y	1979 - 1981	44	20	136	1.24	S	40	12.2	4.6	4.2	A	VG	11.5	B
NEWTON (19)	8Y-O														
NIMROD	2Y-Y	1968 - 1971	12	23	181	2.30	S	42	11.5	3.3	4.4	B	G	8.4	
NINTH LANCER	2Y-O	1964 - 1967	-21	24	nd	nd	S	34	11.5	2.5	3.8	A	G	10.2	
NIPHETOS	2W-W	1955 - 1967	15	10	238	2.28	SB	34	10.0	3.0	4.5	A	G	9.0	
NIRVANA	7W-W	1955 - 1967	-1	23	326	2.53	E	40	6.0	1.1	1.9	B	M	8.8	
NISSA	2W-Y	1955 - 1963	14	15	308	2.40	S	54	10.0	3.6	4.0	A	M	nd	
NIVETH	5W-W	1955 - 1967	7	27	264	1.82	E	28	7.0	2.2	2.9	B	M	9.0	
NOBLE DIAMOND	3W-GYR	1955 - 1963	66	51	215	1.33	SB	52	9.0	0.8	2.3	A	VG	nd	
NOBLESSE	2Y-O	1972 - 1972	45	29	125	1.79	nd	nd	nd	nd	nd	nd	nd	nd	
NODDY	2W														B
NOR-NOR	2Y-Y	1955 - 1963	0	8	268	1.87	SB	30	8.0	2.6	2.4	B	M	nd	
NORTH CAPE	2W-W	1964 - 1967	19	22	192	2.60	S	33	9.5	2.7	3.6	B	G	9.2	
NORTHERN LIGHT	2W-O	1972 - 1981	2	36	153	1.83	SB	40	11.9	2.0	4.1	A	VG	9.0	S
NORVAL	2W-O	1982 - 1984	66	40	nd	2.00	B	36	11.0	1.7	4.2	A	VG	nd	
NORWAY	1W-Y	1972 - 1973	16	16	168	1.20	S	32	11.0	5.0	5.4	B	M	9.3	
NUAGE	2W-W	1964 - 1973	22	11	142	1.98	S	32	11.0	4.5	3.9	A	G	6.6	
NURSEMAID	3W-O	1955 - 1963	-4	29	280	1.87	SB	50	10.0	1.0	2.7	A	M	nd	
NYORA	3W-YYR	1976 - 1977	-18	21	62	1.34	SB	41	10.0	1.2	2.9	A	VG	8.4	
N. CYPRI	10	1965 - 1969	-61	23	191	0.18	nd	nd	nd	nd	nd	nd	nd	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
N. JONQUILLA (ADS)	10Y-Y	1955 - 1963	-80	25	657	6.13	S	38	4.0	0.4	1.1	B	M	nd	
N. OBVALLARIS	10	1955 - 1967	7	-8	447	2.82	E	31	7.0	3.1	3.1	A	G	9.8	
N. OCHROLEUCUS	10														
(syn. TAZETTA OCHROLEUCUS)															
N. POETICUS	10W-YR	1955 - 1967	45	68	272	1.78	SB	58	7.0	0.4	1.7	A	G	10.6	
RECURVUS (ADS)															
N. POETICUS RADIIFLORUS	10W-GYR	1955 - 1967	24	46	336	2.75	SB	48	8.0	0.5	1.8	A	P	8.2	
POETARUM (ADS)															
N. PSEUDONARCISSUS	10Y-Y	1964 - 1967	34	9	332	1.90	S	26	11.0	4.7	3.9	A	P	7.2	
GAYI (ADS)															
N. TAZETTA LACTICOLOR	10W-Y	1955 - 1963	0	3	439	1.87	S	18	3.0	0.5	0.8	B	P	nd	
(syn. CANALICULATUS) (ADS)															
N. x MEDIOLUTEUS	10W-Y	1983 - 1985	-25	56	284	1.56	S	36	6.2	0.6	1.4	B	VG	nd	
(syn. BIFLORUS) (ADS)															
OCARINO (19)	4Y-Y	1979 - 1985	17	49	183	2.08	S	44	11.2	2.7	nd	A	G	8.3	
OCEAN MIST	1W-W	1976 - 1978	43	17	112	1.74	SB	33	11.0	4.5	4.3	A	G	7.1	S
ODENSE	2W-YYO	1971 - 1987	15	35	168	3.21	S	39	10.6	2.3	4.6	A	M	8.8	
OKLAHOMA	1W-Y	1968 - 1971	54	25	160	1.90	S	33	13.0	5.2	5.4	B	G	7.4	BS
OLDERFLEET	2W-P	1982 - 1982	94	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	V
OLIVER CROMWELL	1W-Y	1955 - 1963	-4	12	170	1.87	S	35	12.0	4.4	5.0	A	M	nd	
OLIVET	2W-W	1972 - 1980	-6	74	115	1.82	SB	45	11.9	3.0	5.1	A	G	10.2	
OLYMPIA	1Y-Y	1964 - 1967	5	30	190	1.20	S	35	11.0	5.3	5.6	A	G	8.0	
OLYMPIC FIRE															
2Y-R															
OLYMPIC GOLD	1Y-Y	1979 - 1985	34	34	163	1.56	S	35	11.3	5.0	4.4	A	VG	8.2	
OLYMPIC TORCH															
2W-YYO															
OMAROE	2W-Y	1977 - 1980	-10	4	240	2.16	S	39	11.2	3.6	4.0	A	G	9.2	VB

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
ORANGE BELL	2Y-O	1964 - 1967	20	24	nd	nd	SB	43	11.0	3.2	4.0	B	G	7.6	
ORANGE BIRD	2Y-R	1955 - 1963	14	17	221	1.87	SB	40	10.0	2.1	3.8	A	G	7.6	
ORANGE COCKADE	3W-O	1972 - 1972	31	33	263	1.33	SB	30	9.5	2.1	3.5	B	M	8.0	
ORANGE CREST	2Y-YYO	1972 - 1973	34	20	256	3.27	S	41	10.5	2.2	3.9	A	G	9.6	
ORANGE CUP	8Y-YYR	1964 - 1967	-45	14	nd	nd	S	21	5.0	0.6	1.6	B	M	7.9	
ORANGE FAVOURITE	2	1964 - 1967	-15	15	334	1.80	SB	36	9.5	2.5	4.2	A	G	8.2	
ORANGE FIRE	2W-O	1972 - 1972	45	46	70	0.83	SB	24	11.0	2.1	5.7	A	G	7.4	
ORANGE FLAG	2W-O														
ORANGE FRILLED	2Y-O	1964 - 1967	24	12	nd	nd	SB	34	10.0	2.9	4.1	A	G	7.6	
ORANGE GLORY	6Y-O	1955 - 1963	14	5	334	2.93	S	36	8.0	3.4	3.5	A	M	nd	
ORANGE GLOW	2Y-O	1955 - 1963	-4	18	264	2.40	SB	42	10.0	2.6	4.4	A	P	nd	
ORANGE MASTER	2Y-O	1964 - 1967	39	28	nd	nd	SB	37	10.0	3.9	4.6	A	G	9.0	
ORANGE MONARCH	2W-YOO	1964 - 1967	34	27	nd	nd	SB	33	11.0	2.4	5.9	B	VG	6.2	
ORANGE QUEEN	7Y-Y	1955 - 1963	-45	5	227	2.40	S	30	6.0	1.4	1.3	B	P	nd	
ORANGE SHERBERT	2Y-YOY	1984 - 1984	44	49	23	1.25	nd	33	nd	nd	nd	nd	nd	nd	
ORANGE STANDARD	2Y-O	1968 - 1972	18	12	236	2.04	S	31	12.5	2.9	3.8	B	G	9.0	
ORANGERY	11W-POY	1976 - 1982	24	24	184	2.21	S	36	9.6	2.4	6.5	B	G	8.1	
ORANIQUE	2W-YYO	1964 - 1967	51	33	nd	nd	SB	35	10.0	1.5	3.7	A	G	7.0	
ORANJE BRUID	2W-GOO	1955 - 1963	29	25	215	2.40	S	42	10.0	1.8	4.7	A	VG	nd	
ORANJE VAAN	2Y-O	1955 - 1963	-7	17	342	1.87	SB	42	11.0	2.4	3.8	A	G	7.2	
ORATORIO	2W-Y	1977 - 1987	4	16	208	2.47	S	42	10.8	2.9	4.5	A	G	9.4	
ORIENTAL RAY	2Y-YYO	1972 - 1973	26	22	140	2.10	S	36	11.2	3.0	5.4	A	G	9.8	
ORISSA	3Y-YYO	1964 - 1967	-2	33	216	2.50	S	42	8.5	1.3	3.0	B	G	8.0	
ORMEAU	2Y-Y	1968 - 1982	20	32	160	2.26	S	36	10.8	4.3	3.6	A	G	8.6	S
ORNATUS	10W-YYR	1964 - 1967	31	33	244	2.30	S	34	7.0	0.4	1.5	B	P	9.4	

(syn. POETICUS ORNATUS) (16)

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
ORNATUS MAXIMUS (16)	9W-GYR	1964 - 1967	20	30	194	1.40	SB	28	7.5	0.4	1.8	A	G	9.8	
ORONTES	2Y-O	1964 - 1967	5	7	nd	nd	S	44	9.0	2.0	3.1	B	G	9.0	
ORPHEUS	2W-PPY	1979 - 1979	-2	29	207	2.50	S	27	10.5	3.5	3.2	B	G	nd	
OSIRIS	1Y-Y	1984 - 1985	-5	-2	186	2.10	S	34	11.8	5.2	5.3	A	G	7.0	
OSLO	2W-W	1964 - 1967	40	24	nd	nd	SB	24	10.0	2.5	3.4	B	G	9.4	
OSPREY	2W-Y	1955 - 1963	27	40	381	2.13	SB	48	11.0	1.6	3.0	A	M	nd	
OSWEGA	2W-Y														
OTRANTO	3W-YYO	1973 - 1973	9	36	199	2.29	S	46	10.5	1.7	2.2	A	VG	14.0	
PAINTED LADY	2W-O	1955 - 1963	14	31	199	1.87	SB	53	10.0	2.1	3.1	A	G	nd	
PALAVERER	1W-Y	1974 - 1987	18	6	135	2.53	S	41	11.6	4.8	4.1	A	G	9.8	S
PALESTINE	2Y-YOO	1964 - 1967	34	32	178	1.80	SB	44	9.5	1.8	2.6	A	G	8.4	
PALMY DAYS	2Y-YOO	1970 - 1970	28	14	288	1.80	SB	40	12.0	3.0	4.1	A	M	10.8	
PAPILLON BLANC	11W-W	1964 - 1982	43	35	238	2.27	SB	39	11.0	2.5	5.8	A	G	8.0	
PAPRIKA	3W-R	1968 - 1973	50	36	212	2.41	SB	44	9.5	1.3	3.2	A	G	8.1	
PAPUA	4Y-Y	1976 - 1983	-11	33	147	1.97	SB	44	10.2	2.2	nd	A	G	8.7	B
PARACUTIN (15)	2Y-R	1972 - 1972	60	23	159	1.42	S	32	9.5	1.5	4.4	A	VG	9.4	
PARAMARIBO	2Y-R	1964 - 1967	37	22	253	2.30	SB	38	9.5	2.3	4.1	B	G	6.0	
PARCPAT	7Y-O	1955 - 1963	-17	17	317	5.06	E	44	8.0	1.8	2.5	B	M	nd	
PARISIENNE	11W-O														
PARKMORE	2W-W	1964 - 1967	45	9	123	1.90	S	33	11.0	4.2	3.7	A	VG	7.8	B
PARSIFAL	2W-R	1979 - 1985	26	41	270	1.32	S	40	10.8	1.6	4.0	A	VG	7.4	
PASSIONALE	2W-P	1977 - 1981	71	36	179	2.59	S	33	10.9	3.4	4.1	A	G	9.4	
PATAGONIA	2Y-O	1967 - 1987	-8	30	170	2.74	S	45	10.8	2.2	3.3	A	G	10.6	BS
PATHAN	2Y-O	1964 - 1967	47	39	190	2.00	SB	46	11.0	1.7	3.8	A	G	8.6	
PATRIA	1W-Y	1964 - 1967	40	11	nd	nd	S	27	11.5	5.5	4.9	A	G	7.6	
PATRIARCH	2W-O	1955 - 1963	-4	22	317	1.87	SB	52	12.0	2.0	4.8	A	VG	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
PATRICIA	4W-R	1973 - 1975	83	89	189	0.78	SB	42	7.5	0.5	nd	A	G	13.0	
PAUL BUNYON															
PEARL SCHUBERT	2W-Y	1977 - 1979	10	9	148	2.09	S	38	11.8	4.7	4.5	A	G	9.0	B
PEDRO	1Y-Y	1955 - 1963	10	8	199	1.33	S	44	8.0	4.1	3.9	A	M	nd	
PEEPING TOM	6Y-Y	1955 - 1967	17	0	210	3.15	S	34	10.0	4.4	2.9	A	M	11.8	
PEERLESS GOLD															
PEKING	1Y-Y	1955 - 1963	10	2	124	1.87	S	32	12.0	4.6	5.0	B	G	nd	
PEMBRAZE	1Y-Y	1955 - 1963	-7	8	265	1.87	S	41	12.0	4.9	4.1	B	G	nd	
PENBEAGLE	2Y-Y	1955 - 1963	-14	7	299	2.40	E	40	10.0	3.4	4.7	A	VG	7.0	
PENBEARITH	1Y-Y	1973 - 1979	14	12	144	2.03	S	40	11.3	4.8	4.3	A	G	9.9	S
PENCOYS	2Y-YOO	1955 - 1963	-7	15	318	1.87	SB	45	9.0	1.8	3.2	A	G	7.0	
PENCREBAR	4Y-Y	1964 - 1967	11	16	275	1.80	S	22	5.5	1.7	nd	A	M	7.5	
PENGORNIN	2W-Y	1973 - 1973	10	13	72	2.80	S	40	11.0	4.2	4.5	A	M	10.8	S
PENPOL	7Y-Y	1955 - 1967	-44	10	260	2.90	S	46	8.0	1.2	2.4	B	M	9.4	
PENQUITE	2Y-O	1955 - 1967	1	13	186	1.84	SB	40	9.0	1.8	3.1	A	G	7.6	
PENRYN	2W-W	1968 - 1980	30	35	158	2.30	SB	41	10.5	2.0	3.6	A	VG	8.5	
PENSIVE															
PENTEWAN	2Y-Y	1955 - 1987	-9	4	244	2.68	S	47	10.2	3.0	3.5	A	G	9.7	
(?syn. CALIFORNIA)															
PENTREATH	3Y-O	1964 - 1967	-2	14	338	3.00	S	36	9.0	2.2	3.0	B	M	7.6	
PENVER	2Y	1972 - 1972	-25	20	186	2.08	S	40	12.0	3.0	4.0	B	G	12.4	B
PENVOSE	2Y-Y	1955 - 1963	7	15	263	2.40	S	54	10.0	3.8	4.6	B	G	nd	
PEPPER	2Y-O	1955 - 1963	-11	0	461	2.93	S	32	9.0	2.0	2.5	A	M	nd	
PER LAGERQUIST	2Y-YYO	1973 - 1973	-13	18	200	1.75	S	34	8.5	2.0	3.5	B	VG	9.6	S
PERA	3W-R	1964 - 1967	13	39	181	1.90	SB	39	9.0	0.9	2.6	B	G	9.4	
PERDITA	9W-GYR	1976 - 1981	21	52	153	2.20	SB	51	8.3	0.6	2.0	A	G	9.1	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
PERICLES															
1W-W															
PERIDOT	2W-GYY	1972 - 1977	55	48	183	2.10	S	45	9.0	1.4	3.0	B	G	11.0	
PERIMETER	3Y-YYO	1972 - 1981	22	24	132	2.18	S	49	9.6	1.2	2.9	A	G	10.5	BS
PERSONALITY															
2W-Y															
PERSUASION	2W-OOY	1972 - 1980	8	28	79	1.52	SB	36	10.6	1.9	4.1	A	G	8.7	B
PERU	2Y-O	1973 - 1973	-5	21	152	1.83	S	45	10.5	2.0	3.8	A	VG	10.6	
PETER PIPER	2Y-O	1964 - 1967	-5	21	nd	nd	S	32	7.5	1.4	2.3	B	G	8.8	
PETERBOROUGH	2Y-Y	1964 - 1980	26	24	180	2.48	SB	41	10.0	2.4	4.2	B	VG	7.0	
PETILLANT	3W-YYO	1972 - 1972	0	34	148	0.71	SB	35	9.0	1.0	3.0	A	G	10.0	
PETSAMO															
1W-W															
PHILALETUS	2W-WWY	1964 - 1967	-5	18	nd	nd	S	34	9.0	2.4	4.2	A	G	8.0	
PHILLUS CHIDGEY (17)	2Y-Y	1977 - 1980	7	22	269	2.36	S	46	10.6	4.0	4.1	A	G	10.2	
PICADOR	3W-GOO	1955 - 1963	7	39	338	1.87	S	44	8.0	1.0	2.9	B	G	nd	
PICK ME	2Y-O	1977 - 1980	9	9	260	2.11	S	42	10.7	2.4	3.7	B	G	10.9	
PICK UP	11Y-O	1976 - 1982	20	25	170	1.97	SB	37	10.0	2.6	6.8	B	G	8.5	S
PICKWICK	2W-YYO	1970 - 1987	41	41	150	2.22	B	44	10.1	1.8	3.4	A	G	8.8	
PIERROT	3W-R	1968 - 1972	56	45	130	1.09	SB	37	10.0	1.3	2.8	A	G	8.3	S
PIGEON	2W-W	1964 - 1967	13	45	nd	nd	SB	34	9.0	2.0	3.6	A	VG	8.6	
PILGRIM	2W-Y	1964 - 1967	10	18	nd	nd	S	34	9.0	1.0	2.0	B	P	7.4	
PILGRIMAGE	2Y-Y	1955 - 1963	-14	12	352	2.40	E	37	10.0	3.1	3.6	B	VG	nd	
PILOT LIGHT	2Y-O	1968 - 1980	15	40	153	2.12	SB	41	9.9	1.8	3.2	A	G	8.0	S
PINDAR	3W-GWP	1964 - 1967	-9	42	nd	nd	SB	42	8.0	0.7	2.2	A	G	9.8	
PINK BEAUTY	2W-P	1964 - 1967	56	36	194	2.10	SB	30	11.5	3.7	3.3	A	VG	10.4	
PINK CHIFFON	4W-P	1977 - 1984	10	20	148	1.86	S	45	9.8	2.3	nd	A	G	8.4	S
PINK FANCY	2W-P	1955 - 1963	29	31	268	2.93	S	40	10.0	2.7	3.6	B	M	nd	
PINK GIN	2W-GPP	1976 - 1977	57	20	120	1.35	E	29	10.5	3.6	3.8	A	G	7.9	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
PINK GLORY	2W-P	1955 - 1963	39	37	292	1.87	SB	35	12.0	3.4	4.4	A	G	nd	
PINK ISLE	2W-P	1964 - 1967	-21	25	nd	nd	S	30	10.5	2.9	3.1	B	G	9.2	
PINK PACER	2W-P	1976 - 1987	-8	1	236	2.96	S	39	10.5	3.4	2.9	A	M	7.8	
PINK PROMISE	2W-P	1972 - 1980	11	8	89	1.43	S	36	11.2	3.8	3.8	B	G	7.9	B
PINK RIM	2W-YYP	1955 - 1981	30	24	170	2.28	S	42	10.0	2.6	3.8	A	G	9.6	
PINK SELECT	2W-OOP	1964 - 1967	5	32	nd	nd	S	34	9.5	3.1	4.1	B	G	7.8	
PINK SMILES	2W-P	1968 - 1982	46	24	200	2.22	SB	42	12.0	4.0	5.3	A	VG	7.3	
PINK SPRITE	2W-P														
PINKBEN	2W-P	1955 - 1963	4	24	252	2.93	S	46	9.0	3.1	2.8	A	VG	nd	
PINOCCHIO	2W-GOO	1955 - 1963	-4	20	186	2.40	S	58	9.0	1.7	3.1	B	VG	nd	
PINWHEEL	2Y-Y														
PINZA	2Y-YOR	1977 - 1987	-5	15	187	2.57	E	46	10.0	2.2	2.6	A	M	11.1	S
PIONEER	1W-Y	1976 - 1981	7	0	87	1.68	S	35	11.0	5.0	5.1	A	G	9.8	
PIRATE KING	2W-O	1970 - 1987	11	28	166	2.24	B	47	10.9	2.0	2.8	A	G	9.1	
PIXIE'S PARLOUR	3W-Y	1977 - 1987	34	31	203	2.61	B	45	9.6	1.2	2.6	A	M	8.0	V
PLANET	4Y-Y	1980 - 1985	1	10	122	2.22	S	46	12.7	4.2	nd	A	G	9.9	BS
PLANTINAL	3W-GWW														
PLAYBOY	2Y-O														
PLEASANT (18)	2Y-R	1955 - 1982	-19	8	227	2.44	SB	44	10.0	2.8	3.8	A	G	9.4	
PLEASING	2Y-Y	1955 - 1963	-7	18	374	1.87	SB	48	9.0	3.2	3.3	A	G	8.2	
PLENUS	4W-W	1955 - 1971	54	61	231	1.90	S	53	8.0	nd	nd	A	G	11.2	
(syn. POETICUS FLORE PLENO)															
PLUVIUS	2Y-O	1964 - 1967	40	39	188	2.60	S	34	12.5	4.2	4.2	B	G	8.6	
POINT BARROW	1W-W	1973 - 1973	64	24	202	3.00	nd	nd	nd	nd	nd	nd	nd	nd	B
POLAR ICE	3W-W	1955 - 1963	14	37	278	1.87	SB	50	9.0	0.8	2.2	A	M	nd	
POLAR SEA	3W-GWW	1964 - 1967	49	55	nd	nd	S	42	8.0	0.6	2.0	A	M	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
POLGLASE	8W-R	1955 - 1963	-31	15	320	1.87	SB	44	7.0	0.9	1.7	A	M	nd	
POLINDRA	2W-Y	1955 - 1981	-12	18	210	2.40	SB	58	11.0	2.9	4.0	A	G	7.0	B
POLNESK	7Y-Y	1955 - 1967	-16	22	240	3.02	SB	50	8.0	1.2	2.1	B	G	8.0	
POMONA	3W-GYO	1955 - 1963	0	17	300	2.40	SB	48	10.0	1.4	3.7	A	M	nd	
PONDEROSA	1Y-Y	1973 - 1987	15	16	145	2.40	S	43	11.1	4.8	5.2	A	M	10.7	S
PONTRESINA	2W-Y	1972 - 1980	6	34	172	1.99	S	42	10.1	1.8	4.7	B	G	9.3	
PORTADOWN	1Y-Y	1976 - 1981	35	23	135	1.93	S	41	10.8	4.6	4.1	A	G	8.2	B
PORTAL (15)	2W-WPP	1976 - 1980	10	22	212	2.08	S	43	11.1	2.8	5.2	A	G	8.9	S
PORTARLINGTON	1Y-Y	1964 - 1981	2	22	188	1.92	S	40	10.0	4.2	4.6	A	G	7.0	
PORTCHAPEL (16)	7Y-O	1973 - 1980	-34	13	198	3.31	S	37	7.5	1.6	2.1	B	G	9.9	
PORTHGWARRA	2W-Y	1973 - 1973	43	27	195	2.25	S	47	10.0	4.2	4.3	B	G	12.4	
PORTHILLY	2Y-O	1955 - 1967	-14	16	372	2.65	SB	46	10.0	1.9	3.2	A	M	9.2	
PORTMARNOCK	2Y-Y	1968 - 1971	51	33	184	1.80	S	38	11.0	3.1	4.9	A	G	9.2	S
PORTRUSH	3W-W	1964 - 1967	49	51	205	1.50	SB	50	8.5	0.9	1.9	A	VG	7.4	
POSTMISTRESS	2W-R	1967 - 1987	-8	31	165	2.30	B	43	10.7	1.5	2.6	A	G	8.3	S
PREAMBLE	1W-Y	1968 - 1971	75	25	216	2.30	S	33	11.5	4.6	4.2	A	VG	9.6	S
PRECENTOR	1Y-Y	1973 - 1977	7	13	212	1.78	S	45	10.7	4.6	5.0	A	G	9.5	S
PREFIX	6Y-Y	1976 - 1976	41	-3	56	0.36	S	23	7.5	2.7	2.2	A	G	nd	V
PRESIDENT LEBRUN	1W	1955 - 1963	-4	15	196	1.33	S	36	10.0	4.5	5.0	B	VG	nd	
PRESTIGE	1W-W														B
PRETORIA	1Y-Y	1955 - 1963	-7	17	168	1.87	S	38	12.0	4.9	4.9	A	VG	nd	
PRIDE OF CORNWALL	8W-YR	1955 - 1963	-28	17	231	2.40	SB	48	8.0	0.7	1.9	A	M	nd	
PRIDE OF ERIN	3W-YR	1964 - 1967	0	41	nd	nd	SB	46	10.0	1.0	2.9	A	VG	9.8	
PRIMROSE BEAUTY	4Y	1955 - 1982	23	41	214	1.96	E	50	6.0	0.6	nd	A	G	9.2	
PRIMROSE PHOENIX	4Y-Y	1955 - 1963	-4	17	171	1.87	SB	39	8.0	1.4	nd	A	P	6.2	
PRINCE REGENT	2Y-O	1964 - 1967	17	33	176	2.30	S	47	10.0	2.2	3.5	B	G	9.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
PRINCEPS	1W-Y	1955 - 1967	20	8	304	2.40	S	30	12.0	4.8	4.0	A	P	9.2	
PRINCESS BEATRIX	2W-R	1968 - 1981	18	28	155	2.15	SB	32	10.2	1.6	4.0	A	G	7.5	B
PRINCESS OF WALES	3W-WWO	1955 - 1963	7	20	259	2.40	SB	42	9.0	1.0	2.9	A	M	nd	
PRINCESS VICTORIA	2Y	1964 - 1967	-21	15	nd	nd	SB	47	11.0	2.1	3.7	A	M	9.2	
PRINCIPAL	1Y-Y	1955 - 1963	36	24	138	1.33	S	44	10.0	4.7	4.2	A	M	nd	
PRINS CARNAVAL	11Y-OOY	1973 - 1973	57	32	124	1.79	S	35	10.0	2.1	3.8	A	M	10.0	
PRIORITY	1Y-Y	1955 - 1972	7	12	166	1.58	E	37	10.0	4.1	4.5	A	VG	8.9	
PRISK	7W														
PRIVATEER	3W-O	1972 - 1973	39	30	152	2.57	SB	34	11.0	1.4	3.4	A	G	8.9	
PROFESSOR EINSTEIN	2W-R	1964 - 1967	40	24	nd	nd	SB	28	10.5	1.6	4.9	A	G	8.6	
PROFILE	2W-Y	1976 - 1987	0	41	131	2.18	S	49	11.1	2.9	3.7	A	M	10.1	
PROLIFIC	1Y-Y	1983 - 1983	36	17	nd	2.00	S	36	12.5	4.8	6.1	nd	G	7.0	
PROLOGUE	1W-Y	1973 - 1987	34	5	138	2.70	E	36	10.5	4.5	3.5	A	M	8.7	
PROMPTITUDE	2W-Y	1964 - 1973	21	10	182	1.60	S	36	10.5	3.0	3.8	B	G	10.0	
PROSPERO	3W-R	1968 - 1972	60	42	128	0.92	SB	39	8.8	1.0	3.0	A	G	8.8	
PROVERB	2Y-Y	1972 - 1976	-6	24	123	2.44	S	42	11.0	3.7	4.8	B	G	7.8	VS
PROWESS	2W-Y	1976 - 1980	1	40	175	2.28	SB	41	10.5	1.9	4.6	B	G	9.2	
PTOLEMY	1W-Y	1973 - 1975	28	-3	232	1.21	S	40	11.2	4.9	4.7	A	G	6.9	
PURITY	2W-W	1972 - 1972	66	34	145	1.10	S	32	11.0	3.3	3.9	A	VG	9.8	
PYTCHLEY	3W-R	1964 - 1967	37	36	211	1.80	SB	45	9.0	1.0	2.6	A	VG	7.4	
QUEEN JULIANA	1W-W	1972 - 1972	31	26	97	1.73	S	34	12.5	5.4	5.5	A	G	10.6	
QUEEN OF BICOLORS	1W-Y	1955 - 1981	-4	14	196	1.78	S	44	11.0	4.3	5.0	A	G	8.4	B
QUEEN OF HEARTS	3W-O	1955 - 1963	17	23	382	1.33	E	51	8.0	1.0	2.1	A	M	nd	
QUEEN OF NARCISSI	3W-YYR	1955 - 1980	-4	33	270	2.39	SB	58	11.0	0.9	2.6	A	VG	9.0	
QUEEN OF STARS	3W-YYR	1955 - 1963	-14	24	354	2.40	SB	52	10.0	1.1	2.9	A	G	nd	
QUEEN OF THE NORTH	3W-Y	1955 - 1963	-20	17	295	2.93	SB	47	9.0	1.2	2.0	A	M	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
QUEEN OF ULSTER	1W-Y	1964 - 1967	69	44	nd	nd	S	30	11.0	5.1	5.0	B	G	7.0	
QUEEN'S COUNSEL	1W-W														B
QUEEN'S LADY	1W	1964 - 1967	61	33	nd	nd	S	29	10.0	4.8	4.1	A	G	nd	
QUEENSLAND	2W-P	1968 - 1982	68	36	167	2.35	SB	34	10.0	2.7	3.7	A	G	9.7	B
QUEBETTA	3W-O	1964 - 1980	5	33	246	2.23	SB	45	10.0	1.4	2.9	B	G	9.0	
QUIRINUS	2Y-O	1972 - 1984	25	35	169	2.69	SB	47	11.2	2.2	3.6	A	G	10.4	BS
RADIATION	2W-P														
RAFFREY	2W-WWY	1977 - 1980	31	28	199	1.98	S	34	10.0	2.4	4.0	B	G	9.3	S
RAINBOW	2W-WWP	1979 - 1985	17	35	197	2.18	S	36	10.8	2.5	4.0	A	VG	9.1	
RAMESES	2W-R	1979 - 1985	2	34	244	1.97	S	46	11.2	1.9	3.3	A	G	9.6	
RAMILLIES	2Y-O	1955 - 1963	0	15	296	1.87	SB	46	10.0	1.9	3.8	A	M	nd	
RAMOAN	3W-GYO	1972 - 1972	60	-1	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
RANGATIKEI	2	1976 - 1977	29	16	294	1.90	SB	42	10.8	3.4	3.9	A	G	8.5	
RAPALLO	3Y-YR	1964 - 1967	40	22	179	2.50	SB	38	8.5	1.2	3.4	A	G	8.0	
RASHEE	1W-W	1970 - 1987	38	38	158	1.97	S	38	11.8	4.6	4.3	A	G	10.6	S
RATHAVAN	2W-GWW														
RATHGAR	2W-WWY	1972 - 1980	34	43	118	1.90	S	42	11.0	4.1	4.7	B	G	9.5	
RATHKENNY	1W-Y	1955 - 1982	15	35	205	2.52	S	41	10.0	4.4	4.0	B	G	7.4	
RATHOWEN FLAME	2Y-O	1982 - 1982	82	-23	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
REALM	2Y-Y	1976 - 1981	6	2	155	2.09	SB	39	10.8	3.1	5.0	A	G	9.7	
RED ABBOT	2W-O	1955 - 1963	17	37	399	2.40	SB	51	10.0	1.5	2.6	A	G	nd	
RED APRIL	2W-R	1964 - 1973	24	25	228	2.12	SB	43	10.5	2.2	3.1	A	G	7.8	
RED BAY	2Y-R	1981 - 1981	39	54	278	2.44	SB	46	10.5	1.9	3.0	nd	G	10.6	
RED BEACON (16)	3W-O	1955 - 1963	-4	26	271	1.87	SB	41	8.0	0.9	2.3	A	P	nd	
RED BIRD	2W-O	1955 - 1963	-4	35	342	1.87	SB	48	11.0	1.7	3.3	A	G	nd	
RED BUTTON	2Y-O	1968 - 1973	19	32	131	2.18	S	41	10.2	1.8	3.6	A	G	8.1	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
RED COTTAGE	2W-YJR	1973 - 1987	-5	22	206	2.39	B	43	9.7	1.3	2.7	A	M	9.5	
RED CROSS	2Y-YOO	1955 - 1963	14	23	351	2.40	SB	42	11.0	2.5	3.9	A	M	nd	
RED DEVON	2Y-O	1955 - 1987	15	15	194	2.63	SB	44	11.8	2.9	3.9	A	VG	7.6	
RED GAUNTLET	3W-O	1955 - 1963	-14	43	334	1.60	EB	44	9.0	1.2	2.4	A	P	nd	
RED GOBLET	2Y-O	1955 - 1981	24	22	279	2.33	E	48	9.0	2.5	1.9	A	VG	11.4	
RED GUARD	8YO-O	1955 - 1967	2	34	212	2.95	S	62	8.0	0.8	1.7	A	M	8.9	
RED HACKLE	2W-YOO	1955 - 1963	-4	40	275	1.87	SB	53	11.0	1.5	2.6	A	G	nd	
RED IDOL	2Y-R	1968 - 1981	27	33	226	2.00	SB	46	10.5	1.9	2.6	A	VG	11.0	
RED LAKE	3W-OOR	1972 - 1974	49	47	190	2.24	SB	41	10.3	0.8	3.4	A	G	10.7	
RED RANGER	2Y-O	1964 - 1982	14	28	237	2.30	SB	49	11.0	1.8	3.9	A	G	10.4	B
RED RASCAL	2Y-R														
RED RIBAND	2Y-YJR	1968 - 1981	40	30	216	2.27	SB	43	11.2	2.3	3.6	A	G	7.2	
RED RIM	9W-YJR	1955 - 1967	46	40	272	2.35	SB	51	8.0	0.6	2.4	A	G	12.4	
RED SATIN	2W-R	1975 - 1975	27	4	nd	nd	SB	35	9.5	2.5	2.9	A	G	7.2	V
RED SQUIRREL	2Y-O	1968 - 1982	14	42	222	3.04	SB	42	10.5	2.2	3.0	B	G	7.8	
RED SUNRISE	2Y-O	1983 - 1983	5	21	nd	1.80	S	36	13.0	3.6	3.5	nd	G	nd	
REDMAN	2Y-R	1984 - 1984	0	-2	187	1.50	S	33	nd	nd	nd	nd	nd	nd	
REDMARLEY	2Y-O	1955 - 1963	0	35	334	1.87	SB	46	11.0	3.1	3.8	A	M	nd	
REDSTART	3W-GWP	1973 - 1980	39	40	221	2.47	SB	44	10.7	1.4	3.4	B	G	9.6	
REGALITY	3W-YOO	1968 - 1982	34	48	166	2.18	SB	48	11.5	1.1	2.8	A	VG	8.4	S
REGINALD DIXON	2Y-O	1955 - 1963	14	21	208	2.40	SB	50	10.0	1.9	3.7	A	G	nd	
REMBRANDT	1Y-Y	1955 - 1982	-2	8	244	2.36	S	36	11.9	5.2	5.5	B	G	9.3	
RENE DE CHALONS	2Y-YOO	1955 - 1963	-4	29	244	2.40	SB	52	12.0	3.5	5.0	A	M	7.0	
RENEE DONALDSON	2Y-O	1972 - 1972	53	29	186	2.04	SB	33	11.5	1.7	4.4	A	G	8.4	
REPRIEVE	3W-GWY														
REPROOF	2Y-R	1974 - 1975	40	31	355	1.59	S	37	10.5	2.2	3.2	A	G	9.1	V

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vascular life (days)	Disease notes
RESERVE	2Y-O	1968 - 1973	46	46	176	2.51	SB	36	8.8	1.9	2.6	A	G	9.2	B
RETREAT	2Y-R														V
REVELRY	2Y-O	1955 - 1982	26	27	288	2.31	SB	50	10.0	2.1	3.2	A	VG	9.7	
REWA	2W-O	1955 - 1963	17	15	299	1.87	SB	57	9.0	2.2	4.0	A	G	nd	
REYNOLDSTOWN	2Y-O	1955 - 1963	-20	17	268	2.40	S	56	10.0	2.0	2.3	A	M	nd	
RIBER	1W-W														
RICHARD TAUBER	8W-Y	1955 - 1963	-14	16	15	1.33	S	36	7.0	0.8	2.0	A	G	nd	
RICHHILL	2Y-YYR														
RIDEAU HALL	3W-YYO	1955 - 1973	14	38	277	2.27	SB	54	10.0	1.2	3.0	A	G	7.1	
RIESENBECK	2W-YYO	1955 - 1967	-7	24	314	2.08	SB	42	10.0	2.4	4.9	A	G	7.8	
RIJNVELD'S EARLY SENSATION	1Y-Y	1955 - 1984	-7	-13	192	1.42	E	27	10.1	4.0	4.6	A	M	7.3	
RIMA	1W-P	1974 - 1987	37	24	192	2.22	S	37	11.2	4.6	4.3	A	P	7.7	S
RING O' ROSES	2Y-YYR	1955 - 1963	-11	34	290	1.87	SB	56	10.0	2.0	3.4	A	G	nd	
RINGMASTER	2Y-YYO	1972 - 1977	52	24	118	1.98	S	34	9.5	1.8	4.0	A	G	9.6	V
RINSEY	3W-W	1964 - 1967	47	35	166	2.20	SB	34	9.5	1.4	2.8	B	VG	7.8	
RIOTOUS	4Y-Y	1976 - 1979	30	18	182	1.88	S	39	10.6	3.3	nd	A	G	8.2	B
RIPPLING WATERS	5W-W	1955 - 1967	28	31	180	2.28	SB	34	9.0	2.6	2.0	A	G	8.4	
RISTIN	1Y-Y	1976 - 1982	15	-3	138	1.80	S	37	9.4	4.0	2.7	A	VG	10.2	B
RIVA	2W-W	1955 - 1963	17	20	167	1.87	S	44	10.0	4.0	4.4	A	M	nd	
ROB ROY	3W-R	1976 - 1981	-4	31	138	2.13	SB	50	10.4	1.5	2.8	A	VG	10.2	B
ROBERTA	1W	1968 - 1982	-6	18	178	1.99	S	38	11.3	4.5	4.9	A	G	8.3	S
ROBESPIERRE	2W-O	1955 - 1963	-4	20	377	2.40	SB	49	11.0	1.6	3.1	A	M	nd	
ROBEY	1Y-Y	1955 - 1981	2	10	248	2.34	S	36	11.0	5.2	5.6	B	VG	8.3	
ROCKET	2W	1968 - 1972	48	48	84	1.62	SB	38	9.8	1.0	2.3	A	G	10.6	
ROCOCO	2W	1955 - 1963	14	11	258	1.87	SB	47	12.0	2.7	6.7	A	VG	nd	
RODOMONT	2Y-O	1955 - 1981	8	24	313	2.66	SB	52	10.0	2.2	2.5	A	G	8.5	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
ROGER	6Y-O														
ROJMOND	2W-O														
ROMAINE	1W-W														
ROMAN CANDLE	2W-P	1976 - 1980	-8	22	173	2.42	SB	40	12.2	3.8	4.2	B	G	9.5	
ROMAN TILE	2W-WYP	1977 - 1980	40	22	156	1.83	SB	39	10.7	2.2	3.9	B	G	9.3	S
ROMANTICA	2O-O														
ROMANUS	4W-O	1964 - 1964	nd	nd	-28	nd	nd	nd	nd	nd	nd	nd	nd	nd	
(syn. ? DOUBLE ROMAN)															
ROMEO	8Y-O	1968 - 1987	-2	38	154	2.82	S	37	6.1	0.5	1.4	A	M	9.5	
ROSA VAN LIMA	2W-P	1964 - 1967	1	11	228	2.30	SB	34	11.5	2.9	3.4	A	G	6.8	
ROSARIO	2W-P	1964 - 1967	49	29	nd	nd	S	31	11.0	3.9	4.5	B	G	5.8	
ROSDEW	2W-P	1976 - 1977	60	27	180	2.11	S	38	8.8	2.2	2.8	A	G	10.8	
ROSE BROCADE	2W-P	1972 - 1974	82	42	106	0.65	S	24	9.0	3.2	4.2	A	G	8.4	
ROSE CAPRICE	2W-GPP	1972 - 1972	79	35	46	0.36	S	19	11.0	3.4	4.6	A	G	nd	
ROSE NOBLE	2W-P	1976 - 1980	34	36	156	2.34	S	40	11.3	2.4	3.0	B	G	8.8	B
ROSE OF CUAN	2W														
ROSE OF MAY	4W-W	1964 - 1967	51	63	124	1.50	S	42	7.0	nd	nd	A	G	8.0	
ROSE OF TRALEE	2W-P	1955 - 1967	60	48	194	1.80	SB	41	10.0	3.8	3.4	B	G	9.0	
ROSE RIBBON	2W-YYP	1972 - 1977	19	30	159	1.92	S	40	10.2	2.3	3.7	A	G	8.9	
ROSE ROYALE	2W-P	1982 - 1984	46	30	nd	2.25	S	31	10.3	3.0	3.7	B	G	nd	
ROSEANNA	2W-P	1964 - 1967	56	51	nd	nd	S	25	10.0	2.9	2.9	B	G	10.2	
ROSEATE HUES	2W-YYP	1968 - 1977	59	33	87	1.73	S	30	11.2	3.4	5.2	B	G	9.1	BS
ROSEBUD	3W-O	1955 - 1963	-14	32	303	1.87	SB	60	11.0	1.0	2.9	A	G	nd	
ROSEDEW	2W-P	1979 - 1979	80	28	225	2.00	S	29	13.0	4.5	4.3	A	G	nd	
ROSEWELL	2W-P	1964 - 1967	17	36	237	3.00	SB	29	11.0	3.4	3.6	A	G	10.6	
ROSEWORTHY	2W-P														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
ROSEYARDS	2W-P														
ROSIE MCNABB	2W-P	1976 - 1980	30	6	266	1.73	S	25	10.2	3.6	3.8	A	G	9.5	
ROSSINI	2W-O	1979 - 1985	6	35	169	1.23	S	46	12.2	1.9	3.7	A	VG	10.5	
ROSTOV	2W-W	1964 - 1968	17	33	223	1.40	SB	27	11.0	4.0	4.4	B	G	8.2	
ROSY DIAMOND	2W-WWP	1964 - 1973	30	41	172	1.82	SB	33	10.0	2.6	3.0	A	G	8.0	
ROSY PINK	2W	1976 - 1980	27	14	136	2.36	SB	30	9.8	3.3	4.1	B	G	10.3	
ROSY SUNRISE	2W-P	1955 - 1963	29	37	266	1.87	SB	46	11.0	3.8	3.8	A	G	8.0	
ROTONDE	3W-Y														VB
ROUGE	2Y-O	1955 - 1963	-20	8	527	2.40	S	36	11.0	2.0	3.4	A	G	nd	
ROVER	1Y-Y	1973 - 1975	-10	2	222	1.27	S	46	11.0	5.2	3.8	A	G	10.4	V
ROWALLANE	1Y-Y	1973 - 1973	50	28	201	1.78	S	28	10.5	4.3	4.4	A	VG	10.2	
ROWENA LEE TEAGLE	2W-ORR	1964 - 1967	-21	11	nd	nd	SB	34	11.5	2.0	4.4	A	G	7.4	
ROXANE	1W-W	1955 - 1963	14	11	97	1.33	S	30	12.0	4.9	5.4	B	G	nd	
ROYAL ARMOUR	1Y-Y	1972 - 1984	4	-2	140	1.85	S	42	10.8	4.7	4.9	A	G	8.1	
ROYAL CHARGER	2Y-O														
ROYAL CHARM	2Y-ORR	1979 - 1985	4	33	337	2.01	S	42	11.8	2.3	2.5	A	G	9.0	
ROYAL CROWN	2W-O														
ROYAL DELIGHT	1Y-Y	1976 - 1979	-7	-8	102	1.78	S	34	11.3	5.2	4.9	A	G	8.4	
ROYAL GOLD	1Y-Y	1977 - 1981	30	13	139	1.90	S	34	13.3	5.9	5.7	A	VG	12.2	S
ROYAL MAIL	2Y-O	1955 - 1981	-2	20	254	2.96	SB	50	10.0	1.8	3.8	B	G	10.2	
ROYAL OAK	1Y-Y														
ROYAL ORANGE	2W-O	1955 - 1963	14	17	401	1.33	SB	50	11.0	2.2	3.6	A	VG	9.0	
ROYAL PORCELAIN	2W-WWP	1979 - 1979	80	32	309	1.33	SB	32	10.5	3.3	4.1	B	G	nd	
ROYAL RANSOM	2Y-O	1955 - 1963	14	17	401	1.33	SB	50	11.0	2.2	3.6	A	VG	9.0	
ROYAL REGIMENT	2W-O	1979 - 1985	23	38	194	1.62	B	47	10.3	1.4	3.6	A	VG	9.2	B
ROYAL ROMANCE	2	1976 - 1980	0	12	190	2.08	SB	46	9.8	1.9	2.7	A	G	10.2	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
ROYAL SEAL	2W-R	1968 - 1972	64	52	91	1.47	SB	40	11.0	1.1	3.3	A	G	8.7	
ROYAL WINDSOR	2Y-YRR	1964 - 1972	10	30	108	2.12	SB	36	11.2	2.3	4.1	B	G	9.2	
ROYALIST	1Y-Y	1955 - 1963	63	24	146	1.07	E	38	10.0	4.2	4.5	A	G	6.0	
RUBICON	2Y-Y														
RUBRA	2W-Y	1955 - 1963	-4	26	399	2.40	S	57	10.0	1.4	3.2	B	M	nd	
RUBY KING	2	1964 - 1972	14	10	158	2.20	SB	38	12.0	2.5	4.2	A	G	8.0	
RUBY TAIL	2W-R	1972 - 1972	8	34	107	1.71	SB	35	11.5	2.1	4.8	A	G	8.4	
RUMBA	2Y-O	1968 - 1981	35	24	174	2.94	SB	38	11.8	2.6	5.2	B	G	7.4	
RUSSET	3Y-R	1968 - 1972	28	45	164	1.19	SB	44	10.5	1.3	3.3	A	VG	8.5	
RUSTOM PASHA	2Y-O	1955 - 1967	5	7	197	2.48	S	41	11.0	2.5	3.4	A	G	8.8	
SACAJAWEA	2Y-YYO	1967 - 1987	7	1	162	2.12	S	43	11.1	2.8	4.0	A	G	9.1	
SAILING LIGHT	2W-YYO	1964 - 1967	8	22	nd	nd	SB	32	10.0	1.8	4.2	A	G	8.6	
SAINTE AGNES	8W-O	1955 - 1984	-2	21	245	3.12	S	47	6.9	0.6	1.9	A	M	9.0	
SAINTE BRENDAN	2W-W	1968 - 1972	48	32	122	1.74	S	34	12.0	5.0	5.4	A	VG	9.4	BS
SAINTE BRIDE	1W-W														B
SAINTE EGWIN	2Y-Y	1955 - 1963	-14	21	143	1.87	SB	54	10.0	2.0	2.9	A	G	nd	
SAINTE ISSEY	2Y-Y	1955 - 1973	-16	7	216	1.95	E	46	10.0	3.5	4.0	A	G	9.2	
SAINTE IVES	2Y-Y	1955 - 1963	-20	7	300	1.87	E	46	10.0	2.9	3.9	A	G	nd	
SAINTE JUST	3W-YYO	1955 - 1963	-14	17	292	1.87	SB	52	10.0	1.6	3.2	A	M	nd	
SAINTE KEVERNE	2Y-Y	1955 - 1987	4	6	182	2.81	E	44	10.3	3.6	3.7	A	VG	10.0	
SAINTE KEYNE	8W-O	1955 - 1982	-36	10	183	2.75	SB	52	7.0	0.7	2.1	A	M	9.7	
SAINTE LOUIS	3Y-YOO	1955 - 1963	-4	36	185	1.87	SB	52	12.0	1.6	3.8	A	M	nd	
SAINTE MARTIN	2	1955 - 1963	-7	-7	137	1.33	E	46	10.0	1.9	3.0	A	P	nd	
SAINTE MARY	1W-W														
SAINTE MORITZ	2W-W														
SAINTEFIELD	2W-P	1973 - 1973	17	24	192	2.75	S	28	11.0	3.0	3.7	B	G	10.6	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
SALMON TROUT	2W-P	1972 - 1972	31	30	93	1.29	S	30	12.0	3.9	4.0	B	VG	10.0	
SALOME	2W-PPY	1972 - 1987	35	36	174	2.69	E	34	10.5	3.4	3.5	A	G	9.3	
SALTASH	2Y-O	1955 - 1963	-7	15	203	1.87	SB	50	12.0	2.7	3.1	A	G	nd	
SAMITE	1W-W	1955 - 1963	52	39	183	1.87	S	41	10.0	4.0	3.5	A	G	8.0	
SANTA CLAUS	4W-W	1982 - 1984	78	61	nd	2.17	S	47	9.0	nd	nd	A	G	nd	
SANTA LUCIA	2W-W														
SANTIAM	2Y-YYO	1972 - 1977	28	14	129	2.25	S	44	10.8	2.5	4.6	A	G	8.6	
SARABANDE	9W-GOR	1964 - 1967	31	33	210	2.30	SB	37	9.0	0.5	2.0	A	G	9.0	
SARAH LANCASHIRE	3W	1968 - 1972	-11	42	156	1.82	SB	45	9.2	1.2	2.5	A	G	7.2	
SARCHEDON	9W-GYR	1955 - 1982	27	41	249	2.32	SB	54	10.0	0.5	1.9	A	G	10.6	
SARGASSO	2Y-YOO	1955 - 1981	-2	28	182	2.44	SB	55	10.0	2.3	3.7	A	VG	9.7	
SASSENHEIM'S GIANT	1Y-Y	1964 - 1967	-5	17	nd	nd	S	35	13.0	4.9	5.7	B	G	nd	
SATIN QUEEN	2W-YYO														
SCANDAL	3W-O	1968 - 1980	6	39	162	2.10	SB	46	10.0	1.1	2.7	A	G	10.3	
SCAPA	1W-W	1968 - 1971	75	32	154	1.70	S	36	11.0	4.8	4.6	A	G	7.5	BS
SCARLET ELEGANCE	2Y-R	1955 - 1973	3	16	241	1.74	SB	43	10.2	1.9	3.1	A	G	9.0	
SCARLET GEM	8Y-O	1955 - 1963	-45	20	365	2.40	S	42	6.0	0.4	1.6	A	M	nd	
SCARLET LEADER	2Y-O	1955 - 1967	1	22	202	2.06	SB	42	10.0	1.6	3.2	B	G	8.0	
SCARLET PIMPERNEL	2W-O	1955 - 1963	0	23	335	1.87	SB	50	9.0	1.2	2.9	A	G	10.0	
SCARLET ROYAL	2Y-R	1972 - 1977	51	29	98	2.24	SB	39	11.3	2.3	4.0	B	G	9.5	
SCARLETT O'HARA	2Y-R	1964 - 1984	11	3	151	2.31	SB	42	11.3	2.9	3.8	A	G	8.6	
SCILLY WHITE	8W-Y	1964 - 1967	-83	4	nd	nd	nd	nd	nd	nd	nd	nd	nd	nd	
SCINTILLA	2W-R	1964 - 1967	19	21	314	2.10	SB	41	9.0	1.3	3.7	A	G	8.0	
SCORCHER	2Y-O	1977 - 1987	0	-5	163	2.33	S	36	11.5	2.8	4.7	A	M	9.9	B
SCORPION	2Y-R	1964 - 1967	10	9	276	2.90	SB	40	9.0	2.3	3.6	A	M	8.4	
SCOTCH GOLD	1Y-Y	1973 - 1973	24	21	116	1.62	S	38	10.0	4.4	4.8	A	G	8.0	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
SCRABO	2Y-O	1964 - 1981	14	22	194	2.82	SB	38	9.5	1.8	4.3	B	G	7.0	
SEA GREEN (16)															
SEA URCHIN	2W-Y	1964 - 1967	-45	16	nd	nd	S	39	12.0	3.6	4.4	A	G	7.0	
SEAFORD	2W-W	1972 - 1972	95	46	68	0.86	S	22	11.0	4.0	4.2	B	G	nd	
SEAGULL	3W-YYO	1955 - 1967	6	24	306	2.90	S	54	10.0	1.3	2.5	A	P	8.2	
SEALING WAX (15)	2Y-O	1968 - 1987	-15	13	228	2.38	B	48	11.1	2.3	2.6	A	G	11.2	
SEBASTOPOL	2W-Y	1968 - 1981	46	28	158	3.36	S	38	10.0	3.2	4.4	A	G	9.4	S
SELM LAGERLOF	2W-YOO	1955 - 1963	-4	26	319	2.93	SB	48	11.0	1.9	4.6	A	M	nd	
SELTAN															
SEMIRAMIS	2W-Y	1973 - 1973	17	20	276	1.79	S	41	11.0	2.5	5.9	A	G	9.2	S
SEMPRE AVANTI	2W-O	1955 - 1973	37	27	262	2.47	S	45	10.7	2.0	4.0	A	M	8.1	
SERAGLIO	3Y-YYR	1955 - 1967	8	28	249	2.08	SB	57	10.0	1.3	3.4	A	VG	8.7	
SHAGREEN (16)	3W-GWW	1968 - 1971	80	61	156	1.60	SB	40	8.5	1.2	2.6	A	G	8.2	
SHAH	7Y-Y	1955 - 1982	-3	8	208	2.56	E	42	10.0	3.5	3.8	A	G	8.8	
SHALDON	2W-Y	1973 - 1973	7	21	163	1.21	S	25	12.0	3.4	5.9	B	VG	8.4	
SHANACH															
SHANGHAI	2Y-Y	1968 - 1981	28	18	145	2.07	S	38	12.0	3.4	5.4	A	G	9.6	
SHANNON	2W-W	1964 - 1967	19	24	194	1.90	S	29	11.5	3.9	3.5	B	G	8.8	
SHANTALLOW	3W-GGY	1976 - 1978	34	44	108	1.92	SB	42	9.0	1.0	2.8	A	G	7.9	
SHEEROE															
SHEIK	2W-YOY	1982 - 1984	53	44	nd	2.88	B	40	10.8	1.5	3.4	A	G	nd	
SHEPHARD	3W-OR	1973 - 1973	59	36	150	2.33	S	37	9.5	1.0	2.3	B	G	8.6	
SHEPPY															
SHINING WATERS	2W-W	1964 - 1967	19	9	nd	nd	S	28	11.0	3.2	4.6	A	G	8.2	
SHIRLEY WYNESS	2W-YYP	1964 - 1967	40	24	233	2.20	S	32	9.5	2.9	4.1	B	G	8.8	
SHOT SILK	5W-W	1955 - 1967	10	26	219	2.04	E	38	9.0	2.4	2.3	A	M	9.2	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
SHOWMAN	2W-Y	1964 - 1967	51	32	nd	nd	SB	26	10.0	2.6	5.4	B	G	7.8	
SIAM	2W-P	1964 - 1967	51	32	181	1.80	S	28	10.0	3.4	5.0	B	G	7.0	
SIBERIA	2W-W														
SIDELIGHT	9W-GOR	1964 - 1967	55	62	nd	nd	S	36	7.0	0.5	1.9	A	G	8.6	
SIDNEY TORCH	2Y-YOO	1964 - 1982	20	16	198	3.02	SB	42	10.5	1.3	4.3	A	G	8.0	
SIGNAL LIGHT	2W-O	1968 - 1973	32	30	87	1.61	SB	41	11.2	1.9	4.6	A	G	9.2	BS
SIGRID UNSET	3W-YOO	1955 - 1963	-4	20	289	1.87	SB	50	10.0	1.6	3.5	B	G	nd	
SILENT BEAUTY	3W-YJR	1979 - 1981	39	38	301	2.18	SB	36	10.0	1.2	3.0	B	VG	9.0	
SILENT GLOW	3W-R	1979 - 1981	44	42	251	1.37	SB	40	8.2	1.0	2.6	B	G	11.0	
SILENT MORN	3W-YJO	1979 - 1981	22	34	300	1.75	SB	38	10.0	1.2	2.7	B	G	8.3	
SILVER BUGLE	2W-W	1955 - 1963	-7	4	417	1.87	S	34	11.0	3.0	4.5	B	G	8.4	
SILVER CHIMES	8W-W	1955 - 1967	-46	30	146	1.00	SB	28	6.0	1.2	1.7	A	G	7.2	
SILVER LINING	2W-W	1955 - 1963	52	10	184	1.87	S	41	11.0	3.0	4.5	B	VG	10.0	
SILVER MOON (17)	2W-O	1971 - 1987	13	37	158	2.32	S	39	11.3	2.4	5.1	A	G	9.1	S
SILVER PENNY	3W-O	1955 - 1963	49	37	187	1.87	S	52	8.0	0.8	2.6	A	P	nd	
SILVER PRINCESS	3W-W	1964 - 1971	29	59	173	1.20	S	41	8.7	0.6	1.9	A	G	10.6	
SILVER SALVER	3W-GWW	1955 - 1967	28	47	259	1.56	SB	42	8.0	1.0	2.5	A	M	6.0	
SILVER SAND (NL)		1967 - 1987	39	38	167	2.11	S	43	9.9	1.6	3.4	A	G	10.6	BS
SILVER STANDARD	2W-W	1970 - 1987	17	3	194	2.40	S	42	11.8	2.7	5.1	A	G	9.5	
SILVER WEDDING	1W-W														
SILVERTON	1W-W	1977 - 1977	43	15	75	0.33	S	23	9.5	4.2	3.9	A	M	nd	V
SILVESTER	11W-Y	1968 - 1968	84	41	272	1.20	S	26	9.0	1.4	3.5	B	G	8.0	
SILVRETTA	1Y-Y														
SINCERITY	1W-Y	1964 - 1967	34	37	179	2.30	S	30	9.5	4.7	4.3	B	G	8.8	
SIR PERCY	2Y-O	1973 - 1973	9	24	175	2.33	S	33	9.5	1.9	4.2	B	G	10.0	
SIR SAMUEL	2W-P	1976 - 1987	19	14	165	1.95	E	32	10.3	3.9	2.8	A	M	9.3	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
SIR WATKIN	2Y-Y	1955 - 1967	16	13	215	2.43	S	42	10.0	2.8	3.1	A	M	9.0	
SIR WINSTON CHURCHILL	4W-O	1982 - 1982	53	34	220	1.79	S	26	6.0	0.8	nd	nd	G	9.2	
SISTER REINALDE	2W-YO	1955 - 1963	39	22	507	1.87	S	46	11.0	2.8	4.5	A	G	nd	
SLEMISH	2W-W														B
SLEVEEN	2W-W														
SLIEVEBOY	1Y-Y	1973 - 1973	57	32	144	2.04	S	32	11.0	5.1	5.0	A	G	10.8	VS
SLIGO	2Y-Y	1964 - 1967	13	31	133	1.70	S	31	10.0	4.1	4.7	B	G	7.0	
SMARAGD	2W	1964 - 1967	11	26	nd	nd	SB	36	11.5	2.3	5.0	B	G	7.0	
SMILING MAESTRO	2Y-R	1976 - 1980	10	18	171	2.78	SB	46	11.5	2.1	4.6	B	G	9.5	
SMILING MARY	3W	1972 - 1973	33	42	142	1.62	SB	34	10.2	1.0	3.0	A	G	10.9	
SMILING QUEEN	2W-YO	1955 - 1963	14	20	347	2.93	SB	46	10.0	2.0	4.0	B	M	nd	
SMYRNA (16)	9W-GOO	1955 - 1963	21	56	213	1.87	SB	42	7.0	0.5	1.9	A	G	nd	
SNEEUWPRINSES	3W-YO	1964 - 1973	13	29	288	3.24	SB	32	9.3	1.2	3.0	A	M	9.7	
SNOW BUNTING	7W-Y														B
SNOW DREAM	2W-W														B
SNOW GEM	3W-O	1970 - 1987	-2	24	187	1.94	B	45	10.1	1.1	3.0	A	M	8.4	
SNOW KING	9W-YR	1964 - 1967	20	46	nd	nd	SB	39	9.0	0.5	1.8	A	G	11.8	
SNOWBALL	4W-W	1972 - 1972	45	34	5	0.75	S	34	10.0	2.0	nd	A	G	8.0	
SNOWBIRD	5W-W	1955 - 1967	1	28	201	1.82	SB	34	9.0	2.5	2.0	A	M	8.3	
SNOWLINE	2W-W														
SNOWSHILL	2W-W	1977 - 1980	41	34	184	2.24	SB	46	9.5	1.9	3.4	A	G	9.3	B
SOIREE	2W-Y														
SOLARIO	2W-Y	1955 - 1963	14	15	261	2.40	S	46	11.0	3.8	5.3	A	M	nd	
SOLE MIO	1Y-Y	1955 - 1967	-8	8	198	1.94	E	38	11.0	4.9	5.5	A	G	8.5	
SOLE PONENTE	1Y-Y	1973 - 1979	22	12	130	2.13	S	40	12.1	5.4	5.5	A	G	9.8	
SOLINUS	2W-P	1976 - 1980	25	13	92	1.88	S	32	9.8	3.5	3.0	B	G	8.0	B

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
SOLOGNE	2Y-O														
SOLVEIG	1W-Y	1955 - 1963	-4	21	331	2.40	E	38	12.0	4.7	4.9	B	G	nd	
SOMALI	2Y-YOO	1955 - 1967	32	53	223	2.40	SB	44	8.0	0.5	2.0	A	G	9.6	B
SONATA	9W-GYR	1973 - 1982	30	48	166	2.31	nd	nd	nd	nd	nd	nd	nd	nd	
SONIA SLOAN	2W-OOY	1977 - 1977	57	14	377	1.00	S	37	8.5	3.1	2.1	A	G	9.0	
SORBET	11W-YYO	1979 - 1984	18	32	217	2.08	SB	37	10.0	3.0	5.3	B	M	8.0	
SOUTH POLE	1W-W	1973 - 1973	50	28	177	2.21	S	34	11.5	5.0	5.8	A	VG	8.0	
SPACE AGE	2Y-Y	1976 - 1980	16	14	147	2.44	SB	43	10.7	3.3	3.7	A	VG	7.6	
SPANISH GOLD	1Y-Y														
SPARKLING EYE	8W-GOO	1964 - 1967	34	40	192	2.10	SB	34	6.5	0.7	1.7	A	G	8.1	
SPECTATOR	2W-YYO														B
SPELLBINDER (15)	1Y-WWG	1964 - 1982	44	16	182	2.13	s	39	11.4	5.1	5.2	A	g	7.7	S
SPELTER	2Y-O	1964 - 1967	-36	24	nd	nd	S	31	10.5	2.7	4.0	B	VG	11.2	
SPENCER TRACY	2W-O	1955 - 1973	16	26	231	2.30	SB	47	9.7	1.6	3.8	A	VG	9.3	
SPICK AND SPAN	2Y-YYO	1964 - 1973	1	28	188	2.18	SB	39	9.5	1.5	2.7	A	VG	10.0	
SPITZBERGEN	1W-Y	1955 - 1982	21	29	188	2.28	E	43	10.0	4.4	4.4	A	G	7.5	B
SPLIT	11W-W	1964 - 1967	23	26	nd	nd	S	33	10.0	2.6	6.2	B	M	6.8	
SPRING BEAUTY	3Y-O	1955 - 1949	-14	15	260	1.87	S	50	10.0	1.6	3.6	B	M	nd	
SPRING FEVER	2Y-O														
SPRING GLORY	1W-Y	1955 - 1967	4	12	197	1.42	S	32	11.0	5.0	4.9	B	P	7.6	
SPRING MORNING	1Y-Y	1968 - 1982	34	-4	224	2.17	S	26	11.7	4.6	5.3	A	G	8.4	
SPRY	2Y-R	1968 - 1973	23	38	156	1.38	SB	40	10.0	1.6	2.2	A	VG	1.2	
SQUIRE	1Y-Y	1979 - 1984	18	17	152	2.39	S	38	12.9	5.4	4.9	A	VG	9.5	B
STADIUM	2W-Y	1964 - 1967	47	9	158	1.90	S	30	11.5	4.3	6.6	B	G	6.0	
STAINLESS	2W-W	1982 - 1985	40	40	nd	1.57	B	34	9.3	1.9	3.6	A	G	10.1	
STANDARD BEARER	1Y-Y														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
STANDARD VALUE	1Y-Y	1964 - 1987	23	22	163	2.39	SB	42	12.3	4.7	5.2	A	G	7.4	
STATUE	2W-Y	1972 - 1984	25	16	134	2.01	S	51	14.0	3.3	4.0	nd	VG	7.0	
STELLA	2W-Y	1964 - 1967	-15	21	nd	nd	S	35	9.0	1.4	2.0	B	P	nd	
STERLING	1Y-Y	1968 - 1982	29	17	157	2.17	S	37	11.7	5.4	4.6	A	VG	7.8	
STILTON	9W-YR	1973 - 1982	13	46	130	1.72	SB	49	8.4	0.7	2.2	A	G	9.1	
STOREEN	9W-YO	1955 - 1963	27	34	287	2.40	S	50	8.0	0.5	2.0	A	G	nd	
STORMONT	1W-W														
STOUT LAD	2Y-Y														
STRAIGHT	1W-Y	1968 - 1982	26	29	160	2.26	S	35	11.4	4.7	5.2	A	G	8.6	S
STRAIGHT FLUSH	2Y-O	1972 - 1973	-16	13	156	2.21	S	38	8.8	2.0	3.1	A	VG	11.2	S
STRAY PINK	2W-P	1964 - 1967	49	32	215	1.40	S	33	9.0	2.6	3.2	A	G	9.0	
STRESA	1W-W	1955 - 1963	-4	9	245	1.87	S	37	12.0	5.3	5.0	A	G	nd	
STROMBOLI	2W-O	1970 - 1987	8	35	182	2.25	B	44	10.2	1.8	4.2	A	G	9.3	B
STRONGHOLD	1Y-Y	1964 - 1967	26	22	152	1.90	S	31	11.5	5.4	5.4	B	G	7.2	
SUCCESSOR	1Y-Y	1964 - 1967	20	30	115	0.40	S	30	10.5	4.6	4.4	A	G	6.4	
SUDA	2W-O	1955 - 1963	0	20	201	2.40	SB	48	10.0	3.7	3.4	A	M	nd	
SUDAN	2Y-R	1955 - 1981	-18	23	228	2.27	SB	49	10.0	1.8	1.9	A	G	9.2	
SUILVEN	3W-W	1973 - 1973	-17	32	326	1.62	S	40	10.5	1.5	3.3	B	VG	11.0	
SULPHUR	1Y-Y	1955 - 1967	10	3	289	2.82	S	36	11.0	4.5	4.4	A	G	8.4	
SULPHUR PRINCE	1Y-Y	1955 - 1963	-11	31	182	1.87	S	42	10.0	4.7	4.0	B	G	nd	
SUMATRA	2Y-O	1955 - 1963	-11	5	184	1.87	SB	34	11.0	2.8	4.9	B	VG	nd	
SUN CHARIOT	2Y-O	1968 - 1982	1	24	170	2.84	S	43	12.5	2.2	3.8	A	G	11.4	
SUN DANCE	1Y-Y	1955 - 1981	21	7	153	1.80	S	42	10.0	4.0	5.0	A	VG	7.3	
SUN FIRE	3Y-R	1982 - 1984	4	48	206	2.31	S	50	8.6	1.1	2.1	A	G	8.8	
SUN SALUTE	2Y-O														
SUNBURST	4Y-Y	1977 - 1978	43	18	513	1.41	S	39	12.8	4.2	3.0	A	G	8.4	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
SUNCREST	2														
SUNLIT HOURS	2Y-Y	1973 - 1975	23	20	274	2.25	S	40	12.8	4.6	4.5	A	G	9.1	V
SUNNY BROOK	2W-O	1955 - 1963	29	37	209	1.87	SB	48	9.0	1.7	3.7	A	M	nd	
SUNRIDGE	2Y-OOY	1976 - 1977	18	9	147	3.00	SB	46	9.0	2.2	3.2	A	G	8.6	
SUNRISE	3W-YYO	1964 - 1967	5	11	265	3.60	SB	37	8.0	1.0	2.3	A	P	9.2	
SUNSILK	2Y-Y	1973 - 1973	9	14	222	2.54	E	39	10.0	3.3	4.0	A	G	7.2	S
SUNSTAR	3W-R	1955 - 1967	61	52	224	1.12	SB	48	9.0	0.9	2.4	A	VG	8.2	
SVENGALI	1W-Y	1955 - 1963	-20	0	248	1.87	S	38	10.0	4.6	5.4	B	P	nd	
SWANSDOWN	4W-W	1955 - 1963	14	46	177	1.87	SB	40	8.0	1.8	nd	A	G	nd	
SWEET ADARE	1W-Y	1955 - 1963	14	10	234	2.40	S	31	10.0	3.7	4.9	B	M	nd	
SWEET CHARITY	2W-YYO	1981 - 1985	19	29	243	1.63	S	34	11.3	2.8	4.7	A	VG	11.8	
SWEET MEMORY	2W-P	1972 - 1984	18	21	147	2.01	S	39	10.4	2.3	5.1	A	G	9.3	S
SWEETHEART	2W-YYO	1955 - 1963	-7	25	359	2.40	SB	50	10.0	1.8	4.1	A	VG	nd	
SWEETNESS	7Y-Y	1955 - 1967	-34	12	486	4.25	E	48	8.0	2.2	2.1	A	G	9.4	
SWISS MISS	2W-P														
SYLVESTER	1W-W	1967 - 1968	82	34	196	1.00	S	24	11.8	5.1	6.0	A	G	6.6	
SYLVIA O'NEILL	3W-WWY	1972 - 1972	53	39	84	0.96	SB	33	10.0	1.5	3.2	A	G	7.2	
SYRACUSE	3W-WWY	1977 - 1981	-2	37	195	2.12	SB	41	10.9	1.5	3.3	B	VG	10.1	
TABASCO	2														V
TAIN	1W-W														
TAKORADI	4W-W	1977 - 1984	-3	35	116	1.60	S	43	10.3	2.9	nd	B	G	8.5	B
TALLAND	2Y	1968 - 1987	-1	16	209	3.02	S	46	10.6	2.8	4.0	A	G	7.7	
TAMAR FIRE	4Y-R	1979 - 1985	-8	14	149	1.89	S	31	8.1	1.7	nd	A	G	8.6	S
TAMARA	2Y-Y	1984 - 1984	-8	-27	350	1.46	S	34	11.0	4.5	4.5	nd	M	5.8	
TAMINO	2Y-O	1955 - 1963	0	23	302	1.33	SB	42	10.0	1.5	3.5	A	G	9.2	
TAMSYN	1Y-Y	1984 - 1984	-22	-9	nd	0.83	E	38	10.0	4.0	4.5	nd	G	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
TANNAGHMORE	1W-Y	1982 - 1984	77	37	nd	0.58	B	43	nd	nd	nd	A	nd	nd	
TANNHAUSER	2Y-YYO	1968 - 1982	13	23	176	3.18	SB	40	9.3	2.1	4.7	A	G	9.0	
TARZAN	2W-O	1964 - 1967	-2	40	nd	nd	SB	32	14.0	3.0	5.2	B	G	7.6	
TAWNY GOLD	1Y-Y	1972 - 1975	32	7	161	2.89	S	32	12.7	5.4	4.6	A	VG	8.8	
TAWNY LAD	2Y-O	1984 - 1984	20	38	nd	2.00	S	36	10.0	1.9	2.8	A	G	nd	
TEAROSE PHOENIX	4Y-Y	1955 - 1963	37	26	173	1.87	SB	42	9.0	1.4	nd	A	P	nd	
TEBESSA	3W-R	1955 - 1963	14	37	241	2.40	SB	52	9.0	1.1	3.5	A	G	nd	
TEBOURBA	3W-R	1955 - 1963	0	37	198	1.87	SB	51	10.0	1.0	2.8	A	G	nd	
TECOMA	2Y-O	1968 - 1987	0	-6	150	2.16	S	44	11.7	3.5	4.6	A	M	9.0	B
TEHERAN	2Y-O	1964 - 1967	19	38	209	3.10	SB	45	9.0	1.5	3.4	A	VG	11.0	
TELAMONIUS PLENUS	4Y-Y	1955 - 1967	27	1	175	1.94	S	30	11.0	nd	nd	A	P	7.0	
TELSTAR	1W-W	1977 - 1978	25	14	265	2.56	S	30	12.0	3.4	6.1	B	G	9.0	
TEMPLEMORE	2W-W	1968 - 1973	78	44	142	1.80	SB	34	11.2	3.2	4.2	A	G	10.8	
TEMPLEPATRICK	2W-W														
TEMPURA	2Y-Y	1975 - 1975	21	29	nd	nd	SB	30	10.5	3.5	4.0	A	G	nd	V
TENEDOS	2W-Y	1955 - 1967	-8	18	162	2.08	S	47	12.0	4.1	4.1	A	G	8.8	
TEXAS	4Y-O	1955 - 1963	-4	27	525	1.87	SB	46	10.0	2.0	nd	A	G	nd	
THALIA	5W-W	1955 - 1967	2	24	213	2.08	S	33	9.0	2.1	2.6	A	P	8.2	
THE FIRST	1Y-Y	1955 - 1973	-3	-2	240	2.25	S	42	10.3	4.7	4.7	A	M	6.7	
THE GIFT	1Y-Y	1977 - 1978	32	12	262	2.23	S	40	9.5	4.6	4.0	A	G	8.3	
THE KNAVE	6W-Y	1970 - 1970	21	9	206	1.43	S	22	nd	nd	nd	B	nd	10.4	
THE SUN	3W-O	1955 - 1963	-20	51	305	2.40	SB	48	8.0	0.5	1.7	B	M	nd	
THERAPIA	3W-YYO	1955 - 1963	14	31	267	1.33	SB	47	11.0	1.9	4.4	A	VG	nd	
THERM	3Y-O	1964 - 1967	5	18	nd	nd	SB	43	9.5	1.3	2.8	A	G	8.8	
THETIS	9W-R														
THOMAS HARDY	9W-R														

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
THOMAS MARK	2Y-YOO	1955 - 1963	-20	10	417	1.87	S	34	10.0	2.6	5.4	B	M	nd	
THRILLER	2Y-R	1972 - 1977	27	16	143	2.10	SB	42	9.8	2.2	4.3	A	G	8.7	S
THURSO	1W-W														B
TIBET	2W-W	1964 - 1987	8	16	165	2.54	S	41	1.2	3.9	5.9	A	G	10.7	
TILLA DURIEUX	3W-GYO	1955 - 1963	21	46	287	1.33	SB	38	9.0	0.9	2.5	A	VG	nd	
TINKER	2Y-O	1955 - 1987	-4	20	244	2.55	S	57	10.1	2.4	2.8	B	G	8.2	
TINO ROSSI	2Y-O	1955 - 1963	-20	15	367	1.87	SB	47	10.0	2.1	4.3	A	G	nd	
TINSEL	3W-WWY	1964 - 1967	26	34	nd	nd	SB	34	9.0	1.4	2.8	B	G	7.4	
TINTORETTO	1Y-Y	1955 - 1963	10	12	184	1.33	S	41	10.0	4.1	4.2	A	G	nd	
TITANIA (16)	6W-W														
TITCH	1Y-Y	1972 - 1973	-9	-24	115	1.29	S	32	11.2	4.8	5.2	A	G	13.0	B
TITOKI	1	1976 - 1978	46	21	225	1.37	S	40	10.3	4.5	4.7	A	M	8.7	
TITTLE-TAITLE (16)	7Y-GYY														
TIVOLI	2Y-R	1968 - 1972	12	38	136	1.96	SB	38	10.8	2.0	3.8	B	G	9.6	S
TOLEDO	2														
TOORAK GOLD	2Y-Y	1968 - 1987	-12	-16	191	2.01	S	44	10.7	3.0	4.3	A	M	9.3	B
TOPAZ	7W-Y														
TOPIC	2W-P	1955 - 1963	60	28	87	1.87	SB	40	12.0	4.0	4.7	A	M	9.0	
TOP-NOTCHER	1Y-Y	1976 - 1981	8	7	101	1.44	S	44	10.9	5.1	4.9	A	G	11.1	
TOPSCORE	2	1968 - 1981	8	14	187	2.38	S	36	10.0	2.2	3.6	A	G	7.6	V
TORCHDANCE	2Y-Y	1968 - 1981	8	32	177	2.59	S	42	10.7	3.2	4.2	B	G	10.6	B
TOREADOR	3W-R	1970 - 1987	-4	31	179	2.20	B	44	10.2	1.4	3.2	A	G	8.7	B
TORERO	2Y-O														V
TORNAMONA	2W-W	1968 - 1982	69	60	132	2.06	SB	39	11.0	3.6	4.0	A	G	9.7	BS
TORQUAY	2Y-O	1976 - 1977	26	29	206	2.15	SB	46	8.8	1.2	3.1	A	G	8.2	
TOSCANINI	2W-WWP	1955 - 1963	29	17	179	1.87	SB	40	10.0	3.1	3.8	B	G	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
TOUJOURS	1Y-Y	1983 - 1985	72	46	64	0.84	S	32	11.8	4.7	0.3	B	G	nd	
TRAMORE	2W-Y	1968 - 1972	51	15	187	1.76	E	37	11.8	4.6	4.8	A	G	8.2	B
TREDORE	3Y-O	1955 - 1963	16	29	340	2.40	SB	50	11.0	1.4	3.0	A	M	nd	
TREEN	2Y-OOR	1972 - 1972	53	23	98	2.04	SB	39	11.0	2.8	4.4	A	G	8.8	
TREFFRY	3W-O	1955 - 1963	17	20	293	1.87	S	42	9.0	1.1	2.9	A	M	nd	
TRENANCE	2Y-Y	1955 - 1973	-14	4	179	1.80	S	42	10.0	3.5	4.3	A	G	7.5	
TRENTHON	1Y-Y	1976 - 1981	24	21	127	2.15	S	42	11.0	4.3	3.6	A	G	9.3	
TRENOON	2Y-Y														
TRESAMBLE	5W-W	1955 - 1967	-18	34	244	2.35	SB	42	10.0	2.2	2.8	B	G	8.6	
TRESITHICK (18)	2Y-Y														
TRESKERBY	3Y-R	1955 - 1963	17	23	225	2.40	SB	44	11.0	1.3	2.4	A	M	nd	
TRESKEWES	5Y-Y	1964 - 1967	-2	23	nd	nd	S	29	10.0	4.6	4.3	A	M	6.4	
TRESSERVE	1Y-Y	1955 - 1967	-14	15	157	1.62	S	40	12.0	5.5	5.6	A	G	7.0	
TREVISKY	2Y-O	1955 - 1963	-20	17	351	2.40	S	42	10.0	1.7	2.8	A	G	nd	
TREVITHIAN	7Y-Y	1955 - 1967	-36	14	279	2.34	S	51	7.0	1.4	2.2	B	M	8.0	
TREWINCE	2Y-Y	1955 - 1963	-7	10	261	2.40	SB	45	11.0	3.4	4.3	A	G	nd	
TREWIRGIE	6Y-Y	1955 - 1963	21	20	193	1.33	S	34	9.0	4.1	3.4	A	G	nd	
TREWITHEN	1Y-Y	1977 - 1987	10	10	162	2.21	E	38	11.2	4.5	4.3	A	G	7.6	
TRIANON	3W-Y	1973 - 1973	62	28	245	2.79	nd	nd	nd	nd	nd	nd	nd	nd	S
TRIFINE	2Y-O	1974 - 1987	7	16	208	2.27	S	41	12.3	3.0	3.6	A	G	9.4	
TROCADERO	1W-Y														
TROSTAN	1W-Y	1968 - 1981	57	28	128	2.08	S	38	10.8	5.0	5.0	A	G	8.5	S
TROUBADOUR	2W-P	1976 - 1980	40	15	167	2.02	S	37	11.9	3.1	3.2	B	VG	9.2	B
TROUSSEAU	1W-Y	1955 - 1984	14	7	175	2.26	S	38	10.2	4.2	4.2	A	G	8.1	
TRUELOVE (18)	3W-R	1964 - 1974	14	36	204	1.08	S	46	8.6	1.1	2.8	A	G	9.5	
TRUMPETER	1Y-Y	1970 - 1987	4	13	153	2.28	S	44	11.3	4.9	4.4	A	G	8.8	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase life (days)	Disease notes
TRUTH	2W-W	1955 - 1963	-4	8	107	1.33	S	38	11.0	3.9	4.2	A	VG	nd	
TRYST	2W-W	1964 - 1967	51	42	163	1.60	SB	42	9.0	1.8	2.6	A	VG	7.8	
TUDOR KING	1W-Y	1972 - 1972	45	30	63	0.71	S	28	11.5	5.2	5.2	B	G	8.0	
TUDOR MINSTREL	2W-Y	1968 - 1972	43	31	197	2.08	SB	39	12.8	2.7	4.6	B	VG	9.5	
TULLYGARLEY	2Y-R	1964 - 1967	5	32	nd	nd	SB	42	10.0	1.7	3.4	A	G	8.2	
TULLYGLASS	2W-WWY	1972 - 1973	37	25	122	1.75	S	40	10.8	2.6	3.8	A	VG	9.6	S
TULLYROE	2W-R	1970 - 1987	-10	38	160	2.05	B	49	10.4	1.5	2.8	A	G	10.7	
TULLYROSE	2W-P	1976 - 1977	72	40	139	1.70	S	28	10.2	3.2	3.8	A	G	8.5	
TULLYROYAL	2W-P	1982 - 1984	54	48	nd	3.50	S	39	11.5	3.3	3.5	A	G	nd	
TULYAR	3W-R	1972 - 1975	32	42	142	0.86	S	38	9.2	0.8	2.5	A	VG	10.1	
TUNIS	2W-WWY	1955 - 1963	-11	15	206	1.87	E	46	12.0	3.7	4.2	B	G	nd	
TURIN	3W-GOO	1955 - 1963	14	39	277	1.87	SB	62	10.0	0.8	2.7	A	M	nd	
TUSKAR LIGHT	2W-YYR														
TUTANKHAMUN	2W-GWW	1981 - 1983	18	23	242	1.78	S	35	12.4	4.4	4.6	A	VG	10.0	
TWINK	4W-O	1955 - 1963	-7	12	340	1.87	S	44	10.0	1.5	nd	A	P	nd	
ULSTER BEAUTY	2W-WWY	1968 - 1982	40	18	170	2.15	S	40	10.2	3.3	3.8	A	G	9.5	
ULSTER MAIDEN	2W-R	1977 - 1980	19	44	188	1.70	SB	41	10.8	1.3	3.2	B	VG	9.2	
ULSTER PRINCE	1Y-Y	1968 - 1982	18	16	164	2.08	S	42	11.2	5.0	4.4	A	G	8.8	S
ULSTER QUEEN	1W-W	1982 - 1984	58	32	nd	2.00	S	30	12.5	4.7	4.8	A	VG	nd	
ULTIMUS	2Y-O	1967 - 1987	17	55	157	1.91	B	53	12.2	2.2	4.1	B	G	9.2	
UNIQUE	4W-Y	1983 - 1985	-12	40	nd	0.84	S	36	11.8	2.9	nd	A	G	nd	B
UNSURPASSABLE	1Y-Y	1955 - 1984	3	8	177	1.97	S	38	12.1	5.6	5.6	A	G	7.7	
URANIA	3	1973 - 1973	17	18	196	1.58	S	36	10.5	1.9	4.0	A	VG	9.6	
URSA	2W-WPP	1972 - 1975	24	20	355	1.38	S	39	11.5	4.2	3.9	A	G	9.9	
VAHSEL BAY	2W-W	1968 - 1973	66	30	172	1.66	S	39	12.8	4.3	6.0	A	G	7.6	
VALENCIA	4Y-O	1955 - 1963	49	38	124	1.33	SB	42	12.0	2.4	nd	B	G	nd	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Peduncle diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
VALLANT SPARK	2Y-O	1964 - 1983	-13	5	212	2.24	SB	45	10.4	3.0	3.5	A	G	9.5	
VAN WAVEREN'S GIANT	1Y-Y	1955 - 1967	10	16	166	2.14	S	38	11.0	5.1	4.5	B	P	8.2	
VAN WERELD'S FAVOURITE	1W-Y	1955 - 1963	52	16	209	1.33	S	34	11.0	4.9	5.1	A	VG	nd	
VARIANT	2Y-R	1972 - 1977	-6	23	154	3.31	S	38	11.0	2.4	4.8	B	G	12.0	S
VARNA	3Y-YOO	1955 - 1980	3	24	172	2.36	SB	50	10.0	1.5	3.0	A	G	10.0	B
VELVETHEEN	2Y-Y	1964 - 1981	7	17	158	2.92	S	44	11.0	2.7	3.3	A	VG	8.2	
VENUS	2W-YYO														
VERDI	1W	1964 - 1967	19	54	170	1.70	S	34	11.2	5.2	5.2	A	G	8.8	
VERGER	3W-R	1955 - 1984	8	30	218	2.38	S	52	9.8	0.9	2.8	A	M	9.6	
VERONA	3W-W	1977 - 1981	44	39	212	2.18	S	44	9.9	1.4	4.0	B	G	9.4	
VICLAND	2W	1968 - 1981	14	28	181	2.43	SB	41	11.2	2.3	4.2	A	G	9.4	S
VICTOR BORGE	1W-Y														
VICTORIA	1W-Y	1964 - 1967	49	17	185	2.20	S	32	9.5	4.4	4.2	A	M	7.0	
VICTORIOUS	2W-Y	1955 - 1984	10	3	212	2.48	S	45	10.1	3.5	3.6	A	G	7.4	
VIGIL	1W-W	1972 - 1973	44	28	146	1.60	S	28	12.0	5.1	5.0	A	VG	10.2	S
VIKING	1Y-Y	1968 - 1987	14	22	153	2.37	S	39	11.6	4.7	3.9	A	G	9.6	
VILLAGE BEAUTY	3W-YY	1964 - 1967	36	30	nd	nd	S	31	8.5	1.2	2.6	B	M	9.0	
VILNA	2W-W	1972 - 1972	-11	19	310	2.12	S	38	12.0	4.0	4.3	B	G	8.8	
VIRTUE	2Y-Y	1955 - 1963	7	30	161	1.33	E	34	10.0	3.8	3.9	B	VG	10.2	
VIVEX	2Y-YYO	1972 - 1972	60	29	23	0.67	S	24	9.5	1.6	3.9	B	G	8.0	S
VIXI	2Y-Y	1974 - 1974	53	38	239	0.90	S	35	9.0	3.2	3.7	A	G	11.0	V
VOLGA	2W-W	1964 - 1967	34	27	nd	nd	SB	33	10.0	3.2	4.0	B	G	nd	
VOLTERRA	2Y-O	1972 - 1972	-11	1	169	0.36	S	32	10.5	2.2	3.3	B	M	nd	
VULCAN	2Y-O	1971 - 1987	4	16	146	2.15	S	44	10.2	2.6	2.9	A	G	11.1	
W. P. MILNER	1W-W	1955 - 1963	36	10	130	1.33	E	22	7.0	3.1	2.5	A	P	nd	
WAHKEENA	2W-Y	1976 - 1980	22	23	176	2.30	SB	40	11.5	4.1	3.8	B	G	9.1	

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
WALT DISNEY	2Y-O	1955 - 1981	1	12	171	2.54	SB	46	12.0	2.5	3.1	A	VG	9.8	B
WANDIN GLORY	1Y-Y	1972 - 1973	20	12	112	1.88	S	40	10.2	4.1	5.2	A	G	9.3	
WARIALDA	1Y-Y	1976 - 1980	16	3	149	1.00	S	39	12.5	5.0	5.8	A	G	nd	
WARLOCK	2W-YYO	1955 - 1963	-14	20	184	2.13	SB	54	11.0	1.9	3.5	A	G	nd	
WARM SPRINGS	2Y-YYO	1955 - 1963	-20	37	255	1.87	SB	50	11.0	2.0	4.7	B	M	nd	
WARRIOR	2Y-O	1964 - 1967	19	24	nd	nd	S	39	9.5	1.9	3.0	A	G	9.8	
WARWICK	1Y-Y	1972 - 1973	20	23	242	1.62	S	39	10.2	4.8	4.4	A	G	8.0	
WATERPERRY (19)	7W-YPP	1964 - 1967	-12	15	nd	nd	S	29	8.0	2.7	2.7	B	M	7.9	
WATERVILLE	2W-W	1964 - 1967	40	34	222	1.10	SB	32	9.5	1.9	3.9	A	VG	9.0	
WEARDALE PERFECTION	1W-Y	1955 - 1963	52	18	183	2.93	E	24	8.0	3.6	3.2	B	P	9.4	
WEDDING BELL	2W-W	1968 - 1982	52	23	175	2.18	S	38	11.4	3.7	4.0	A	VG	8.5	BS
WEINER BLUT	2W-O	1975 - 1975	8	18	nd	nd	SB	41	10.5	2.2	4.0	A	G	9.2	V
WEISSHORN	1W-W	1972 - 1972	60	34	183	1.62	S	33	11.0	4.9	5.4	B	G	9.0	
WELCOME (NL)		1976 - 1987	9	-4	167	2.07	S	38	12.8	5.3	6.1	A	G	8.3	V
WELL-BORN	2W-W	1964 - 1980	23	28	110	1.82	SB	33	10.0	4.2	4.5	B	G	6.6	B
WEXFORD	2Y-Y	1964 - 1967	-9	17	nd	nd	S	34	10.5	4.4	5.1	B	G	6.0	
WHANG-HI (15)	6Y-O														
WHITE CHIEF	1W-W	1973 - 1987	15	30	143	2.14	S	41	12.9	4.8	5.1	A	G	10.1	V
WHITE CITY	3W-W	1955 - 1963	27	29	256	2.40	SB	52	10.0	1.0	2.5	A	M	nd	
WHITE EMPEROR	1W-W	1955 - 1967	1	20	164	2.43	S	33	10.0	3.9	3.3	B	P	8.0	
WHITE GOLD	2W-Y	1980 - 1985	12	5	149	2.21	S	38	12.0	4.1	5.9	A	VG	8.1	
WHITE LADY	3W-Y	1955 - 1967	1	29	324	3.13	SB	56	10.0	1.2	2.5	A	P	8.8	
WHITE LION	4W-WYY	1955 - 1984	-2	28	177	1.82	SB	47	10.5	2.3	nd	A	G	8.7	
WHITE MAGIC	2W-GWW	1972 - 1977	54	28	118	2.04	S	34	10.5	4.2	3.8	B	G	9.8	
WHITE MAIDEN	2W-W														
WHITE MARVEL	4W-W	1977 - 1980	0	34	168	2.46	SB	35	9.1	2.1	3.6	A	M	8.3	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower subs-tance	Vase-life (days)	Disease notes
WHITE NILE	2W-W	1955 - 1967	1	8	218	2.43	SB	38	10.0	3.5	3.6	A	M	6.5	
WHITE PLUME	2W-W	1968 - 1973	46	48	159	1.50	SB	41	10.9	1.7	3.1	A	G	8.6	S
WHITE PRINCE	1W-W	1973 - 1977	47	26	87	1.86	SB	34	12.2	4.8	4.5	B	VG	9.8	S
WHITE PROSPECT	1W-W	1973 - 1973	43	25	123	1.17	S	30	11.5	5.5	5.4	B	VG	9.8	S
WHITE QUEEN	2W-W	1955 - 1963	14	15	443	1.87	SB	42	10.0	2.1	3.8	A	M	nd	
WHITE SAIL	4W-W	1968 - 1984	57	58	249	2.71	S	48	7.8	0.5	nd	A	M	10.8	
WHITE SPIRE	2W-W	1972 - 1980	40	30	89	2.04	SB	39	10.5	1.9	4.4	A	G	9.0	
WHITE SPRITE	3W-W	1978 - 1978	-31	43	192	2.05	SB	41	10.0	1.0	2.5	A	VG	8.4	
WHITE TARTAR	1W-W														
WHITE TRIUMPHATOR	1W-W	1955 - 1963	29	23	265	1.33	S	39	12.0	5.5	5.2	A	VG	nd	
WHITE WAX	1W-W	1964 - 1967	51	40	nd	nd	S	27	9.5	4.3	3.7	B	G	nd	
WHITEHEAD	2W-W	1970 - 1985	33	35	156	1.80	S	40	10.8	3.1	3.7	A	VG	10.4	
WHITELEY GEM	2Y-O	1955 - 1967	-1	2	212	2.55	SB	44	9.0	2.6	3.1	A	M	8.8	
WHITE'S HYBRID	8W-YYR														
WHITEWELL	2W-Y	1955 - 1963	-7	22	206	1.87	S	46	10.0	1.5	2.8	A	M	nd	
WIELAND WAGNER	2W-Y														
WILD ROSE	2W-P	1955 - 1963	90	39	192	1.87	SB	34	10.0	1.9	2.7	B	M	nd	
WILL. SCARLETT	2W-O	1964 - 1967	40	51	169	1.50	SB	32	8.5	1.6	3.2	B	M	6.0	
WILLEM MENGELBERG	1	1964 - 1967	-2	12	172	1.60	S	34	13.5	5.8	5.7	A	VG	7.4	
WILLEMSTAD	2W-O	1964 - 1967	34	22	336	2.90	SB	40	10.0	1.6	3.4	A	G	8.0	
WILLIAM FARMER	2Y-Y	1982 - 1984	51	20	nd	1.45	S	34	11.8	4.5	5.3	A	VG	13.0	
WILLIAM THE SILENT	1Y-Y	1964 - 1967	10	4	186	2.20	S	33	11.5	5.7	5.6	A	G	7.0	B
WILLOWFIELD	3W-R	1964 - 1967	11	36	342	2.50	SB	45	11.0	1.4	3.3	A	G	9.6	
WIN ALL	3Y-YYR	1972 - 1972	45	23	64	1.33	SB	36	11.0	1.3	4.3	A	G	9.6	
WINCHESTER	2Y-YOO	1968 - 1981	7	22	176	2.54	SB	44	10.0	2.2	3.1	A	G	11.0	S
WINDBLOWN	4W-Y	1979 - 1979	5	22	143	1.69	S	35	10.0	3.6	nd	B	M	7.6	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
WINGADDEE	2W-OOY	1973 - 1987	-3	6	167	2.37	B	46	9.3	2.2	3.3	A	G	8.5	
WINGED VICTORY															
WINIFRED VAN GRAVEN	3W-YJR	1964 - 1984	11	25	230	2.13	SB	40	10.4	1.2	2.8	A	G	9.1	
WINKFIELD'S DOWER	1Y-Y	1955 - 1963	-11	15	267	1.87	S	44	11.0	5.0	5.2	B	G	nd	
WINNIE WEEDON	9W-YJR	1955 - 1963	-7	22	349	1.87	SB	42	8.0	0.7	2.2	A	P	nd	
WINNIPEG	2Y-YYO	1968 - 1982	57	34	185	2.38	SB	40	10.8	1.8	3.8	A	G	8.8	S
WINTER BELLS	8W-Y	1981 - 1981	-53	24	nd	0.33	SB	27	4.5	0.5	1.2	nd	M	nd	
WINTER GOLD	1Y-Y	1955 - 1967	7	1	288	2.60	S	38	9.0	4.2	3.7	A	M	7.8	
WINTERMOON	1W-Y	1976 - 1980	-8	15	187	1.26	S	38	12.2	4.9	5.0	A	G	10.4	B
WINWICK	1W-W	1977 - 1981	23	7	165	1.82	S	33	10.3	4.1	4.0	A	G	8.2	B
WISDOM	3W-W	1955 - 1981	24	37	211	2.43	SB	46	10.0	3.2	1.2	A	G	9.0	
WODAN															
WOODCOCK	6Y-Y	1955 - 1967	-6	6	299	2.93	S	32	11.0	4.1	3.3	B	G	10.4	
WOODGREEN	2W-Y	1967 - 1987	31	-1	151	2.08	S	37	11.2	4.1	3.6	A	G	10.1	B
WOODLAND PRINCE	3W-Y	1979 - 1985	13	35	212	2.38	B	42	9.9	1.5	3.0	A	G	9.2	
WOODLAND STAR	3W-R	1979 - 1985	-7	42	366	1.61	S	43	9.2	1.0	2.8	A	VG	8.3	V
WOODPECKER	2W	1968 - 1972	60	36	176	1.80	SB	38	11.2	2.0	3.2	A	G	9.7	
WOODVALE	2W-WWY	1968 - 1977	56	34	144	2.08	S	37	11.2	4.1	3.6	B	VG	8.8	S
WOONAN	2Y-R	1976 - 1977	40	22	201	2.31	S	46	9.0	1.6	2.6	A	M	7.8	
WRESTLER	1Y-Y	1955 - 1963	0	7	218	1.87	S	38	10.0	4.7	5.4	A	G	nd	
YALTA	3W-YJR	1955 - 1980	6	28	203	2.03	SB	56	10.0	1.2	3.4	A	VG	7.0	
YANKEE CLIPPER	2Y-YYO	1968 - 1981	38	19	159	2.56	SB	41	11.0	2.2	4.3	A	M	8.1	
YELLOW BEAUTY	1Y-Y	1955 - 1963	7	1	324	2.40	S	44	11.0	5.0	4.0	A	M	nd	
YELLOW CHEERFULNESS	4Y-Y	1955 - 1984	24	41	220	2.45	E	42	5.8	0.6	nd	A	G	9.4	
YELLOW CLOUD	1	1964 - 1967	26	10	149	2.00	S	31	11.5	5.1	5.5	A	G	7.0	
YELLOW DAZZLER	1Y-Y	1972 - 1987	20	26	146	2.15	S	53	11.7	4.3	4.0	A	G	9.7	S

Cultivar	Division	Years in trial	Emergence date	First flower date	% weight increase	Flowers per bulb	Leaf poise	Stem length (cm)	Perianth diam. (cm)	Corona length (cm)	Corona diam. (cm)	Flower position	Flower substance	Vase-life (days)	Disease notes
YELLOW IDOL	1Y-Y	1964 - 1967	-5	24	158	1.10	S	38	11.5	5.0	4.5	A	VG	8.0	
YELLOW MOON	2Y-Y	1964 - 1976	27	27	182	2.31	SB	47	9.5	2.6	3.6	A	G	8.2	V
YELLOW POPPY	2Y-YYO	1955 - 1963	21	26	156	2.40	SB	44	10.0	1.8	3.0	A	M	nd	
YELLOW SUN	2Y-Y	1955 - 1981	-4	4	181	2.49	SB	39	12.3	4.4	4.8	B	G	10.6	
YELLOW SUNSET	1Y-Y	1978 - 1982	15	12	227	1.06	S	34	11.5	5.3	5.7	A	G	10.6	
YELLOW TRIUMPHATOR	1	1976 - 1981	20	15	121	2.16	S	40	13.5	5.6	6.0	A	G	9.0	V
YEO	2W	1977 - 1978	12	8	150	1.83	S	48	10.8	1.8	3.9	A	M	10.6	
YORNUP	2Y-Y	1970 - 1987	-23	-7	171	2.34	E	49	12.2	2.9	4.8	nd	P	8.5	
ZANITA	7Y-GYY	1964 - 1967	-28	37	nd	nd	SB	44	8.0	1.3	2.3	B	M	nd	
ZARAH LEANDER	2Y-R	1955 - 1963	14	31	264	1.87	SB	46	10.0	1.7	3.7	A	VG	nd	
ZEPHYR	2														
ZERO	2W-W	1964 - 1967	5	22	nd	nd	S	34	12.5	4.5	4.5	B	VG	8.0	
ZEST	1W-Y														
ZIRCON	2W-Y	1972 - 1980	13	9	154	2.42	SB	41	11.4	3.2	4.8	A	G	8.9	

FLOWER DESCRIPTION

The proportions of the flowers can be gauged from the measurements of perianth diameter and corona (trumpet or cup) diameter and length (see table), but they also need to be considered in conjunction with the written descriptions. Cultivars with pointed, star-shaped perianths will have an actual diameter far greater than it appears to be (eg, Sun Chariot). If the segments do not overlap, the perianth will appear thin and poor (eg, Orange Standard). Segments need to be stiff enough to avoid being floppy. For comparison of flower shapes, see Figures 1 and 2.

Generally speaking, a well balanced trumpet cultivar (Division 1) has a corona which is roughly as long as it is wide (eg, Dungannon, Golden Cockerel). Cultivars with very broad mouths do not look as attractive (eg, Magnificence, Dinton Giant). The perianth should give the trumpet a good background and a diameter of 10 to 12 cm is satisfactory, provided the segments are broad enough to overlap each other.

Large-cup cultivars (Division 2) vary considerably in shape and size, many having perianths as large as those of trumpet cultivars, and a diameter of 10 to 12 cm is again satisfactory, provided that the segments are broad enough to overlap each other. In Division 2, yellow-cupped cultivars have coronas which do not vary greatly in shape, and they are often of trumpet proportions (eg, Orange Leitrim). Orange or red-cupped cultivars vary greatly in size and shape, and may be straight sided (eg, Red Idol, Ceylon, Cardigan), bowl-shaped (eg, Balalaika, Bombay), or almost flat (eg, Blair House, Ice Follies). There are considerable differences in the form of the edge of the corona, which may be crenate (eg, Dinton Giant, Ark Royal), lightly frilled (eg, Coral Island, Forthright), lobed (eg, Corofin, Ann Abbott), or heavily and deeply frilled (eg, Ice Follies, Joke Fulmer, Princess Beatrix and Belisana).

Small-cup cultivars (Division 3) do not vary so much in size and shape. They usually have smaller perianths, and a diameter of 10 cm is a suitable size for market if the segments are broad enough to overlap each other. It is important that they should have well-shaped perianths because the coronas are so small. They usually look well balanced whether the corona is bowl-shaped, flat or straight-sided.

Double cultivars (Division 4) with one flower to a stem need to be of a reasonable size, with a stem strong enough to support the weight. A diameter of 10 to 11 cm is suitable, and the flower itself should be neat with well arranged segments (eg, White Lion, Gay Time). Cultivars with more than one flower per stem can have small individual flowers (eg, Cheerfulness).

Tazetta cultivars (Division 8) with only two or three florets per stem (eg, St Agnes, Glorious) need florets of reasonable size, not less than 7 cm diameter. Cultivars with many florets (eg, Silver Chimes, Geranium) are acceptable with smaller florets provided they are over 5 cm diameter.

Poeticus cultivars (Division 9) are usually rather small, but if the perianth segments are rounded and overlap well they will look quite substantial. The smallest acceptable diameter is 8 cm.

The descriptions given in this report are of mature flowers, usually picked when the flowers were showing colour. If flowers are picked very green (pencil stage), the perianth colours would often be paler and the corona colours not so intense or poorly developed. 'Crepey' refers to the texture of the perianth segments. 'Hooded' refers to cultivars whose perianths incurve and do not flatten out even when mature. 'Fluted' refers to the corona where it is creased vertically from the edge to the centre; this usually occurs in cultivars of Divisions 2 and 3.

The RHS Colour Chart is not used to describe colours since narcissus often vary and change colour with age and temperature. The area in which they are grown can also influence the colour, particularly with some orange, red and pink cups. Colour is often more intense where growth is slow but regular (eg, best in the Isles of Scilly, good in Cornwall but less intense in Eastern England). The names of some colours used are 'primrose' - soft pale yellow; 'sulphur' - pale creamy yellow; 'lemon' - sharp clear yellow; 'gold' - rich yellow tinged orange; 'poeticus white' - pure sparkling white as found in most poeticus hybrids; 'burnet orange' - orange with coppery tone. Other colour names are self-explanatory. Colours of coronas are usually described from the centre or base leading out to the edge. Where the colours are described as 'shading' (eg, Bombay), the base colour merges gradually into the colour at the edge. Where the colour is described as 'changing' (eg, Corofin) the base colour changes abruptly to the colour at the edge.

Colour development can vary considerably when the flower opens on the plant out of doors; certain conditions of warmth and a fairly high humidity seem to be required for full colouring, particularly for those flowers opening with lemon coronas and developing into cream or buff (eg, Spellbinder). Flowers cut and placed in water in a warm room will usually develop their colour more rapidly and reliably than those left on the plant. Many white trumpets and cups open yellow and develop cream or white, often more successfully in water than on the plant (eg, Mount Hood, Woodvale and Kilrea). Conversely, some red cups will intensify their colour with age often better on the plant than in water (eg, Fortune and Fortright). The perianths of many flowers open deep cream and develop to white. This usually happens quite rapidly regardless of whether they are left on the plant or are in water (eg, Bravura, Actaea, Corofin, Gay Time and Belisana). The perianths of some yellow cultivars develop with age to cream, sometimes a muddy colour, usually leaving a yellow base to the perianth (eg, Hollywood, Golden Bracelet and Rumba). This development of colour is perhaps more noticeable in water than on the plant.

FLOWER DESCRIPTIONS

- ABABA** 2W-O
Perianth white tinged with pink, twisted. Segments pointed, overlapping at base. Corona orange, slightly paler at base, short, open mouthed, crenate, frilly at rim. Overall appearance crumpled. Neck grows badly in water.
- ABADAN** 2W-YRR
Perianth pure white with sheen, creased, curled in at edges. Segments flat, rounded, well overlapping. Corona deep lemon with bright orange-red rim for 1/3 its length, fairly flat, fluted, deeply frilled.
- ABALONE** 2W-YYP
Perianth pale lemon. Segments broad based and overlapping. Segments broadly ovate with central ridge. Corona yellow, flared and frilled. Neck grows badly in water.
- ABEL TASMAN** 11W-WYO
Perianth white. Segments broad based and overlapping. Segments broadly ovate, curling over at edges. Split corona segments white with central yellow streak flattened against perianth. Neck grows badly in water.
- ABELARD** 2Y-YOO
Perianth deep primrose yellow, creased. Segments flat, pointed, overlapping. Corona orange, shading a little paler at base, fluted, with white blotches on edge.
- ACCENT** 2W-P
Perianth white. Segments broad and overlapping with central ridge. Corona frilled and rolled yellow/orange going to pink at edges. Neck grows in water.
- ACCOLADE** 3W-R
Perianth white. Segments broad, slightly pointed, overlapping and curled over at edges. Corona cup shaped and burnt orange in colour. Neck grows in water.
- ACROPOLIS** 4W-R
Perianth white. Segments broad, round and overlapping. Double perianth segments interspersed with orange petaloids. Neck grows in water.
- ACTAEA** 9W-YRR
Perianth white, stained yellow base, creased, sometimes buckled and folded back at edges. Segments flat, rounded, well overlapping. Corona deep lemon with bright orange-red rim for 1/3 its length, fairly flat, fluted, deeply frilled. Neck grows very badly in water.
- ACTRICE** 2Y-O
Perianth primrose yellow, broad ovate overlapping segments. Corona orange frilled cup. Neck grows in water.
- ADA FINCH** 1W-W
Perianth creamy white, creased, crêpey. Segments slightly twisted, pointed, overlapping. Corona opens creamy lemon, develops cream with some white blotches on the edge, deeply cut crenate, flanged. Neck grows in water.
- ADAM** 3Y-O
Perianth opens soft pale sulphur yellow, develops cream, slightly creased, edges curl in. Segments flat, rounded well overlapping. Corona deep orange-red. Neck grows in water. Stiff and sturdy.
- ADAMANT** 2Y-YYO
Perianth orange-gold, crêpey texture. Segments slightly twisted, pointed, overlapping, one segment often caught in cup. Corona bright orange, almost straight.
- ADELA** 2W-Y
Perianth white, stained yellow at base, creased, segments inclined to fold back lengthways, pointed outline. Segments flat, pointed, overlapping. Corona clear pale lemon yellow.
- ADMIRATION** 8Y-ORR
Perianth primrose yellow, very creased, curled back at edges. Segments slightly twisted, rounded, overlapping. Corona bright orange red with greenish gold base, fluted.
- ADORNMENT** 2Y-O
Perianth canary yellow. Segments broad ovate with central ridge. Corona apricot colour flared and frilled. Neck grows badly in water.
- ADVANCE GUARD** 1Y-Y
Perianth greenish yellow with green back. Segments twisted, pointed, separated. Corona deep golden yellow, widespread mouth, deeply cut edge.
- ADVENTURE** 2Y-Y
Perianth soft sulphur yellow, slightly creased. Segments slightly twisted, pointed, broad, overlapping. Corona bright clear lemon yellow, of almost trumpet proportion.

- AEROLITE** 2Y-Y
Perianth pale creamy sulphur yellow, slightly creased. Segments flat, pointed, broad, overlapping. Corona clear pale lemon yellow, of almost trumpet proportion.
- AFLAME** 3W-YOO
Perianth creamy white, stained yellow at base. Segments flat, pointed overlapping. Corona bright deep red shading to orange base, fluted. Corona edge dies first in water.
- AFTER ALL** 3W-YYR
Perianth white. Segments broad based and pointed. Corona medium sized flared cup; canary yellow, with orange frilled edge. Neck grows in water.
- AGATHON** 2Y-Y
Perianth deep primrose yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona deep clear lemon yellow, open mouthed. Well proportioned.
- AGNES ANDREAE** 2W-O
Perianth poeticus white, creased, edges curl in, crêpey texture, circular outline. Segments flat, rounded, broad, well overlapping. Corona apricot orange with narrow rim of creamy apricot, deeply fluted, almost flat. Neck grows in water.
- AGNES MONTEFIORE** 2W-WWY
Perianth creamy white stained yellow at base, slightly creased, pointed outline. Segments slightly twisted, pointed, overlapping. Corona bright clear yellow, develops cream, open mouthed. Neck grows a little in water.
- AGORA** 2W-O
Perianth poeticus white, smooth, edges slightly incurled. Segments broad, rounded, overlapping at base. Corona soft orange, flat, only slightly lobed. Neck grows badly in water.
- AHOY** 11W-Y
Perianth creamy white, pointed outline, top three segments incurved. Segments flat, pointed separated. Corona sulphur yellow, split into six segments, incurved. Top heavy appearance.
- AIR MARSHALL** 2Y-O
Discarded - Very low flower production.
- AIRCASTLE** 3W-Y
Perianth white. Segments broadly ovate and overlapping. Corona yellow getting paler towards edge. Corona flared and frilled. Neck grows badly in water.
- ALABASTER** 4W-W
Perianth poeticus white. Segments smooth, round and 1/2 overlapping. - green at back of base - raised centre line - inner segments tubular and buckled, mixed with short corona segments. Corona white, with greenish tinge. Frilled, tightly packed. Neck grows badly in water.
- ALAMEIN** 2Y-O
Perianth bright clear lemon yellow, crêpey texture, odd segments creased and caught in corona. Segments flat, rounded, well overlapping. Corona deep bright orange red, bright colour contrast. Neck grows in water.
- ALARM** 2Y-O
Discarded - Short.
- ALAYNE** 2W-YYO
Perianth creamy white, deeply creased, often notched, sometimes buckled. Segments flat, pointed, overlapping. Corona light greenish lemon base 1/2 shading to bright orange edge with some white blotches on edge, fluted bowl shape. Neck grows badly in water.
- ALBANY** 8W-YOO
Perianth creamy white, stained yellow at base. Segments flat, pointed, broad, overlapping. Corona bright deep orange shading to orange gold base with slight greenish gold eye. Neck grows in water.
- ALBATROSS** 3W-YYO
Perianth creamy white, creased, curled in at edges. Segments twisted, pointed, separated. Corona greenish lemon with thin line of bright orange on edge, fluted. Neck grows in water.
- ALBERT SCHWEIZER** 1Y-Y
Discarded - Rough flower.
- ALCATRAZ** 1Y-Y
No description available.
- ALDERGROVE** 2W-Y
Discarded - Short.
- ALEPPO** 3W-R
Perianth creamy white, stained yellow at base, creased. Segments flat, rounded, well overlapping. Corona orange gold changing to bright orange red edge for half its length, fluted. Neck grows in water.

- ALEX FLEMING** 2W-O
Perianth white. Segments broadly ovate, overlapping and with a central ridge. Corona medium sized frilled orange cup. Neck grows badly in water.
- ALGECIRAS** 3W-R
Perianth white, smooth, edges curl in, circular outline. Segments flat, rounded, well overlapping. Corona deep orange red, with small green gold eye, fluted, almost flat. Neck grows stiffly in water.
- ALGIERS** 3W-O
Perianth opens cream develops white, stained yellow base, creased, edges curl in at tips, pointed outline. Segments flat, pointed, overlapping. Corona bright orange-red fluted, frilled, lobed, bowl shape. Neck grows stiffly in water.
- ALICANTE** 2W-O
Perianth creamy white, stained yellow base, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona deep orange-red throughout, occasional white blotches on edge, fluted, frilled, lobed, bowl shape. Neck grown badly but stiffly in water.
- ALIGHT** 3Y-O
Perianth soft yellow, suffused apricot at base, slightly creased. Segments flat, pointed, overlapping. Corona deep burnt orange, fluted. Stiff.
- ALL GLORY** 1W-Y
Perianth muddy creamy white, creased, crêpey. Segments flat, pointed, overlapping. Corona creamy lemon, crenate, flanged. Neck grows in water.
- ALLUREMENT** 2W-P
Perianth poeticus white, slightly twisted. Segments broad, rounded, overlapped well, some notched. Corona yellow at base becoming salmon pink, flat, rim deeply cut and frilled. Neck grows in water.
- ALPHONS DIEPENBROCK** 1Y-Y
Discarded - Untidy flower, poor poise.
- ALPINE EAGLE** 1W-Y
Perianth creamy white, creased, pointed outline. Segments flat, pointed, overlapping. Corona creamy primrose, rolled back edge, rather untidy.
- ALROI** 2Y-Y
Perianth soft pale lemon yellow, crêpey texture. Segments flat, pointed, broad, well overlapping. Corona deep clear lemon yellow. Neck very short and brittle.
- ALTYRE** 3W-W
Perianth creamy white, crêpey texture, edges curl in. Segments flat, pointed, broad, overlapping. Corona cream with narrow deeper cream edge, fluted. Hangs head badly, floppy. Neck grows very badly in water.
- ALVERADO** 2W-Y
No description available.
- ALYCIDON** 1W-W
Perianth creamy white, slightly creased. Segments flat, pointed, broad, overlapping, shovel shape. Corona cream, lightly frilled, flanged, neat edge.
- AMATEUR** 3W-O
Perianth white, creased, one segment often buckled, reflexed, circular outline. Segments flat, rounded, well overlapping. Corona deep apricot orange, deeply fluted, very flat.
- AMBASSADOR** 1W-W
Perianth creamy white, cream at base, a little creased. Segments flat, pointed, broad, well overlapping. Corona cream, neat.
- AMBER FLASH** 2W-O
Discarded - Flowers below foliage.
- AMBERGATE** 2O-O
Perianth gold, edges curl in at tips. Segments flat, overlapping. Corona orange-red, short and flat, slightly frilled. Neck grows badly in water.
- AMBERLEY** 2Y-Y
Perianth bright clear lemon yellow, smooth, pointed outline. Segments flat, pointed, overlapping. Corona deep clear lemon yellow. Very stiff and sturdy.
- AMBOYNA** 1W-Y
Perianth creamy white, stained yellow at base. Segments twisted pointed, overlapping. Corona light lemon yellow, edge deeply cut and lobed.
- AMIABLE** 3Y-O
Perianth primrose yellow, creased, top 3 segments flat, lower 3 segments reflexed. Segments flat, pointed, overlapping. Corona deep burnt orange, flat, deeply fluted.

AMOR Perianth creamy white, stained at base. Outer segments broad, inner segments narrower and curled inwards, smooth, round and overlapping. Corona rich lemon. Flat cup shaped, fluted and deeply lobed. Neck grows in water.	3W-YYO
ANACAPRI Perianth white, stained yellow base, circular outline. Segments rounded, overlapping at base, bottom segments broad, smooth, curled in at edges, top segments wrinkled bottom edges. Corona deep warm yellow, top one-third rich orange, expanded, finely fluted. Neck grows badly in water.	3W-YYR
ANAKIM Discarded - Badly proportioned flower, flowers below foliage.	1W-Y
ANDALUSIA Discarded - Short, small, cyclamineus hybrid.	6Y-O
ANDES Discarded - Short, very susceptible to basal rot.	1W-W
ANDROCLES Perianth lemon yellow - paler at tips. Segments pointed, crêpey and slightly inflexed. Corona canary yellow, flared frilled and funnel shaped. Neck grows badly in water.	4W-W
ANGELINE Perianth creamy white, edges curl in, a little creased, one segment often buckled. Segments flat, pointed, overlapping. Corona greyish cream with thin pencil line of bright apricot orange, slightly fluted, bowl shape. Damages easily in wind.	3W-WWY
ANGEL'S WINGS Perianth white, some green on back, a little creased. Segments flat, pointed, broad, well overlapping. Corona open primrose yellow, quickly develops to pure white.	2W-W
ANN ABBOTT Perianth white, sometimes notched and buckled. Segments slightly twisted, rounded, overlapping. Corona creamy white tinged peach with narrow band of bright salmon, fluted, frilled, deeply lobed, deep bowl shape. Stiff. Neck grows in water.	2W-P
ANNAPURNA Perianth creamy white, creased, crêpey, pointed outline. Segments flat, pointed, overlapping. Corona creamy white, untidy edge. Rough and coarse.	1W-W
ANNE FRANK No description available.	4W-O
ANTRON Perianth bright yellow. Segments broad and overlapping. Corona small, frilled cup and dark orange. Neck grows badly in water.	2Y
ANTWERP Perianth deep clear lemon yellow, crêpey texture. Segments pointed, broad, overlapping, shovel shape. Corona deeper lemon yellow, rather straight sided.	1Y-Y
ANZENA Discarded - Short, rough, flowers below foliage.	2W-Y
APPROVAL Perianth sulphur, slightly creased. Segments flat, pointed, overlapping, top 3 segments inclined to incurve, edges curl in. Corona bright orange shading to deeper orange red edge.	2Y-OOR
APRICOT BEAUTY Perianth creamy white, creased, pointed outline. Segments flat, pointed, overlapping. Corona cream shading to narrow apricot edge, fluted, very broad mouth, rough, badly proportioned. Neck grows in water.	2W-O
APRICOT DISTINCTION Perianth opens yellow, suffused pinkish red, develops quickly to pale apricot, with paler tips and centre stripe. Segments flat, rounded, overlapping. Corona deep burnt orange red, flat, deeply fluted.	3Y-R
APRICOT SENSATION Perianth creamy white. Segments broad and well overlapped. Corona frilled/flared. Corona orange with a lemon yellow centre and a pale yellow edge. Neck grows badly in water.	2W-Y
APRIL HARVEST Perianth white. Outer segments very broadly ovate, inner ovate, and overlapping. Corona large flared frilled trumpet pale yellow in colour.	1W-Y
APRIL MERIT Perianth white. Segments broad and overlapping. Corona large, flared trumpet white and frilled. Neck grows in water.	2W-W

- APRIL MESSAGE** 1W-W
Perianth white. Segments broad and well overlapped. Corona flared trumpet with frilled edge. Neck grows in water. Weak stems.
- APRIL ROSE** 2W-P
Perianth white. Segments broad and overlapping. Corona large flared funnel with frilled edge. Corona pink, darker at rim, with yellow eye. Neck grows in water.
- APRIL TEARS** 5Y-Y
Perianth sulphur yellow, smooth, reflexed. Segments flat, rounded, barely overlapping. Corona creamy sulphur yellow, shade paler at base, smooth, goblet shape. Neat, well balanced.
- ARANJUEZ** 2Y-YYO
Perianth soft sulphur yellow, creased, one segment often buckled. Segments flat, rounded, well overlapping. Corona gold, with edge of bright orange gold for 1/3 of its length.
- ARBAR** 2W-O
Perianth white. Segments broad and slightly overlapping. Corona flared, frilled, canary yellow with orange band near rim. Rim canary yellow.
- ARCADIA** 2W-O
Perianth cream, smooth, stained yellow at base. Segments rounded, some notched, overlapping. Corona deep orange, short, open mouthed to flat, fluted giving wavy edge. Neck grows in water.
- ARCTIC DORIC** 2W-WW
Perianth white. Segments broad and well overlapping. Corona open mouthed and frilled also white in colour.
- ARCTIC GOLD** 1Y-Y
Perianth rich bright gold, double-triangle shape. Segments rounded, overlapping at base; bottom segments broad, very smooth, curled in at edges, top segments twisted. Corona dark warm gold, paler at base, fluted, crenate edge, slender. Scented. Neck grows badly in water. Weak stems.
- ARCTIC MOON** 2W-W
Perianth creamy white, creased, often notched. Segments flat, pointed, overlapping. Corona opens primrose, develops buff then becomes deep cream, crenate, flanged. Stiff.
- ARCTIC PRINCE** 1W-W
Perianth white, creased. Segments flat, pointed, overlapping. Corona white, frilled. Neck grows in water.
- ARDARA** 2W-W
Perianth creamy white, slightly creased, crêpey, edges curl in a little, pointed outline. Segments flat, pointed, overlapping. Corona pale cream, slightly fluted, frilled, lobed. Stiff.
- ARDBANE** 2W-W
Perianth white. Segments broad and well overlapping. Corona open mouthed and frilled - white in colour.
- ARDELINIS** 1W-W
Perianth creamy white, perfect double triangle. Segments flat, pointed, broad, overlapping. Corona creamy white with rolled-back rim. Well balanced.
- ARDOUR** 3Y-R
Perianth canary yellow, paler at tips. Segments broad, well overlapping and slightly curled over at edges. Corona flared and frilled, bowl-shaped and dark orange in colour. Neck grows in water.
- ARGENT** 4W-Y
Perianth segments cream, flat, pointed, barely overlapping, two complete rows. Corona segments pale gold, narrow, twisted, concentrated in centre of flower. Weak stem. Neck grows badly in water.
- ARGOSY** 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona flared funnel, frilled and same colour as perianth.
- ARGUROS** 3W-GOR
Perianth creamy white, stained yellow base, creased, one segment often buckled. Segments flat, pointed, overlapping. Corona moss green base, shading through orange to lemon with narrow orange rim, fluted. Neck grows badly in water. Poor poise.
- ARGYLL** 1Y-Y
Perianth pale lemon, slightly creased, crêpey. Segments slightly twisted, pointed, overlapping. Corona deeper lemon, fairly broad mouth, frilled, flanged. Stiff.
- ARK ROYAL** 1Y-Y
Perianth bright lemon, slightly creased, crêpey. Segments flat, pointed, broad, overlapping. Corona deeper lemon, crenate, flanged. Stiff.

ARLINGTON	2W-YOY
Perianth creamy white, one segment creased. Segments flat, rounded, overlapping. Corona bright orange shading to lemon base with thin gold edge, almost flat, fluted, deeply frilled. A large proportion of flowers have too many segments.	
ARMADA	2Y-O
Perianth clear lemon yellow, slightly creased. Segments flat, pointed, overlapping. Corona bright burnt orange throughout, ageing to deeper colour, bowl shape. Stiff and sturdy.	
ARMAGH	1Y-Y
Perianth lemon yellow. Segments broad and overlapping. Corona open mouthed, flared and frilled, same colour as perianth. Neck grows in water.	
ARNHEM	3Y-O
Perianth opens primrose, develops cream stained yellow base, paler at tips, creased, crêpey, circular outline. Segments flat, rounded, overlapping. Corona deep orange red, fluted, frilled, lobed, almost flat.	
ARRANMORE	1Y-Y
Discarded - Short.	
ARRIBA	2Y-O
Discarded - Flowers below foliage.	
ARTILLERY	3W-R
Perianth poeticus white, stained yellow at base, smooth. Segments incurling, some notched, would fit into a double triangle, ovate. Corona bright orange, flat, crenate with ragged edge. Flower faces upwards. Neck grows in water.	
ARTOIS	1Y-Y
Discarded - Short, badly proportioned.	
ARUBA	2Y
Discarded - Short, flowers below foliage.	
ASCOT	4Y-Y
Perianth canary/primrose yellow - paler at tips. Segments 1 1/2 complete rows. Rest intermixed with corona. Corona orange/yellow mixed - loose centre but symmetrical. Segments round and well overlapping. Corona edge frilled. Weak neck. Neck grows in water.	
ASKELON	1W-W
Perianth white, creased. Segments flat, pointed, overlapping. Corona cream, neat shape.	
ASPASIA	8W-Y
Perianth creamy white, slightly creased. Segments flat, rounded, overlapping. Corona clear lemon yellow, slightly deeper at edge, almost flat, fluted. Style half way between short and long stamens. Neck grows in water.	
ATATURK	2Y-O
Discarded - Rough, perianth catches in corona.	
ATHFORD	1Y-Y
Perianth clear lemon yellow, creased. Segments flat, pointed, overlapping. Corona deeper lemon yellow.	
ALTAS	1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona large flared frilled funnel. Same colour as perianth.	
ATTRACTIVE	1Y-Y
No description available.	
AUDA	4Y-Y
Perianth lemon yellow. Double perianth segments broad and overlapping. Perianth segments intermixed with corona segments. Corona colour canary yellow. Neck grows badly in water.	
AUDUBON	3W-WWR
Perianth white. Segments broad and overlapping. Corona flared, frilled cup, white with an orange rim. Neck grows badly in water.	
AUGUST GREEN	1W-Y
No description available.	
AUGUST PINK	2W-P
Perianth white. Segments broad and overlapping. Corona open mouthed, flared and frilled. Corona pink with yellow eye. Neck grows badly in water.	
AURELIA	7Y-Y
Perianth clear lemon yellow, smooth. Segments flat, pointed, overlapping. Corona deeper clear lemon yellow, very smooth and neat. Neck grows in water.	

- AVALANCHE** 8W-Y
Discarded - Better suited to Isles of Scilly.
- AVE** 2W-W
Perianth white, very smooth. Segments flat, pointed, broad, overlapping. Corona opens creamy primrose with paler edge developing to creamy white throughout, with clean cut edge.
- AVENGER** 2W-O
Perianth white, flat, edges slightly incurling. Segments broad, rounded and overlapping. Corona bright orange, flat, crenate, edge slightly notched. Flower tends to hang down. Neck grows badly in water.
- AVIEMORE** 2Y-YOO
Perianth pale clear sulphur yellow, with paler tips, slightly creased, edges curl in a little. Segments flat, pointed, overlapping. Corona gold shading to orange edge for 2/3 of its length, a little fluted.
- AYALA** 2Y-O
Perianth soft pale lemon, overlaid unevenly with bright gold. Segments well rounded, curved in at edges, overlapping. Corona deep orange with reddish tinge, fluted, bowl-shaped.
- BABY DOLL** 6Y-Y
Perianth bright lemon yellow, reflexed, smooth. Segments flat, pointed, broad, well overlapping. Corona deeper bright lemon yellow. Well proportioned.
- BABY MOON** 7Y-Y
Perianth deep lemon yellow, smooth. Segments flat, pointed, separated. Corona deeper golden yellow, smooth, almost flat.
- BACCARAT** 11Y-Y
Perianth sulphur yellow. Segments flat, pointed, broad, overlapping. Corona lemon yellow, split into 6 segments lying almost flat against the perianth.
- BACH** 1Y-Y
Perianth deep primrose, some green at back. Segments slightly twisted, pointed, overlapping. Corona deep rich gold, evenly crenate.
- BACKHOUSE'S GIANT** 2Y-O
Perianth sulphur yellow, some green on back, creased. Segments very twisted, pointed, barely overlapping, edges curl back. Corona bright solid orange throughout, trumpet shape, fluted, deeply lobed edge. Badly proportioned, neck grows badly in water.
- BADGER** 2Y-O
Perianth deep lemon tinged slight orange at base, creased, crêpey. Segments flat, pointed, overlapping. Corona deep burnt orange, fluted, almost straight sided, frilled. Neck grows badly in water.
- BAHRAM** 2Y-O
Perianth light lemon, developing paler at tips, creased, crêpey, edges curl in, pointed outline. Segments pointed, flat, overlapping. Corona deep burnt orange paler at base, fluted, lightly frilled, bowl shape. Neck grows stiffly in water.
- BALALAIKA** 2Y-YYR
Perianth pale clear lemon, slightly creased, circular outline, edges curl in. Segments flat, pointed, overlapping. Corona greenish gold with narrow rim of orange red, fluted, lightly frilled, bowl shape.
- BALAMARA** 2Y-Y
Perianth deep lemon, smooth. Segments small and rounded, overlapping at base. Corona deep lemon, flanged, lightly frilled. Neck grows badly in water.
- BALI HAI** 4W-R
Perianth white. Double perianth segments very broad and overlapping. Corona split and intermixed with perianth segments. Corona segments frilled, yellow with orange edge. Neck grows badly in water.
- BALLINTOY** 2Y-R
Perianth lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona bright orange red, a little fluted, bowl shape. Neck grows in water.
- BALLOO** 2W-W
Discarded - Short, rough flower.
- BALLYCASTLE** 3W-WWO
Perianth creamy white, edges curl in, creased, inclined to reflex. Segments flat, pointed, well overlapping. Corona small grey eye, changing to greyish cream, with bright apricot orange band for 1/3 its length, fluted, nearly flat. Neck grows in water.
- BALLYFERIS** 1W-Y
Perianth creamy white, yellowish on back, slightly creased. Segments flat, pointed, broad, well overlapping, shovel shaped, forming double triangle. Corona light lemon yellow, paler at base.

- BALLYGARVEY** 1W-Y
Perianth creamy white. Segments broad and overlapping. Corona open mouthed canary yellow and frilled.
- BALLYMARLOW** 2Y-R
Perianth light lemon yellow, crêpey texture. Segments flat, pointed, overlapping. Corona bright orange red throughout, fluted, goblet shape. Neck grows stiffly in water.
- BALLYMARTIN** 2W-GYY
Perianth white. Segments round, overlapping and reflexed. Corona flared, frilled cup with lemon yellow rim. Neck grows in water.
- BALLYSILLAN** 3Y-R
Perianth clear pale lemon yellow. Segments flat, pointed, overlapping. Corona deep orange red, fluted, bowl shape, good colour contrast.
- BALLYWALTER** 1W-Y
Perianth white. Segments broad, overlapping and inflexed. Corona lemon yellow, open mouthed and frilled. Neck grows in water.
- BAMBI** 1W-Y
Perianth greenish cream, star-shape, creased. Segments twisted, pointed, separated. Corona lemon yellow, shading paler at base, stiff.
- BAMBINO** 2Y-O
Perianth deep bright lemon yellow, smooth. Segments flat, pointed, overlapping. Corona bright orange, developing to deeper colour with age. Neat and stiff.
- BANBRIDGE** 1Y-Y
Perianth bright lemon yellow, smooth. Segments flat, pointed, well overlapping. Corona deeper lemon yellow, almost straight sided.
- BANDSMAN** 2Y-Y
Perianth pale lemon yellow, crêpey texture, rounded shape. Segments flat, pointed, broad, overlapping. Corona deep lemon yellow.
- BANN** 2W-P
Perianth creamy white, smooth. Segments flat, pointed, broad, overlapping. Corona opens lemon yellow with salmon pink rim, develops to pale pinky apricot with deeper rim, ages to pale buff cream with deeper edge. Well balanced.
- BANTAM** 2Y-OOR
Perianth lemon yellow, smooth, circular outline. Segments flat, rounded, well overlapping. Corona deep coppery red for half its length, shading to greenish orange base. Very small, neat and stiff.
- BANTRY** 3W-YYR
Perianth creamy white, odd segments creased. Segments flat, rounded, well overlapping. Corona greenish lemon, changing to deep bright orange edge for 1/3 of its length, sometimes white blotches on edge, fluted. Neck grows in water. Stiff.
- BARBARA ANN SCOTT** 3W-YYO
Perianth poeticus white, creased, edges curl in, often buckled. Segments flat, rounded, overlapping. Corona greenish lemon changing to orange edge for 1/5 of its length, fluted, saucer shape. Corona edge dies first.
- BARDIA** 2Y-O
Perianth primrose yellow, one segment often creased and caught in corona. Segments flat, rounded, broad, well overlapping. Corona bright orange red, a little paler at base, fluted.
- BARLOW** 6Y-YWW
Perianth canary yellow. Segments acute slightly overlapping and reflexed. Corona cylinder shaped, flared and frilled, same colour as perianth.
- BARRETT BROWNING** 3W-O
Perianth opens primrose, quickly develops creamy white, stained slight yellow at base, a little creased, circular outline. Segments flat, rounded, overlapping. Corona bright orange red throughout, fluted, bowl shape. Neck grows badly in water.
- BARTIZAN** 2Y-R
Perianth pale primrose yellow, crêpey texture, star shaped. Segments flat, pointed, barely overlapping. Corona bright orange red.
- BARTLEY** 6Y-Y
Discarded - Short, cyclamineus hybrid.
- BASHFUL** 2W-W
Perianth creamy white, triangular, one segment sometimes buckled. Segments flat, pointed, broad, overlapping. Corona opens creamy primrose, develops pinkish tone, then turns white. Stiff and sturdy.

BASTION 1Y-Y
Perianth very bright rich lemon yellow, smooth. Segments flat, pointed, overlapping. Corona deeper lemon yellow. Well balanced flower. Neck grows in water.

BATH'S FLAME 3Y-O
Perianth pale lemon yellow, fading with age, spidery looking. Segments twisted, pointed, barely overlapping. Corona deep lemon with brilliant red edge for 1/4 of its length.

BATTLEAXE 2Y-O
Perianth primrose yellow, a little paler at tips, creased and notched at edges. Segments flat, pointed, overlapping. Corona orange.

BAWNBOY 1Y-Y
Discarded - Flowers below foliage.

BAYARD 1Y-Y
Discarded - Small.

BAYEUX 1Y-Y
Perianth clear lemon yellow, slightly creased. Segments flat, pointed, overlapping. Corona deep lemon yellow, flanged, neat.

BEALITA 3Y-O
Perianth pale yellow. Segments broad and overlapping. Corona dark orange, frilled cup. Neck grows badly in water.

BEATEN GOLD 1Y-Y
Perianth deep lemon, a little creased, pointed outline. Segments slightly twisted, pointed, overlapping. Corona dark lemon, crenate.

BEAU MONDE 2W-O
Discarded - Short, rough, flowers below foliage, very short vase life.

BEDOUIIN 3W-O
Perianth cream, stained yellow at base, pointed outline. Segments flat, pointed, barely overlapping. Corona light gold shading to orange for half its length. Neck grows in water.

BEECHER STOWE 3W
Perianth creamy white, very creased, crêpey texture. Segments slightly twisted, rounded, well overlapping. Corona deep apricot orange, fluted.

BEEFEATER 2Y-O
Discarded - Perianth catches in corona, floppy.

BEERSHEBA 1W-W
Perianth white, pointed, star shaped, slightly creased. Segments flat, pointed, overlapping. Corona white, long and slender.

BEGUILDY 3Y-GGR
Discarded - Severe virus in this stock.

BEIRUT 2W-YYR
Perianth creamy white, stained yellow at base, creased. Segments flat, rounded, overlapping. Corona orange with deep red edge for 2/3 of its length, fluted.

BELISANA 2W-O
Perianth opens primrose, develops cream stained yellow for 1/3 at base, creased, crêpey, slightly reflexed. Segments flat, rounded, broad, well overlapping. Corona deep gold base shading to bright orange for half its length, variable colour, deeply fluted, thick double frilled edge, bowl shape. Stiff, bold.

BELLA VISTA 2W-ROO
Perianth white, stained yellow base, slightly creased, edges curl in a little, pointed outline. Segments flat, pointed, overlapping. Corona greenish lemon base, changing to band of bright orange red for 1/3 its length, occasional white blotches on edge, frilled, fluted, almost flat. Neck grows stiffly in water.

BELLE CHINOISE 3W-O
Perianth cream, overlaid apricot pink with white tips, colour fades immediately on opening, edges curl in, but flattens with age. Segments flat, pointed, barely overlapping. Corona deep orange with greenish orange eye, almost flat, crinkled and deeply fluted.

BELLEEK 2W-P
Perianth creamy white, creased, crêpey texture, sometimes caught in corona. Segments slightly twisted, pointed, overlapping. Corona opens primrose yellow, develops creamy peach, fades to cream with age, fluted, open mouth. Dumpy.

BELLEVUE	2W-O
Perianth white, a little creased, edges curl in, sometimes notched. Segments flat, pointed, overlapping. Corona bright apricot orange with slightly paler edge, occasional white blotches on edge, fluted, deep bowl shape. Poor poise. Neck grows badly in water.	
BELTANY	1Y-Y
Discarded - Small.	
BELTRIM	2W-GPP
No description available.	
BELYTA	2W
Discarded - Poor colour, rough, flowers below foliage.	
BENE	1Y-Y
No description available.	
BENEDICTION	3W-W
Perianth white. Segments broadly ovate and well overlapped. Corona small, frilled cup, white with green eye. Neck grows badly in water.	
BENGHAZI	2Y-YOO
Perianth bright lemon yellow, smooth. Segments flat, pointed, overlapping. Corona rich deep orange red throughout, fluted, straight sided. Neck grows stiffly in water.	
BERGEN	2W-W
Perianth creamy white. Segments flat, shovel shaped, overlapping. Corona opens cream, develops pure white with shaded cream base. Very stiff and sturdy.	
BERIMA	2Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona wide mouthed, frilled and same colour as perianth. Neck grows in water.	
BERMUDA	2Y-O
Perianth soft primrose yellow, crêpey texture, edges notched. Segments flat, pointed, broad, overlapping. Corona bright burnt orange shading to orange base, small. Neck grows in water. Flower rather large and floppy.	
BERNARDINO	2W-YYO
Perianth creamy white, touch of green and apricot at base, creased, floppy. Segments flat, pointed, barely overlapping. Corona greenish lemon shading to apricot edge for 1/3 of its length, with slightly paler edge. Neck grows in water.	
BERTHA ATEN	2W-O
Perianth creamy white, creased. Segments flat, rounded, well overlapping. Corona bright sealing-wax red, deep frilled and lobed right into centre, deeply fluted. Stiff and sturdy.	
BERYL	6W-YYO
Perianth pale creamy lemon, developing creamy white stained yellow at base, reflexed. Segments flat, pointed, overlapping. Corona gold shading to orange edge for 1/2 of its length. Very long style.	
BESSBROOK	2W-W
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and lemon yellow. Neck grows in water.	
BETTER TIMES	2Y-YYO
Perianth pale lemon yellow, creased, crêpey texture, one segment often buckled, inclined to reflex. Segments flat, pointed, broad, overlapping. Corona bright deep orange, shading to orange-gold base. Flower rather large and floppy.	
BIANCA	1W-W
Discarded - Rough flower, very susceptible to basal rot.	
BIBKIE	2W-Y
Perianth lemony white. Segments broad and overlapping. Corona frilled cup and lemon yellow in colour.	
BIG CHIEF	1Y-Y
No description available.	
BIG KEITH	1Y-Y
No description available.	
BIG WIG	4W-Y
Perianth white. Segments double, corona yellow, split and interspersed with white perianth segments. Neck grows badly in water.	
BILEE	1Y-Y
Perianth lemon yellow. Segments pointed and slightly overlapping. Corona large open mouthed, frilled, flared and canary yellow.	

- BILL WEBBER** 2W-O
Perianth white. Segments broad, ovate and overlapping. Corona canary yellow, open mouthed and frilled. Neck grows badly in water.
- BINDA** 1W-P
No description available.
- BINKIE** 2Y-W
Perianth pale sulphur yellow, fading back to almost white at the base. Segments pointed, broad, overlapping. Corona opens clear sulphur yellow then fades back to lemonade white with thin rim of yellow.
- BIRD OF DAWNING** 1Y-Y
Perianth deep lemon, slightly twisted. Segments pointed, overlapping. Corona deep lemon, straight with edge rolled back, only slightly frilled. Neck grows stiff in water.
- BIRMA** 3
Perianth soft lemon yellow, reflexed, edges curl inwards, smooth. Segments flat, pointed, overlapping. Corona brilliant deep red, fluted, goblet shape. Corona edge dies first.
- BISMARCK** 2W-W
Perianth creamy white, deeper at base. Segments flat, pointed, overlapping. Corona pale creamy lemon, paler at base. Neat.
- BITHYNIA** 3W-WWO
Perianth white. Segments broadly ovate and overlapping. Corona small bowl shaped and frilled, white with orange/yellow rim. Neck grows badly in water.
- BIZERTA** 2W-Y
Perianth white, one segment usually creased and buckled. Segments flat, pointed, broad, well overlapping. Corona opens lemon, develops gold shaded orange, ages to orange throughout - variable colour.
- BLACK PRINCE** 9W-GYR
Perianth pure white, stained slight yellow at base, reflexed. Segments flat, rounded, overlapping. Corona greenish lemon changing to deepest red edge for 1/5 of its length, with a greenish gold eye. Style longer than longest stamens.
- BLAIR HOUSE** 3W-O
Perianth white, slightly creased, circular outline. Segments flat, rounded, broad, well overlapping. Corona bright apricot orange with narrow rim of creamy apricot, fluted, frilled, lobed, nearly flat. Neck grows in water.
- BLANQUET** 4W-Y
Discarded - Small.
- BLARIS** 2W-P
Perianth white. Segments broad based, slightly pointed and overlapping. Corona frilled cup, dark pink in colour. Neck grows badly in water.
- BLARNEY** 3W-OOY
Perianth creamy white, smooth, star shaped. Segments flat, pointed, broad, overlapping. Corona bright apricot with narrow rim of buff apricot, almost flat, fluted. Neck grows in water.
- BLARNEY'S DAUGHTER** 2W-OOY
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona deep apricot orange with narrow rim of cream, fluted, lightly frilled, shallow bowl shape. Neck grows stiffly in water.
- BLEASBY GORSE** 1Y-Y
Perianth lemon, slightly twisted. Segments small, overlapping, would fit into a double triangle if straight. Corona deep lemon, long, narrow, flanged with ragged rim, deeply lobed. Hollow stems. Neck grows in water.
- BLLENHEIM** 1Y-Y
No description available.
- BLISS** 1W-W
Perianth creamy white, crêpey. Segments twisted, pointed, overlapping. Corona opens creamy lemon, develops cream, frilled, flanged, neat edge, trumpet proportion.
- BLOEMFONTEIN** 2Y-YYO
Perianth bright yellow. Segments broad and overlapping. Corona frilled, small open mouthed, yellow in centre and orange at rim. Neck grows badly in water.
- BLUSH** 2W-P
Perianth creamy white, deeply creased, edges curl in. Segments flat, pointed, overlapping. Corona pale creamy buff with faint tinge of pink, muddy colours.
- BLUSTER** 2Y-R
Perianth mid lemon suffused slight gold at base, developing paler at tips, slightly creased. Segments flat, pointed, overlapping, sometimes buckled. Corona bright orange, paler at base, fluted, lightly frilled, bowl shape. Neck grows badly in water.

- BO PEEP** 2W-P
Perianth white. Segments broad with slight overlap. Inner whorl has edges curled inwards. Corona small cup with frilled edge, colour - orange with light pink centre.
- BOBOLINK** 2W-YYO
Perianth cream, creased. Segments broad, coming to a point, well overlapping. Corona lemon, edge develops broad band of creamy apricot, flat, very fluted giving deeply lobed edge. Neck grows in water.
- BODILLY** 2W-Y
Perianth white, stained yellow at base. Segments flat, pointed, overlapping. Corona clear lemon yellow, develops thin cream edge. Neat.
- BOKHARA** 2Y-O
Perianth pale lemon yellow, creased and sometimes buckled. Segments flat, pointed, overlapping. Corona pale orange. General effect, small washed out and floppy.
- BOLTON** 7Y-Y
Perianth clear lemon yellow, very smooth, edges curl in. Segments flat, pointed, broad, overlapping. Corona deeper lemon yellow. Neck grows in water.
- BOMBAY** 2Y-YYR
Perianth mid lemon, creased. Segments flat, pointed, broad, well overlapping. Corona gold base shading to bright orange for half its length, fluted, lobed, bowl shape. Neck grows stiffly in water.
- BONAPARTE** 2Y-Y
Perianth greenish primrose yellow, crêpey texture, often creased. Segments flat, pointed, overlapping. Corona lemon yellow, fluted. Faded looking colour effect.
- BONFIRE** 3Y-O
Perianth creamy primrose, developing to cream, stained yellow at base. Segments slightly twisted, pointed, barely overlapping. Corona orange gold shading to bright orange red edge for 3/5 of its length, deeply fluted, crinkled.
- BONNEVILLE** 2W-W
Discarded - Short, coarse flower.
- BONYTHON** 1W-Y
Perianth creamy white, tinged yellow on back, creased. Segments flat, pointed, broad, overlapping. Corona clear sulphur yellow, rolled back edge.
- BORDER CHIEF** 2Y-O
Perianth bright lemon, slightly creased, crêpey. Segments flat, pointed, overlapping. Corona bright orange, paler at base, fluted, frilled, bowl shape. Neck grows stiffly in water.
- BORDER LEGEND** 2Y-O
Perianth canary yellow. Segments broad and overlapping with central ridge. Corona bowl shaped, flared, frilled, and orange in colour, slightly paler in centre. Neck grows badly in water.
- BOREAS** 2W
Discarded - Flowers below foliage, damages easily in wind.
- BOSWELL** 3W-YYR
Perianth creamy white stained yellow at base, creased, edges curl in. Segments flat, pointed, overlapping. Corona greenish lemon changing to band of bright orange red for 1/3 of its length, shallow bowl shape, fluted. Neck grows stiffly in water.
- BOSWIN** 1W-Y
Perianth creamy white, tinged green on back and base. Segments flat, pointed, overlapping. Corona clear lemon yellow, wide mouth, deeply frilled edge.
- BOTANY BAY** 2Y
Perianth light lemon yellow, creased, edges curled in. Segments flat, pointed, overlapping. Corona light orange, fluted, deep bowl shape. Neck grows badly in water, poor poise.
- BOVAGH** 2W-YYO
Perianth creamy white. Bottom segments ovoid, well rounded; top segments creased, notched, curled in at edges, rounded outline; overlapping at base. Corona sulphur yellow, pale orange, deeply lobed edge.
- BRABAZON** 1Y-Y
Perianth lemon, slightly twisted, slightly hooded. Segments broad, overlapping to 3/4 of length. Corona deep lemon, rolled back with deeply lobed rim. Corona appears large in relation to perianth, good, strong colour.
- BRAEMAR** 1Y-Y
Perianth light lemon yellow, crêpey texture. Segments flat, pointed, overlapping. Corona bright lemon yellow.

- BRAHMS** 2W-O
Perianth pale creamy white, smooth. Segments broad, edges inward curling, overlapping to half length. Corona flat, pale orange at centre becoming deeper at edge, crenate, edge gently frilled. Neck grows stiff in water.
- BRAIDWOOD** 2Y-Y
Perianth white. Segments broad and overlapping - double triangle layers obvious. Corona pale yellow, open mouthed and frilled. Neck grows badly in water.
- BRANDON** 1Y-Y
Perianth, greenish lemon yellow with green back. Segments twisted, pointed, barely overlapping. Corona deep clear lemon yellow, untidy deeply frilled edge, broad in proportion to perianth.
- BRANIEL** 2Y-O
Perianth opens primrose, develops to cream, stained yellow at base and stripes down the centre of each segment, crêpey texture. Segments flat, pointed, overlapping. Corona bright orange with slight gold edge, colour varying with age.
- BRAVADO** 2Y-O
Perianth sulphur with paler tips, creased, crêpey, circular outline. Segments flat, rounded, broad, well overlapping, edges curl in a little. Corona bright orange, fluted, frilled, shallow bowl shape. Neck grows in water.
- BRAVOURE** 1W-Y
Perianth milky white, stained yellow base. Segments rounded, overlapping, fairly smooth, some creasing, bottom segments very broad; double-triangle shape. Corona sharp lemon, cylindrical. Very good flower substance.
- BRAVURA** 3W-O
Perianth opens primrose, develops creamy white, stained yellow base, creased, sometimes notched and buckled, top 3 segments curl in, bottom 3 segments reflex. Segments flat, pointed, overlapping. Corona greenish gold eye changing to bright orange red, fluted, frilled, lobed, nearly flat. Neck grows badly in water.
- BREAD AND CHEESE** 2W-Y
Perianth white stained slightly yellow at base, creased. Segments flat, pointed, broad, overlapping. Corona deep bright lemon yellow ageing to buff primrose with paler edge.
- BREAGE** 2Y-Y
Discarded - Short.
- BREAK OF DAY** 2Y-R
No description available.
- BRIDAL CROWN** 4W-Y
Discarded - Short, flowers below foliage.
- BRIDGET HILL** 2Y-O
Perianth pale lemon, creased, one segment often buckled. Segments flat, pointed, overlapping. Corona gold base shading to bright orange for 1/2 its length, fluted. Neck grows in water.
- BRIGHT BEAUTY** 3W-O
Perianth creamy white, stained yellow at base. Segments flat, pointed, overlapping. Corona bright solid orange red, sometimes with white blotches on edge, fluted, untidy edge.
- BRIGHT CHARM** 3W-O
Perianth creamy white, a little creased, edges curl in. Segments flat, rounded, broad, well overlapping. Corona bright solid orange red, fluted, almost flat. Hangs head.
- BRIGHT CIRCLE** 3W-O
Perianth creamy white, creased, edges curl in. Segments flat, pointed, broad, overlapping. Corona pale orange shading to bright deep orange edge for 2/3 of its length, deeply fluted, almost flat.
- BRIGHT LAD** 2Y-O
Perianth canary yellow. Segments ovate, 1/2 overlapping and crêpey. Corona orange, funnel shaped and frilled. Neck grows badly in water.
- BRIGHT LASS** 2Y-YYR
Perianth canary yellow. Segments broad and overlapping with central ridge. Corona bowl shaped, frilled, mainly yellow with an orange band at rim. Neck grows badly in water.
- BRIGHTLING** 2Y-OOR
Perianth pale creamy primrose yellow, creased, fades with age. Segments flat, rounded, overlapping. Corona deep gold changing abruptly to bright deep orange for 1/2 its length, fluted, open bowl shape.
- BRILLIANCY** 3Y-YYO
Perianth pale sulphur yellow, spidery looking. Segments twisted, pointed, separated. Corona deep yellow shading to narrow bright red edge for 1/4 its length. Neck grows in water.

- BRILLIANT STAR** 11Y-OOY
Perianth creamy white. Segments broad, overlapping and reflexed. Corona split, creamy white with orange stripes. Neck grows badly in water.
- BRIMSTONE** 1Y-Y
Perianth pale creamy sulphur yellow. Segments flat, pointed, overlapping. Corona deep sulphur yellow, paling with age, leaving deeper lemon rim, flanged.
- BRITISH CHARM** 2Y-O
Perianth lemon yellow, paler at tips, creased, edges curl back. Segments flat, pointed, barely overlapping. Corona bright orange red, fluted.
- BROOKFIELD** 2W-W
Perianth white. Segments very broad and well overlapping, double triangle layers obvious. Corona wide mouthed and frilled, same colour as perianth.
- BROOKVILLE** 2W-Y
Perianth creamy white, creased, buckled, crêpey texture. Segments flat, pointed, overlapping. Corona greenish lemon base shading to pale orange edge for 1/2 its length, variable colour, deeply fluted, untidy lobed edge. Rough. Neck grows in water.
- BROUGHSHANE** 1W-W
Perianth creamy white, slightly creased. Segments flat, pointed, broad, well overlapping. Corona creamy white, open mouthed. Large but well balanced. Neck grows in water.
- BRUNSWICK** 2W-Y
Perianth white, sometimes one segment buckled. Segments flat, pointed, broad, overlapping. Corona deep primrose yellow, develops to pale primrose yellow with deeper edge, of almost trumpet proportions. Neat.
- BRUSSELS** 1W-W
Perianth white, creased, top segments inclined to incurve. Segments slightly twisted, pointed, overlapping. Corona creamy white, lightly frilled, rolled back edge, neat. Stiff.
- BRUTUS** 2Y
Perianth bright lemon yellow. Segments twisted, pointed, separated. Corona deeper lemon yellow with untidy deeply frilled edge. Neck grows in water.
- BRYHER** 3W-W
Perianth white. Segments very broadly ovate and well overlapping with central ridge. Corona cup shaped, filled with a green eye. Neck grows badly in water.
- BUCKLER** 2W-ORR
Perianth creamy white, stained yellow at base, pointed outline. Segments flat, reflexed, pointed, barely overlapping. Corona orange, shading to deep rich orange-red edge for 2/3 of its length, fluted.
- BULLA** 1W-Y
Perianth pale lemon white. Segments broad and overlapping. Corona open mouthed and frilled, yellow in colour. Neck grows in water.
- BULLION** 1Y-Y
Perianth pale lemon yellow, a little creased. Segments slightly twisted, pointed, overlapping. Corona bright lemon yellow, edge flanged.
- BUNCO** 2Y-YYR
Discarded - Small.
- BUNCRANA** 2W-O
Perianth white, stained slight yellow at base. Segments flat, rounded, well overlapping. Corona bright orange shading to gold base. Neck grows in water. Stiff and neat.
- BUNDORAN** 2W-P
Discarded - Poor colour, white blotches on corona edge, susceptible to basal rot.
- BURGEMEESTER GOUVERNEUR** 1Y-Y
Perianth soft sulphur yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona bright clear lemon yellow, flanged, neat.
- BURNING HEART** 11Y-OOW
Perianth opens primrose yellow, develops muddy cream, deeper at base, creased. Segments flat, pointed, overlapping. Corona divided into 6 long lobes of deep orange, edged all round with cream, untidy. Neck grows in water.
- BURRATOR** 3W-YYR
Perianth white. Segments broad and overlapping. Corona cup shaped, yellow with orange band around rim. Neck grows badly in water.

- BUSHMILLS** 3W-YYO
Perianth white, slightly incurling. Segments broad, ovate, overlapping at base. Corona lemon at base, orange at edge, small and flat, heavily ribbed giving frilled rim. Neck grows badly in water.
- BUSHTIT** 6Y-Y
Discarded - Short, small, cyclamineus hybrid.
- BUSTER** 2Y
Perianth canary yellow. Segments broad and overlapping with central ridge. Corona funnel shaped, frilled, yellow centre going orange towards rim. Neck grows badly in water.
- BUTTER AND EGGS** 4Y-Y
Perianth pale lemon, flat, pointed, barely overlapping, at least 6 rows complete, intermixed with light orange corona segments. Outer row inclined to reflex. Very full and heavy with weak neck. Neck grows badly in water.
- BUTTER KING** 1Y-Y
Perianth yellow. Segments broad and overlapping, with central ridge. Corona open mouthed and frilled. Same yellow as perianth. Neck grows in water.
- BUTTERCUP** 7Y-Y
Perianth deep clear lemon yellow, edges curl in a little, crêpey texture. Segments flat, pointed, overlapping. Corona deeper clear lemon yellow. Neck grows in water.
- BUTTERSCOTCH** 2Y-Y
Perianth sulphur, slightly creased. Segments long, would fit into a double triangle. Corona sulphur, slightly darker than Perianth, flanged, lightly frilled. Well balanced flower.
- BUZZARD** 2W
Perianth creamy white, creased, one segment buckled. Segments flat, pointed, broad, overlapping. Corona pale lemon yellow, some white blotches on edge, of trumpet proportion.
- C.J.BACKHOUSE** 2Y-O
Perianth pale lemon yellow, very crêpey texture. Segments twisted, pointed, separated. Corona bright orange, shading to orange-gold base, fluted, pinched in. Flower has thin spidery appearance.
- CAEDMON** 9W-YYR
Perianth pure white stained slight yellow at base, often creased. Segments flat, rounded, overlapping. Corona lemon yellow, changing to deep red edge for 1/3 of the length, with a moss green eye, almost flat, fluted. Style very long. Neck grows in water.
- CAERLEON** 2Y-O
Perianth soft sulphur yellow, crêpey texture, slightly creased. Segment flat, pointed, broad, overlapping. Corona light orange, fluted, bowl shape. Stiff.
- CAIRNDHU** 2W-GPP
Perianth white. Segments broadly round and 1/2 overlapping (shovel shaped), slightly creased with turned in edges. Corona moss green throat, pale salmon darkening towards edge. Edge lobed and evenly frilled.
- CAIRO** 2W-R
Perianth creamy white, notched, edges curl in. Segments flat, rounded, overlapping. Corona bright deep orange red, fluted, flat. Very stiff.
- CALABAR** 2Y-Y
Discarded - Poor colour, hooded.
- CALEDONIA** 2W-YYO
Perianth dirty creamy white stained yellow at base, creased. Segments flat, pointed, overlapping. Corona deep gold changing to bright orange red edge for 1/4 its length, almost flat, deeply fluted.
- CALIFORNIA** 2Y-Y
Perianth bright lemon yellow, a little creased. Segments flat, pointed and overlapping. Corona deeper golden yellow. Neat. Possibly synonymous with Pentewan (q.v). Neck grows badly in water.
- CALIFORNIA GOLD** 2Y-O
Perianth bright lemon yellow, creased, circular outline. Segments flat, rounded, well overlapping. Corona bright orange red throughout, fluted. Spoilt by excessively long neck which grows badly in water.
- CAMBER** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed and frilled - same yellow as perianth.
- CAMBRIA** 1Y-Y
Discarded - Short, ugly flower.
- CAMELLIA** 4Y-Y
Segments soft primrose yellow throughout, flat, pointed, overlapping, evenly spaced in the shape of a double camellia.

- CAMELOT** 2Y-Y
Perianth white. Segments broad with slight overlap. Inner whorl has edges curled inwards. Corona small cup with frilled edge, colour - light pink in centre with orange rim.
- CAMERONIAN** 1W-W
Discarded - Hangs head badly.
- CANARYBIRD** 8Y-GOO
Perianth pale sulphur yellow, often creased and buckled. Segments flat, rounded, overlapping. Corona bright orange gold with slight green eye, fluted.
- CANASTA** 11W-Y
Perianth white. Segments broad and overlapping. Corona split, disc shaped, laid back on perianth and frilled. Neck grows badly in water. Weak stems.
- CANDIDA** 4W-Y
Perianth creamy white. Double segments round, 1/2 overlapping, creased and curled in at edges. Segments loosely intermixed with corona. Corona frilled, short and cream coloured. Neck grows in water.
- CANDOUR** 2W-W
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona opens primrose, develops cream, of trumpet proportion.
- CANOPUS** 1W-W
Perianth white, creased, crêpey texture. Segments flat, pointed, broad, well overlapping. Corona creamy white, hangs head badly.
- CANTABILE** 9W-GYR
Perianth pure white, smooth but sometimes creased, reflexed. Segments flat, rounded, well overlapping. Corona greenish lemon changing to orange red edge for 1/6 of its length, with a large moss green eye, flat, fluted. Style very short.
- CANTATRICE** 1W-W
Perianth pure white, smooth. Segments flat, pointed, broad, overlapping. Corona white, slender. Faultless flowers.
- CAPE HORN** 1W-Y
Discarded - Hangs head.
- CAPE KENNEDY** 11W-O
Perianth white. Segments very broad and overlapping. Corona split, orange and white in colour. Neck grows badly in water.
- CAPISCO** 3W-GYO
Perianth poeticus white, yellow at base. Segments cupped, ovate to rounded, well overlapping. Corona greenish yellow at base, shading to lemon, orange rim, very short, straight sided, fluted edge. Neck grows in water.
- CAPPAROE** 3W-R
Perianth white. Segments broad and overlapping, edges turned in. Perianth disc shaped, frilled and orange/red in colour. Neck grows badly in water.
- CARACAS** 2O-O
No description available.
- CARAMEL** 9W-O
Perianth white. Segments broad and overlapping, crêpey with central streak. Corona open mouthed, frilled and pale yellow. Neck grows in water.
- CARAVELLE** 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona flared mouth and frilled. Same colour as perianth. Neck grows in water.
- CARBINEER** 2Y-O
Perianth bright deep lemon, paler at tips, creased. Segments flat, pointed, broad, overlapping. Corona gold shading to bright orange red for 1/2 the length, fluted. Neck grows in water. Stiff and sturdy.
- CARDIGAN** 2W-Y
Perianth creamy white, stained yellow base, sometimes notched, slightly creased, edges curl in. Corona clear lemon, lightly frilled, almost straight sided. Neck grows in water.
- CAREFREE** 1W
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, well overlapping. Corona creamy white with deeper edge, frilled, trumpet proportion.
- CAREYSVILLE** 2W-Y
Perianth white. Segments broad, overlapping and very ridged. Corona open mouthed and frilled, lemon yellow in colour. Neck grows badly in water.

CARGAN 2Y-Y
Perianth bright lemon yellow, smooth. Segments flat, pointed, broad, well overlapping, shovel shape. Corona deeper lemon yellow, of trumpet proportion. Very stiff and neat.

CARLTON 2Y-Y
Perianth soft sulphur, creased. Segments slightly twisted, pointed, broad and overlapping. Corona deep lemon, deeply frilled, crenate open mouth. Neck grows badly in water.

CARMINOWE 3W-R
Perianth pure white, stained yellow at base, a little creased, edges curl in. Segments flat, pointed, overlapping. Corona solid deep red, deeply fluted. Neck grows in water.

CARMOA 2W-YYP
Perianth creamy white, edges curled in. Segments flat, pointed, overlapping. Corona opens deep primrose, developing to pale cream tinted peach with pale peach edge. Neck grows in water.

CARNALEA 2W-WWY
Perianth creamy white, a little creased, edges curl in a little. Segments flat, pointed, overlapping. Corona pale greenish primrose shading to greenish lemon edge, overlaid with light orange for 1/4 of its length, fluted, shallow bowl shape.

CARNATION 4Y-O
Perianth segments cream, flat, pointed, separated, 2 rows. Corona segments orange, intermixed. Rather full flower. Neck grows badly in water.

CARNBEG 2Y-OOR
Perianth canary yellow. Segments broad and overlapping. Corona open mouthed and frilled. Colour same as perianth getting deeper towards rim. Neck grows badly in water.

CARNLOUGH 2W-WWP
Perianth white, slightly creased. Segments flat, pointed, broad, overlapping. Corona pale creamy primrose with narrow pink edge, develops to cream throughout, of trumpet proportion.

CARNMONEY 2W-W
Perianth creamy white, a little creased. Segments flat, pointed, overlapping. Corona opens deep cream, developing to pale cream with narrow edge of white, slightly fluted, deep bowl shape. Neck grows in water.

CARNMOON 3W-WWY
Perianth creamy white, slightly creased, edges curl in. Segments flat, rounded, overlapping. Corona grey green eye changing to cream with narrow rim of primrose developing to buff, deeply fluted, frilled, lobed. Neck grows stiffly in water.

CARNSORE 3W-GR
Perianth opens pale creamy primrose, develops to creamy white. Segments flat, pointed, overlapping. Corona rich orange red shading to small greenish gold base, with greenish eye, fluted. Neck grows in water.

CARO NOME 2W-WPP
Perianth white. Segments broad and overlapping with central ridge. Corona bowl shaped and frilled, starting off white, with a pink rim. Neck grows badly in water.

CAROLINE FOX 1W-Y
Perianth lemony/white. Segments broad and overlapping with central ridge. Corona open mouthed, frilled and canary yellow in colour. Neck grows badly in water.

CARPATICA 3W-O
Perianth creamy white, stained yellow at base, edges curl in a little. Segments flat, rounded, overlapping. Corona bright orange red throughout, fluted, almost flat, stiff. Neck grows badly in water.

CARRICK 2W-P
Perianth white. Segments broad and overlapping. Corona open mouthed and frilled, yellow with pink rim.

CARRICKBEG 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona funnel shaped, frilled and same colour as perianth.

CARRICKMANNON 2W-GPP
No description available.

CARRIGART 3Y-O
Perianth sulphur yellow, a little creased. Segments flat, rounded, overlapping. Corona orange red, shading paler at base, fluted, bowl shape.

CASA D'ORO 1Y-Y
Discarded - Short.

CASABIANCA 2W-Y
Perianth white. Segments broad and overlapping, with ridges. Corona open mouthed, frilled and canary yellow.

- CASABLANCA** 2W-Y
Perianth creamy white, a little creased, edges curl in. Segments flat, pointed, broad, well overlapping. Corona light lemon yellow throughout, almost straight sided. Neat and stiff. Neck grows in water.
- CASSANDRA** 9W-YYR
Perianth white, edges often curled back, thin, floppy. Segments flat, pointed, separated. Corona slight greenish gold eye, shading to gold, with narrow rim of bright orange red, fluted, flat. Style longer than stamens. Neck grows badly in water.
- CASSATA** 11W-W
Perianth creamy white, pointed outline. Segments flat, pointed outline, barely overlapping. Corona cream, split into 6 segments, frilled edges. Neck grows in water.
- CASTELLA** 2W-W
Perianth creamy white, creased, pointed outline. Segments flat, pointed, overlapping. Corona opens primrose yellow, develops to pale cream, almost flat. Rough looking flower.
- CASTLE COOLE** 2W-WWO
Discarded - Hangs head.
- CASTLE KENNEDY** 2W-WWY
Perianth creamy white, crêpey texture, edges curled in. Segments flat, pointed, broad, overlapping. Corona pale cream shading to deeper cream edge for 1/3 its length, grey green eye, fluted, shallow bowl shape. Neck grows in water.
- CASTLE OF MEY** 2W-W
Perianth white. Segments broad and overlapping with central ridge. Corona open mouthed, frilled and same white as perianth.
- CASTLEBAR** 1Y-Y
Perianth deep clear lemon yellow, edges curl in a little. Segments flat, pointed, overlapping. Corona deep intense lemon yellow. Well proportioned, stiff and sturdy.
- CASTLEREAGH** 1Y-Y
Perianth soft lemon yellow, edges curl in slightly. Segments flat, pointed, overlapping. Corona deeper lemon yellow.
- CASTLEROCK** 2Y-R
Perianth mid lemon with paler tips, creased, very crêpey, often caught in the corona, circular outline. Segments flat, rounded, broad, well overlapping, edges curl in a little. Corona orange base shading to orange red, fluted, lobed, bowl shape. Neck grows stiffly in water.
- CASTLEWELLAN** 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona open mouthed and frilled, same yellow as perianth.
- CATHAY** 2Y-R
Perianth strong yellow. Segments rounded, overlapping. Corona orangey-red, expanded, frilled edge. Very good flower substance. Neck grows in water.
- CATRIONA** 2W-YYW
Perianth white. Segments very broadly ovate and overlapping. Corona bowl shaped and frilled. Colour pale lemon getting deeper at rim. Neck grows in water.
- CAVALIER** 3W-YOO
Perianth white. Segments very broadly ovate and well overlapping. Corona small cup, frilled and canary yellow.
- CAVALIERO** 2Y-YOO
Perianth bright clear lemon yellow, perianth often catches in corona, star shaped. Segments flat, pointed overlapping. Corona bright orange red shading to tangerine base, almost straight sided, attractive glow. Very stiff neck.
- CAVODA** 1W-GPP
Perianth creamy white, stained green back base. Segments rounded, overlapping, slightly creased, some notched, curled in at edges. Corona shell pink, paler at edge, expanded, lobed.
- CELEBRITY** 1W-Y
Perianth creamy white tinged yellow at base, crêpey texture, creased. Segments twisted, pointed, overlapping. Corona deep primrose yellow.
- CELESTIAL** 5Y-Y
Discarded - Small, short, triandrus hybrid.
- CERVANTES** 1Y-Y
Perianth pale creamy sulphur yellow, deeper at base, crêpey texture. Segments twisted, pointed, separated. Corona pale lemon yellow, paler at base, slender.

- CEYLON
Perianth bright lemon, smooth, pointed outline. Segments flat, pointed, overlapping. Corona orange gold shading quickly to deep orange, slightly fluted, almost straight sided. Stiff. 2Y-O
- CHAMPAGNE
Perianth white, creased, notched, pointed outline. Segments twisted, pointed, overlapping. Corona pale flesh pink base shading through deeper colour to apricot rim. Ages to paler colour. Ugly. 2W-P
- CHANTERELLE
Discarded - Short, small, flowers below foliage. 11Y-Y
- CHANTILLY
Discarded - Rough, flowers below foliage, susceptible to basal rot. 2W-O
- CHARIOTEER
Perianth yellow. Segments broad and overlapping. Corona funnel shaped and frilled. Corona same colour as perianth. 1Y-Y
- CHARIS
Perianth creamy white, a little creased, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona creamy primrose, with narrow rim of peach for 1/4 of its length, some white blotches on edge, fluted. Neck grows badly in water. 2W-P
- CHARITY MAY
Perianth sulphur yellow, slightly green on back base, well reflexed. Segments flat, pointed, broad, overlapping. Corona pale lemon yellow, slightly waisted. Neat, well proportioned. 6Y-Y
- CHARLES
Perianth opens pale primrose, quickly turning to cream stained lemon at base. Segments flat, pointed, overlapping. Corona orange red shading to greenish gold in the centre. Stiff and sturdy. 3W-YOO
- CHARTER
Perianth cream with pale yellow edges, slightly twisted. Segments rounded, overlapping only slightly. Corona primrose, straight sided, with slightly frilled edge. Neck grows badly in water. Weak stems. 2Y-W
- CHATSWORTH
Perianth creamy white, a little green on back. Segments flat, pointed, broad, overlapping. Corona clear lemon yellow. 1W-Y
- CHEERFULNESS
Perianth creamy white, complete. Segments flat, rounded, overlapping. Corona yellow with thin orange rim, complete, filled in with rounded, twisted cream segments, intermixed with twisted corona and stamen segments. 2-3 florets per stem. Neck grows badly in water. 4W-Y
- CHEERIO
Perianth pale primrose yellow with paler tips, creased but very stiff, hooded. Segments flat, rounded, overlapping. Corona bright orange shading to orange gold base. 2Y-O
- CHELSEA CHINA
Perianth creamy white, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona intense clear bright pink throughout, a little fluted, deep bowl shape. Neck grows in water. 2W-GPP
- CHELSEA DERBY
Perianth white. Segments broad and overlapping, edges turned in. Corona open mouthed, frilled, yellow in centre, with a pink rim. Neck grows badly in water. 2W-P
- CHEMAWA
Perianth yellow. Segments very broadly ovate, overlapping and crêpey. Corona cup shaped, frilled, orange with a yellow rim. Neck grows badly in water. 2Y-OOY
- CHERIE
Perianth creamy white stained yellow at base, smooth. Segments flat, rounded, overlapping. Corona opens pale buff cream with deeper edge, develops to soft peach pink. Neck grows in water. 7W-P
- CHEVALIER
No description available. 1Y-Y
- CHIEF JUSTICE
Perianth creamy white, slightly creased. Segments flat, pointed, broad, well overlapping. Corona pale creamy primrose. Hangs head badly. 1W-Y
- CHIFFON
Perianth white. Segments broad and overlapping and slightly notched. Corona bowl shaped, frilled, starting pale pink in centre getting darker towards rim. Neck grows in water. 2W-P
- CHILLAGOE
Perianth white, slightly twisted. Segments would fit into a double triangle, broad and overlapping at base. Corona white, flanged, slightly frilled at rim. Bold flower. Neck grows in water. Weak stems. 2W-W

CHILLION No description available.	1Y-Y
CHILTERN Perianth creamy white. Segments ovate and overlapping. Corona open mouthed, frilled and lemon yellow in colour. Neck grows in water.	1W-Y
CHINA PINK Discarded - Small, short.	2W-P
CHINESE WHITE Perianth creamy white, creased, crêpey, edges curl in, circular outline. Segments flat, rounded, broad, overlapping. Corona grey eye, changing to creamy white, deeply fluted, frilled, lobed, shallow bowl shape. Hangs head when young. Neck grows badly in water.	3W-W
CHINITA Perianth primrose yellow, develops cream stained yellow at base with age, inclined to reflex, often creased and buckled. Segments flat, rounded, overlapping. Corona deep gold with thin red edge, fluted. Neck grows in water.	8Y-YYR
CHINOOK Perianth creamy white, slightly twisted, creased. Segments broad, ovate, overlapping to 3/4 of length. Corona pale creamy orange at base becoming cream at rim, open mouthed, crenate, rim lobed. Neck grows in water.	2W-Y
CHITRA No description available.	2Y-R
CHMOY Perianth white. Segments broad and overlapping. Corona flared, frilled cylinder, yellow in colour. Neck grows badly in water.	2W-Y
CHOIR BOY Perianth white, with yellow stain in centre. Segments broad and overlapping, slightly notched. Corona cup shaped, frilled and orange in colour. Neck grows badly in water.	2W-O
CHUNGKING Perianth deep lemon yellow, slightly crêpey texture and creased, edges curl in, circular flower. Segments flat, rounded, well overlapping. Corona bright orange red, fluted. Neck grows in water.	3Y-O
CHYNANCE Perianth white. Segments broad overlapping, and ridged. Corona cup shaped, frilled and canary yellow in colour. Neck grows badly in water.	2W
CIBOLA Perianth deep lemon, flower semi-upward pointing, slightly reflexed. Segments edges backward curled. Corona deep lemon, short and flanged, deeply frilled. Neck grows badly in water.	2Y-Y
CICELY Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona opens pale lemon, develops creamy primrose a little paler at base, straight sided. Neck grows badly in water.	2W-W
CIMAROSA Discarded - Short, susceptible to basal rot.	1W-Y
CINEL Perianth canary yellow. Segments broadly ovate and overlapping. Corona bowl shaped, frilled and deep orange. Neck grows badly in water.	2Y-R
CINZANO Discarded - Floppy, poor poise.	2W-O
CITRIX Perianth greenish cream, some green on back, creased, crêpey texture. Segments twisted, pointed, overlapping. Corona pale creamy lemon, developing cream with lemon rim, broad mouth. Neck grows in water.	1Y-Y
CLACKAMAS Discarded - Hangs head.	2W-YYO
CLACKRATTLE Perianth deep lemon yellow with paler tips, slightly creased, crêpey texture. Segments flat, rounded, overlapping. Corona bright coppery red, bowl shaped, fluted.	2Y-O
CLAMOR Perianth creamy white stained yellow at base, creased. Segments flat, rounded, overlapping. Corona lemon-gold changing to band at edge of bright red for 1/3 its length. Stiff and sturdy.	2W-YYR

- CLAVA** 2W-Y
Perianth cream, stained yellow at base, edges curl in, making pointed outline. Segments flat, rounded, overlapping. Corona lemon, shading to pale orange edge for 1/3 its length with slight paler edge.
- CLEENA** 2W-OOY
Perianth opens greenish primrose, develops creamy white, slightly creased. Segments flat, pointed, overlapping. Corona greenish lemon developing apricot, overlaid apricot orange for 1/3 its length with narrow edge of pale apricot, variable colour, occasional white blotches on edge, deep bowl shape.
- CLEMENTINE** 2W-Y
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and very pale yellow in colour. Neck grows in water.
- CLOCKFACE** 3W-YYO
Perianth creamy white, stained yellow base, creased, edges curl in a little. Segments flat, pointed, broad, well overlapping. Corona moss green eye, changing to gold base 2/3, with band of orange red, fluted, frilled, lobed, almost flat. Neck grows badly though stiffly in water.
- CLOGHEEN** 3W-Y
Perianth white. Segments broad and overlapping, edges turned in. Corona cup shaped and frilled. White in centre with a yellow rim. Neck grows in water.
- CLONCARRIG** 1Y-Y
Perianth bright lemon, slightly creased. Segments flat, pointed, broad, well overlapping, shovel shape. Corona deep lemon, lobed, flanged, broad mouth. Stiff.
- CLONEEN** 2W-W
Perianth white. Segments broad and overlapping with central ridge. Corona cup shaped, frilled and white in colour. Neck grows in water.
- CLONMORE** 1Y-Y
Perianth medium lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona deeper lemon yellow. Stiff and neat. Neck grows a little in water.
- CLOUDCAP** 2W-WWP
Perianth white. Segments broad and overlapping. Corona bowl shaped, frilled, white with pink rim. Neck grows in water.
- COLORANGE** 11Y-O
Perianth canary yellow. Segments very broad, ovate, overlapping with slightly paler central ridge. Corona split and intermixed with perianth. Corona segments deep yellow/pale orange, and frilled. Double layers obvious, with segment of corona attached to each perianth segment. Neck grows badly in water.
- COLORATURA** 3W-WWO
Perianth pale cream, slightly hooded, smooth. Segments edges incurling, would fit into double triangle due to in rolling of edges. Corona cream with orange rim, large, flat with frilled rim. Neck grows badly in water.
- COLUMBINE** 3W-WWO
Perianth poeticus white, edges curl in, reflexed, pointed outline. Segments flat, pointed, well overlapping. Corona greyish green, changing to apricot orange for 1/2 its length, slightly fluted, flat. Hangs head badly. Neck grows badly but stiffly in water.
- COMMODORE** 2W-O
Discarded - Hangs head, white blotches on corona edge. Susceptible to basal rot.
- COMPRESSUS** 8W-Y
Perianth white, reflexed. Segments flat, rounded, overlapping. Corona pale lemon yellow, goblet shape. Style half way between short and long stamens.
- COMPTON MACKENZIE** 2W-P
Discarded - Very short vase life, corona edge dies first.
- CONBEG** 2W-R
Discarded - Small, flowers below foliage, rough flower.
- CONFUOCO** 2Y-R
Discarded - Flowers below foliage, very untidy corona.
- CONFUSION** 2W-P
No description available.
- CONGO** 2Y-O
Perianth canary yellow. Segments broad and overlapping with central ridge and edges turned in. Corona bowl shaped, frilled and orange in colour. Neck grows badly in water.
- CONSPICUUS** 3Y-YYO
Perianth pale clear lemon yellow. Segments slightly twisted, pointed, barely overlapping. Corona pale gold with bright orange red edge for 1/4 of its length, small, fluted. Neck grows in water.

- CONTENT** 1W-WWY
Perianth opens pale creamy sulphur, develops to creamy white, slightly creased. Segments flat, pointed, broad, well over-lapping. Corona pale lemon, growing paler with age, long and slender.
- CONTRAPUNT** 2Y-O
Perianth canary yellow. Segments ovate and overlapping, slightly crêpey. Corona cup shaped, frilled and orange in colour. Neck grows in water.
- CONVAIR** 1Y-Y
Perianth yellow. Segments ovate, overlapping with paler central ridge. Corona open mouthed, lobed and same colour as perianth. Neck grows in water.
- COOL CONTRAST** 1W-Y
No description available.
- COOLATTIN** 2W-Y
Perianth white. Segments ovate, overlapping, and slightly notched. Corona cup shaped, frilled, pale yellow with slightly darker band at rim. Neck grows in water.
- COPPELIA** 2W-P
Perianth white. Segments broad and overlapping. Corona cup shaped, crenate, and salmon pink in colour. Neck grows in water.
- COPPER BOWL** 2Y-O
Perianth primrose yellow, deeper at base, creased. Segments flat, pointed, broad, overlapping. Corona bright orange, fluted.
- COPPERSMITH** 2Y-O
Perianth gold overlaid slight orange, creased, crêpey, pointed outline. Segments flat, pointed, overlapping. Corona clear orange throughout, fluted, frilled, lobed, bowl shape. Neck grown stiffly in water.
- CORAL DAWN** 2W-Y
Discarded - Short, floppy flower.
- CORAL ISLAND** 2W-GPP
Perianth creamy white, slightly creased, edges curl in a little at tips, pointed outline. Segments flat, pointed, overlapping. Corona grey green eye, changing to primrose overlaid pale apricot, shading paler towards edge, with cream rim, variable colour, fluted, lightly frilled, bowl shape. Neck grows in water.
- CORBY** 2W-W
Perianth creamy white, a little creased. Segments flat, pointed, overlapping. Corona cream throughout, fluted, deep bowl shape. Hangs head badly.
- CORDIAL** 2W-P
Perianth white. Segments ovate and overlapping. Corona funnel shaped, frilled and pink in colour. Neck grows in water.
- CORINTHIAN** 1Y-Y
Perianth sulphur yellow, green at back. Segments twisted, pointed, overlapping. Corona bright deep lemon yellow, open mouthed, flanged.
- CORNCRAKE** 3W-O
Perianth poeticus white, stained yellow base, slightly creased, circular outline. Segments flat, rounded, well overlapping. Corona moss green eye changing to deep gold base 1/2 changing to band of deep orange red, fluted, frilled, deeply lobed, nearly flat. Neck grows badly but stiffly in water.
- CORNET** 6Y-Y
Perianth clear lemon. Segments pointed, overlapping at base, creased, somewhat reflexed. Corona clear lemon, straight, flared crenate edge.
- CORNISH FIRE** 2Y-O
Perianth pale primrose yellow, one segment often buckled, pointed outline. Segments flat, pointed, overlapping. Corona burnt orange, fluted. Neck grows in water.
- COROFIN** 3W-YR
Perianth opens cream, quickly develops white stained yellow base, slightly creased, circular outline. Segments flat, rounded, broad, well overlapping. Corona bright greenish lemon base 2/3, changing to broad band of bright orange red, deeply fluted, frilled, lobed, nearly flat. Neck grows stiffly in water.
- CORONACH** 3W-R
Perianth poeticus white, stained yellow base, creased, edges curl in, top segments incurve a little. Segments flat, rounded, overlapping. Corona gold green eye, changing to gold and shading to deep orange red at edge, deeply fluted, frilled, lobed, nearly flat. Neck grows badly in water.
- CORONADO** 2Y-P
Perianth white. Segments ovate slightly overlapping and twisted. Corona open mouthed, frilled, yellow and getting deeper at rim.

- CORREGIO** 2W-O
Perianth opens primrose yellow, develops cream stained yellow at base, notched, creased, edges curl in. Segments twisted, pointed, barely overlapping. Corona gold base shading to orange for 1/2 its length, fluted, bowl shape. Neck grows badly in water, poor poise.
- CORRYMEELA** 2W-Y
Discarded - Poor colour.
- CORTINA** 1W-W
Perianth white, creased, edges curl in. Segments flat, pointed, overlapping. Corona cream, open mouthed.
- COTOPAXI** 2Y-O
Perianth lemon yellow, crêpey texture. Segments flat, rounded, well overlapping. Corona light burnt orange, fluted. Very stiff and sturdy.
- COTTERTON** 2W-W
Discarded - Short, hangs head, flowers below foliage, very susceptible to basal rot.
- COTTESMORE** 2Y-YOO
Perianth pale primrose yellow deeper at base. Segments flat, pointed, overlapping. Corona light gold, suffused orange, paler at base, fluted.
- COUNSELLOR** 1Y-Y
Perianth clear lemon yellow, slightly creased. Segments flat, pointed, overlapping. Corona deeper lemon yellow.
- COUNTESS OF STAIR** 1W-Y
Perianth creamy white, a little creased, one segment buckled. Segments flat, pointed, overlapping. Corona cream, rather untidy edge.
- COURAGE** 2W-W
Discarded - Short, flowers below foliage, susceptible to basal rot.
- COURT MARTIAL** 2Y-O
Perianth deep lemon, slightly creased, sometimes notched. Segments flat, pointed, broad, well overlapping. Corona bright burnt orange, slightly fluted, slightly frilled, almost straight sided. Neck grows stiffly in water.
- COVE** 2Y-Y
Perianth primrose, crease down centre to point, very twisted. Segments narrow, overlapping only at base. Corona flat to open mouthed, deep lemon, lightly frilled at edge. Neck grows badly in water.
- COVENT GARDEN** 1Y-Y
Perianth clear lemon yellow. Segments flat, pointed, overlapping. Corona deep golden lemon, flanged.
- COVERACK GLORY** 2Y-Y
Perianth clear light lemon with paler stripe down the centre of each segment, creased, crêpey texture. Segments nearly flat, pointed, broad, overlapping. Corona deep lemon yellow. Neck grows in water.
- COVERACK PERFECTION** 2W-YYO
Perianth creamy white, creased, edges curl in, star shaped. Segments flat, pointed, overlapping. Corona apricot shading through paler colour to greenish lemon base, fairly flat, fluted.
- COVERACK PRIDE** 1Y-Y
Perianth lemon, creased, pointed outline. Segments slightly twisted pointed, overlapping. Corona deeper lemon, crenate, flanged, broad mouth. Neck grows in water.
- CRAGFORD** 8W-O
Perianth creamy white, stained yellow at base, creased and edges folded in. Segments flat, rounded, overlapping. Corona deep red with yellowish green eye, long style.
- CRAIGYWARREN** 2Y-R
Perianth deep bright lemon yellow, paler at tips, smooth. Segments flat, pointed, broad, well overlapping. Corona solid deep red throughout, fluted, bowl shape. Stiff and sturdy.
- CRASKIN** 1Y-Y
Perianth deep clear yellow. Segments slightly twisted, pointed, overlapping, star shaped. Corona deep clear golden yellow, flanged.
- CREAM CUP** 2W-Y
Discarded - Short, poor poise.
- CREAMED HONEY** 1Y-W
Perianth white. Segments broad, overlapping and slightly twisted. Corona open mouthed, frilled and white in colour.
- CREBILLY** 2W-WWY
Perianth creamy white, creased, edges curl in giving star shape appearance. Segments flat, pointed, overlapping. Corona cream with narrow edge of creamy buff, with some white blotches on edge, fluted, shallow bowl shape. Washy colour, poor poise. Neck grows badly in water.

- CRENVER** 3W-YYO
Perianth pure white stained yellow at base, one segment sometimes creased. Segments flat, rounded, well overlapping. Corona greenish lemon with bright orange red edge for 1/3 its length and greenish eye, deeply fluted. Stiff and sturdy.
- CREPELLO** 3W-GWY
Perianth white. Segments rounded and overlapping with a central ridge and edges turning in. Also looks crêpey. Corona cup shaped, frilled, with green eye, segments white with a yellow rim. Neck grows badly in water.
- CRESCENDO** 2Y-YYO
Perianth light lemon yellow, a little creased, very rounded with a nice sheen. Segments flat, rounded, well overlapping. Corona gold shading to bright orange red edge for 2/3 its length, straight sided. Neck grows in water. Neat and stiff.
- CRESCENT** 2Y-R
Perianth bright yellow, crêpey texture, pointed outline. Segments flat, pointed, overlapping, corona bright red, fluted.
- CRINOLINE** 2Y
Discarded - Flowers below foliage.
- CRITERION** 2W-OOY
Perianth creamy white, creased, edges curl in. Segments flat, pointed, overlapping. Corona greenish gold changing to bright orange for 1/4 its length, fluted, shallow bowl shape. Neck grows a little in water.
- CROCK OF GOLD** 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona open mouthed, frilled and same colour as perianth. Neck grows in water.
- CROCUS** 2Y-Y
Perianth deep bright golden yellow, very smooth. Segments flat, pointed, broad, well overlapping. Corona deeper rich golden yellow, of trumpet proportion. Stiff and sturdy.
- CROESUS** 2Y-YYO
Perianth pale cream stained slight yellow at base, fades with age, slightly reflexed. Segments flat, pointed, overlapping. Corona gold shading to burnt orange edge for 1/3 of its length.
- CROMARTY** 1Y-Y
Perianth bright clear lemon yellow, very smooth, stiff, double triangle. Segments shovel shaped, flat, broad, well overlapping. Corona deeper lemon yellow, flanged. Well proportioned.
- CROWN DERBY** 3W-YYO
Perianth creamy white, stained yellow at base, creased, edges curl in, often buckled. Segments flat, pointed, broad, overlapping. Corona deep gold shading to bright orange edge for 1/2 its length, fluted, bowl shape. Neck grows in water.
- CRUMLIN** 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona open mouthed, lobed and same colour as perianth.
- CRUSADER** 2W-O
Perianth white. Segments very broadly ovate and overlapping. Corona bowl shaped, frilled and orange in colour. Neck grows in water.
- CRYSTAL QUEEN** 2W-W
Perianth creamy white, stained slight yellow at base, slightly creased. Segments twisted, pointed, overlapping. Corona bright sulphur yellow, with paler edge.
- CUSHENDALL** 3W-GWW
Perianth poeticus white, sometimes a little creased. Segments flat, rounded, well overlapping. Corona bright green centre with pale creamy buff edge for 1/2 of its length, almost flat. Stiff and sturdy.
- CUTTY SARK** 7Y-Y
Perianth pale sulphur yellow, developing to creamy sulphur, often creased down centre of segments, buckled. Segments flat, pointed, overlapping. Corona sulphur yellow, neat. Neck grows in water.
- CYCLADES** 6Y-Y
Discarded - Small, short, cyclamineus hybrid.
- CYCLONE** 6Y-Y
Discarded - Short, cyclamineus hybrid.
- CYCLOPE** 1Y-Y
Perianth clear lemon. Segments overlapping at base, pointed, outer segments broader, creased, curled in at edges. Corona rich gold, very broad, flared mouth, coarse ragged edge. Large, firm, flower, short stem. Very good flower substance.

- CYNOSURE** 2W-O
Perianth cream stained yellow at base, edges curled back, star shaped. Segments flat, pointed, separated. Corona deep lemon shading to orange edge, pinched in.
- DACTYL** 9W-GYR
Perianth pure white sometimes creased, curled in a little at edges. Segments flat, pointed, broad, well overlapping. Corona greenish lemon changing to deep orange red edge for 1/6 its length, with yellowish green eye. Style a little longer than longest stamens.
- DAISY SCHAFFER** 2W-WYY
Perianth white, slightly creased. Segments flat, pointed, broad, overlapping. Corona pale clear lemon yellow, paler at base, of trumpet proportions.
- DALBOYNE** 2Y-O
Perianth lemon yellow, slightly creased. Segments flat, pointed, overlapping. Corona bright orange, a little paler at base, straight sided. Neat and stiff.
- DALLAS** 3W-GWW
Perianth poeticus white, smooth, circular outline. Segments flat, rounded, broad, overlapping. Corona bright green ½ changing to poeticus white edge, fluted, frilled, almost flat. Neck grows stiffly in water.
- DAMASCUS** 3W-R
Perianth creamy white, slightly creased, edges curled in. Segments flat, pointed, overlapping. Corona deep burnt orange red, fluted, almost flat.
- DAMSON** 2W-O
Perianth opens primrose yellow, quickly changing to dirty cream stained deep yellow at base, crêpey texture. Segments slightly twisted, pointed, broad, overlapping. Corona deep bright orange red. Neck grows in water.
- DANAE** 2Y-O
Perianth rich warm gold, fairly smooth. Segments long, oval, rounded, partly overlapping at base. Corona bright mid orange, wavy, scalloped edge, irregularly lobed. Very good flower substance.
- DANCING PARTNER** 2W-P
Perianth white. Segments broad and overlapping - with edges turned in. Corona flared frilled funnel and pink in colour. Neck grows in water.
- DAPHNE** 4W-W
Perianth segments poeticus white in regular whorls through to centre where they are more pointed. There can be considerable variation from season to season when a proportion of flowers will be only semi-double or even single and show all or part of the poeticus corona from which it sported. Must be almost fully developed before picking as it will not open from bud. Neck grows badly in water.
- DARLINGTON** 2Y-O
Perianth yellow. Segments broad and overlapping. Corona cup shaped, frilled and orange in colour. Neck grows in water.
- DAVID SPRY** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and same colour as perianth. Neck grows in water.
- DAVIOT** 2W-OOY
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona apricot orange, a little paler at base, with narrow band of primrose, straight sided, fluted, almost straight edge. Neck grows stiffly in water.
- DAWLEY** 2Y-Y
Perianth lemon yellow. Segments broad and overlapping - with whitish tips. Corona open mouthed, frilled and same colour as perianth. Neck grows badly in water.
- DAWN** 5W-Y
Perianth creamy white, edges curl in, reflexed, star shaped. Segments flat, pointed, barely overlapping. Corona pale greenish lemon with narrow edge of deeper greenish lemon, almost flat, fluted. Very long protruding style. Neck grows very badly in water.
- DAWSON CITY** 1Y-Y
Perianth soft sulphur yellow, crêpey, slightly hooded. Segments pointed, broad, overlapping. Corona bright lemon, very slightly frilled, edge neatly rolled back.
- DAYDREAM** 2Y-W
Perianth primrose, paler at base, edges incurled. Segments broad, spade shaped, overlapping at base. Corona pale primrose, flanged, rim deeply lobed. Neat flower with attractive colour.
- DEAN** 2W-YOO
Perianth creamy white, edges curl in slightly, very solid. Segments flat, pointed, broad, well overlapping. Corona orange shading to lemon base.

- DEANNA DURBIN 2W-O
Perianth creamy white, stained yellow at base, one segment often creased. Segments flat, pointed, well overlapping. Corona deep apricot orange. Neat. Neck grows in water.
- DEBUTANTE 2W-P
Perianth white. Segments broad and overlapping and slightly notched. Corona cup shaped, frilled and pink in colour. Neck grows badly in water.
- DECENCY 1Y-Y
Perianth deep primrose yellow, some green on back, slightly creased. Segments flat, pointed, broad, overlapping. Corona deep clear yellow, almost straight, flanged, neat edge.
- DECOR 2Y-YOO
Perianth sulphur yellow, very creased and rough, segments often caught in corona. Segments flat, pointed, broad, overlapping. Corona pale orange shading to gold base, deeply frilled. Unbalanced and ugly.
- DEDERANG 1W-Y
Perianth white. Segments broad and overlapping with a central ridge. Corona open mouthed, frilled and yellow in colour.
- DEERFIN 2Y-Y
Perianth pale creamy yellow. Segments broad and overlapping with edges turned in. Corona open mouthed, frilled and same colour as perianth. Neck grows in water.
- DEFIANCE 2Y-O
Perianth yellow. Segments broad and overlapping. Corona cup shaped, frilled and orange in colour. Neck grows badly in water.
- DEIRDRE 3W
Perianth creamy white, smooth, top segments incurve. Segments flat, pointed, well overlapping. Corona large moss green eye, changing to cream with a narrow band of apricot orange for 1/5 of its length, frilled, deeply lobed. Hangs head when young. Neck grows badly in water.
- DELIBES 2Y-O
Perianth sulphur yellow, creased, one or two segments sometimes buckled, circular outline. Segments flat, rounded, broad, well overlapping. Corona bright orange red, paler at base, fluted, almost flat.
- DENHAM 2Y-O
Perianth yellow. Segments broad and overlapping, slightly ridged. Corona cup shaped, frilled and orange in colour. Neck grows in water.
- DEODORA 2W-WWY
Perianth cream, slightly creased. Segments broad, acute, edges slightly incurving, some notched. Corona cream at base becoming deep cream at rim, open mouthed to flat, fluted, frilled. Neck grows badly in water.
- DERVISH 2Y-O
Perianth sulphur yellow, overlaid apricot, deeper at base, creased, pointed outline. Segments flat, pointed, overlapping. Corona orange red, paler at base, goblet shape, small. Neck grows in water.
- DESDEMONA 2W-W
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and same colour as perianth. Neck grows in water.
- DEVILRY 2Y-O
Discarded - Flowers below foliage.
- DEVON LOCH 1W-W
Discarded - Flowers below foliage, hangs head.
- DEW-POND 2W-W
Discarded - Flowers below foliage, hangs head.
- DICK 1W
Perianth cream, stained yellow at base and on back, creased, star shaped. Segments twisted, pointed, barely overlapping. Corona bright lemon yellow, slender.
- DICK WELLBAND 2W-O
Perianth creamy white stained yellow at base. Segments flat, pointed, overlapping. Corona bright burnt orange, deeply lobed edge.
- DILIGENCE 1Y-Y
No description available.
- DILLENBURG 2Y-R
Perianth bright lemon yellow, creased, pointed outline. Segments twisted, pointed, separated, edges curl back. Corona bright orange, untidy deeply lobed edge.

- DINKIE** 3Y-GYR
Perianth sulphur yellow, paler at tips, smooth, circular outline, reflexed. Segments flat, rounded, overlapping. Corona green eye, changing to greenish lemon rim of bright orange red for 1/5 of its length, fluted, bowl shape. Neck grows badly in water.
- DINTON GIANT** 1W-Y
Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona creamy lemon, paler at base, crenate, flanged, broad mouth.
- DIOLITE** 2Y-YYO
Perianth pale lemon, slightly creased, very flat. Segments flat, pointed, overlapping. Corona deep lemon base, shading quickly to bright orange red edge for 1/2 its length, some white blotches on edge, fluted, straight sided. Neck grows in water.
- DIOTIMA** 1Y-Y
Perianth clear yellow, some green at back. Segments flat, pointed, separated, star shaped. Corona deep clear yellow, neat edge.
- DIPLOMAT** 2W-Y
Perianth creamy white, a little creased. Segments flat, pointed, overlapping. Corona opens pale lemon yellow, develops orange buff paler at base, some white blotches on edge, straight sided. Neck grows in water.
- DIVOT** 2W-Y
Perianth creamy white stained yellow at base, pointed outline. Segments slightly twisted, pointed overlapping. Corona deep bright lemon yellow.
- DOCTOR JOHANNES BEIJER** 2W-O
Perianth creamy white, stained yellow at base, deeply creased, buckled, pointed outline. Segments slightly twisted, pointed, overlapping. Corona bright orange red, crumpled, rough. Neck grows in water.
- DOCTOR PERON** 2W-GOO
Perianth creamy white stained yellow at base, edges curl in. Segments pointed, broad, overlapping. Corona bright orange shading to greenish gold centre, deeply lobed edge, fluted.
- DOE ROSS** 1Y-Y
Perianth pale sulphur, slightly twisted. Segments ovate, overlapping, appear translucent. Corona lemon, flanged, neat, frilled edge. Weak, hollow stems. Neck grows badly in water.
- DOLLY MOLLINGER** 11W-OWO
Discarded - Small, short.
- DOMINATOR** 1Y-Y
Perianth bright deep maximus gold, crêpey texture. Segments flat, pointed, overlapping. Corona very deep bright gold with deeply cut edge.
- DONIZETTI** 2W-Y
Discarded - Short, floppy flower.
- DONNA BELLA** 11Y-OOY
Perianth white. Segments very broadly ovate and overlapping. Corona very frilled and deep orange in colour. Neck grows badly in water.
- DONORE** 1Y-Y
Perianth clear lemon, slightly creased. Segments flat, pointed, overlapping. Corona deep lemon, frilled.
- DOUBLE EAGLE** 4W-O
Perianth white. Segments double, broad and overlapping. Corona orange, frilled and intermixed with inner whorl of perianth segments. Neck grows in water.
- DOUBLE EVENT** 4W-Y
Perianth creamy white, flat, rounded, broad, overlapping. 1 row complete with 6-10 narrow pointed, slightly twisted segments in centre, intermixed with corona segments of gold edged cream, circular outline. Neck grows badly in water.
- DOUBLE FORTUNE** 4Y-Y
Discarded - Not consistently double, brittle stem.
- DOUBLE MING** 4W-W
Perianth poeticus white, creased, crêpey texture, flat, rounded, overlapping, one complete row, circular outline. Centre of flower frilled with twisted green corona segments mixed with green stamen segments. Untidy. Neck grows badly in water.
- DOUBLEDALÉ** 4W-Y
Perianth creamy white. Segments rounded, overlapping. One complete row perianth segments, rest intermixed with pale cream corona segments, compact centre, edge frilled. Scented. Neck grows badly in water.

- DOUBLET** 4W-Y
Perianth poeticus white with stained base. Segments broad, round and overlapping. Segments intermixed with sulphur/white frilled corona segments. Compact centre. Neck grows badly in water.
- DOUBTFUL** 3Y-R
Perianth bright lemon, paler at tips, crêpey, bottom segments inclined to reflex, pointed outline. Segments flat, pointed, overlapping. Corona bright orange red, fluted, lightly frilled, small bowl shape. Neck grows stiffly in water.
- DOUMER** 1Y-Y
Perianth soft lemon yellow, creased, crêpey texture. Segments flat, pointed, overlapping. Corona deep bright lemon, flanged.
- DOVE WINGS** 6W-Y
Discarded - Short, small, cyclamineus hybrid.
- DOWNHILL** 3W-W
Discarded - Hangs head, flowers below foliage.
- DOWNPATRICK** 1W-Y
Perianth creamy white, stained yellow base. Segments broad, pointed, overlapping at base, barely creased, curled in at edges, double triangle shape. Corona pale bright lemon, creamy rim, belled edge. Good flower substance but perianth remains hooded.
- DRAGOMAN** 3W-R
Perianth poeticus white, stained yellow at base, sometimes notched, edges curl in, top three segments incurve. Segments flat, rounded, broad, well overlapping. Corona deep sealing-wax red, fluted, almost flat. Neck grows stiffly in water.
- DRAGOON** 3W-R
Perianth pure white stained bright yellow at base, floppy. Segments flat, pointed, separated. Corona bright orange red shading to paler orange base, fluted.
- DREAM GIRL** 2W-YYO
Discarded - Short, flowers below foliage.
- DREAMLIGHT** 3W-GWR
Perianth poeticus white, smooth, edges curl in a little. Segments flat, rounded, well overlapping. Corona greyish lemon with moss green eye and a thin pencil line of apricot orange on edge. Very long style.
- DREEN** 2W-P
Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona cream overlaid pale creamy peach, shading to creamy base, washy colour. Neck grows in water.
- DRENAGH** 2W-P
Perianth white. Segments broad and overlapping. Inner whorl has curled in edges. Corona open mouthed, frilled and pale pink in colour. Neck grows in water.
- DROMONA** 2W-Y
Perianth creamy white, stained at base. Segments round, broad bottoms, 'shovel' shaped, smooth, curled in at edges and 1/2 overlapping. Corona creamy lemon, slightly patchy, darker at base - bowl shaped with lobed, frilled edge. Poor arching neck.
- DROMORE** 2W-Y
Discarded - Perianth catches in corona, untidy, susceptible to basal rot.
- DRUMBOE** 2W-WWP
Perianth white, Segments broad, overlapping pointed. Corona white, tinged pink at edge, somewhat expanded, fluted.
- DUKE OF ANJOU** 1W-Y
Perianth creamy white, stained yellow at base, creased. Segments slightly twisted, pointed, overlapping. Corona bright lemon yellow, deeply lobed, wide mouth.
- DUKE OF BEDFORD** 1W-Y
Perianth creamy white stained yellow at base, star shaped. Segments twisted, pointed, barely overlapping. Corona deep clear lemon yellow.
- DUKE OF WINDSOR** 2W-OOY
Perianth creamy white, creased, edges curl in a little, sometimes caught in corona. Segments flat, rounded, overlapping. Corona apricot orange with narrow rim of creamy apricot, variable colour, fluted, frilled, deeply lobed, shallow bowl shape. Neck grows in water.
- DULCIMER** 9W-GYR
Perianth pure white, edges curl in a little. Segments flat, pointed, overlapping. Corona yellow changing to deepest red edge for 1/3 of its length, with large moss green eye, almost flat, fluted. Style as long as longest stamens. Neck grows in water.

- DUNCAN** 2Y-Y
Perianth creamy white, stained yellow at base, creased, pointed outline. Segments flat, pointed, overlapping. Corona opens clear lemon, develops primrose with cream edge, slightly fluted, lightly frilled, almost straight sided. Neck grows in water.
- DUNDARAVE** 2Y-R
Perianth bright lemon, circular outline. Segments broad, rounded, overlapping at base, creased, crepey, curled in at edges. Corona rich orange with very slight reddish tinge, bowl-shaped, fluted, deeply lobed edge.
- DUNDRUM** 1Y-Y
Perianth clear light lemon. Segments slightly twisted, pointed, broad, overlapping. Corona deeper lemon, crenate, flanged, broad mouth. Stiff.
- DUNEBA** 2Y-R
No description available.
- DUNFANE** 2W-W
Perianth creamy white, a little creased. Segments slightly twisted, pointed, overlapping. Corona opens creamy primrose, develops buff tone then ages to cream, of trumpet proportions. Stiff.
- DUNGANNON** 1Y-Y
Perianth sulphur, creased, crêpey. Segments flat, pointed, overlapping. Corona pale lemon, frilled.
- DUNGIVEN** 1Y-Y
Perianth light lemon, slightly creased. Segments slightly twisted, pointed, broad, well overlapping. Corona lemon, crenate, flanged.
- DUNKELD** 2Y-O
Perianth clear lemon yellow, pales a little with age, creased. Segments flat, rounded, well overlapping. Corona deep burnt orange, sometimes white blotches on edge, shallow, bowl shape, fluted.
- DUNLEWEY** 2W-W
Perianth creamy white, smooth, edges curl in. Segments flat, pointed, overlapping. Corona opens deep primrose yellow, develops buff tint, then turns creamy white.
- DUNLOY** 2W-P
Perianth creamy white, a little creased, edges curl in. Segments flat, pointed, broad, overlapping. Corona creamy peach shading to pale peach edge, variable colour. Neck grows in water.
- DUNLUCE** 1W-W
Perianth pure white, slight green base, a little creased. Segments flat, pointed, broad, well overlapping. Corona pure white. Neat, stiff and sturdy.
- DUNMINNING** 2Y-R
Discarded - Flowers below foliage.
- DUNMURRY** 1/2W-Y
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and canary yellow.
- DUNSEVERICK** 2W-W
Discarded - Severe virus in stock.
- DUTCH MASTER** 1Y-Y
Perianth clear mid lemon, a little green on back, slightly creased. Segments flat, pointed, broad, overlapping. Corona deeper bright lemon, paler at base, crenate, flanged. Neck grows in water.
- EARLY BIRD** 3Y
Perianth pale sulphur, slightly creased, slightly hooded. Segments rounded but coming to a point at the tip, only just overlapping. Corona deep orange, short, open mouthed, deeply cut rim, slightly frilled. Neck grows badly in water.
- EARLY BRIDE** 2W-YOO
Perianth cream, smooth, slightly hooded. Segments broad, pointed, overlapping, edges incurling. Corona egg yolk yellow, rim paler at very edge, almost flat, very fluted, rim very frilly. Neck grows in water.
- EARLY GLORY** 1Y-Y
Perianth pale primrose yellow, greenish back. Segments slightly twisted, pointed, barely overlapping. Corona bright clear gold, flanged. Neat.
- EARLY GOLD** 1Y-Y
Perianth canary yellow. Segments broad and overlapping. Corona wide open mouthed and frilled. Same colour as perianth.
- EARLY HIGHNESS** 1W-Y
Perianth white. Segments broad, overlapping and crêpey. Corona wide open mouthed, frilled and canary yellow in colour.

- EARLY LIGHT** 2Y-O
Perianth bright lemon developing paler at tips, very creased, sometimes notched, often buckled, inclined to remain incurve. Segments flat, pointed, overlapping. Corona bright orange, fluted, frilled, almost straight sided.
- EARLY MIST** 2W-W
Perianth white. Segments broad and overlapping. Corona open mouthed, flanged, frilled and white.
- EARLY PERFECTION** 8W-Y
Perianth white. Segments broad and overlapping. Corona cup shaped, frilled and yellow in colour. Several flowers/stem.
- EARLY PRINCE** 1W-Y
Perianth creamy white. Segments ovate, slightly twisted and overlapping. Corona open mouthed, frilled and yellow in colour. Neck grows in water.
- EARLY SENSATION** 1Y-Y
Perianth sulphur yellow, pointed outline. Segments flat, pointed, overlapping. Corona clear light lemon yellow.
- EARLY SPLENDOUR** 8W-O
Perianth creamy white, stained yellow base, creased, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona moss green eye changing to bright orange red, fluted, lightly frilled. 4-9 florets well spaced.
- EARLY SURPRISE** 3W-R
Perianth opens primrose, develops cream stained yellow at base, creased, notched, pointed outline. Segments flat, pointed, barely overlapping. Corona gold base shading to orange red for 1/2 its length, fluted, nearly flat, uneven lobed edge. Neck grows in water.
- EASTER BONNET** 2W-P
Discarded - Short, flowers below foliage.
- EASTER BRIDE** 1W-W
Perianth white, creased. Segments flat, pointed, broad, well overlapping. Corona whiter than perianth. Neat but hangs head.
- EASTER JOY** 1Y-Y
Perianth sulphur, creased. Segments flat, pointed, overlapping. Corona mid lemon, frilled, flanged, broad mouth, dumpy appearance.
- EASTER MOON** 2W-GWW
Perianth creamy white, smooth. Segments rounded, broad, overlapping. Corona creamy white, short, straight sided, slightly wider at mouth, gently fluted, gently wavy rim. Neck grows badly in water.
- EASTER MORN** 2W-W
Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona opens primrose, developing to creamy peach with narrow cream edge, ageing to cream, of trumpet proportions.
- ECHO** 2W
Discarded - Short.
- EDDY CANZONY** 2W-YYO
Perianth poeticus white, a little creased, circular outline. Segments flat, rounded, broad, well overlapping. Corona deep orange red, paler at base, deeply fluted, deeply cut lobes almost into separate segments, almost flat. Neck grows stiffly in water.
- EDENBANK** 3W-R
Perianth white. Segments broad and overlapping. Corona cup shaped, frilled, yellow in centre with a deep orange rim. Neck grows badly in water.
- EDWARD BUXTON** 3Y-OOR
Perianth pale creamy primrose. Segments flat, rounded, overlapping. Corona deep orange shading to bright red at edge, fluted, lightly frilled. Neck grows badly in water.
- EFFECTIVE** 1W-Y
Perianth cream, stained deep yellow for half its length at base, a little creased. Segments flat, pointed, overlapping. Corona bright intense lemon yellow, slender, good colour contrast.
- EGGS AND BACON** 4W-O
Perianth white. Segments double, ovate and overlapping. Corona double, orange in colour and intermixed with inner perianth segments.
- EL PASO** 2Y-O
Perianth bright clear lemon yellow, creased, crêpey texture. Segments flat, pointed, overlapping. Corona bright deep orange, paler at base, white blotches on edge.
- ELABORATE** 2Y-O
Perianth yellow. Segments broad and overlapping, with ridged texture. Corona cup shaped, frilled, yellow with an orange rim. Neck grows badly in water.

- ELEANOR MAY 4W-R
No description available.
- ELF 2Y-WWP
Discarded - Small, short, flowers below foliage.
- ELGIN 1Y-Y
Perianth clear yellow. Segments pointed, overlapping, hooded, star shaped, crêpey texture. Corona very deep clear lemon yellow, open-mouthed, rolled back edge.
- ELIZABETH BANDON 2W-R
Perianth creamy white stained yellow at base, edges curl in. Segments flat, pointed, overlapping. Corona bright solid deep orange-red.
- ELLESMERE 2Y-YOO
Perianth clear sulphur yellow, edges curled in giving star-shape appearance. Segments flat, pointed, overlapping. Corona bright orange shading to greenish lemon base, open bowl shape, deeply fluted.
- ELMWOOD 2Y-R
Perianth yellow. Segments very broadly ovate and overlapping. Corona open mouthed, frilled and orange in colour. Neck grows in water.
- ELSPETH 3W-YYO
Perianth creamy white, edges curl in. Segments flat, rounded, well overlapping. Corona pale lemon changing to bright orange edge for 1/4 of its length.
- ELVIRA 8W-YYO
Perianth creamy white, slightly creased, curled in at edges. Segments flat, rounded, well overlapping. Corona bright greenish lemon with thin pencil line of orange at edge, almost flat, fluted. Style reaches half way between long and short stamens.
- EMBER 2Y-O
Perianth yellow. Segments ovate and overlapping. Corona cup shaped, frilled and deep orange in colour. Neck grows badly in water.
- EMPEROR 1Y-Y
Perianth pale dirty primrose yellow, deeper yellow strip down centre back, crêpey texture, slightly hooded. Segments twisted, pointed, overlapping. Corona lemon yellow, lobed.
- EMPIRE 2W-WYY
Perianth creamy white, creased, notched edge. Segments flat, pointed, overlapping. Corona opens creamy lemon, develops to pale cream with paler edge.
- EMPRESS 1W-Y
Perianth creamy white, stained yellow at base, creased, pointed outline. Segments flat, pointed, overlapping. Corona very deep lemon yellow, very slender. Wiry and stiff.
- EMPRESS OF IRELAND 1W-W
Perianth white. Segments broad and overlapping, with ridged texture. Corona, open mouthed, frilled and same colour as perianth.
- ENCHANTRESS 2W-P
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled pink with yellow centre.
- ENGADINE 3W-W
Perianth creamy white, edges curl in, crêpey texture, flat, rounded, broad, well overlapping. Corona white, bowl shape, fluted, hangs head badly. Neck grows in water.
- ENNISKILLEN 3W-R
Perianth creamy white, stained yellow base, smooth, edges curl in at tips. Segments flat, pointed, broad, overlapping. Corona small greenish gold eye changing to deep sealing wax red, fluted, lightly frilled, shallow bowl shape. Neck grow stiffly in water.
- ENTERPRISE 4Y-O
Perianth bright lemon, with darker centre line. Double perianth segments round and 1/2 overlapping, and curled in at edges. Corona segments white/orange frilled and loose. Neck grows badly in water. Weak stems.
- ENTRANCEMENT 1Y-W
Perianth pale primrose becoming cream near base, twisted. Segments would fit into a double triangle, small. Corona white, narrow, flanged, edge frilled. Neck grows a little.
- EPIC 9W-YOR
Perianth white, creased a little. Segments flat, rounded, overlapping. Corona greenish lemon with thin pencil line of deep red at edge, and a moss green eye, almost flat, fluted. Long style.
- ERIE 2Y-O
Perianth pale primrose yellow, suffused apricot - variable colour. Segments flat, rounded, overlapping. Corona deep red with white blotches on edge, fluted, double frilled edge.

ERLICHEER 4W-Y
Discarded - Short.

ERNANI 2W-YYO
Discarded - Short, poor colour, susceptible to basal rot.

ESKIMO 2W-W
Perianth white. Segments shovel shape, flat, overlapping. Corona opens primrose, develops primrose-buff.

ESTELLA DE MOL 11W-Y
Discarded - Short.

ETHEL 3W-Y
Perianth white. Segments broad and overlapping. Corona cup shaped, frilled, yellow in centre with orange rim. Neck grows badly in water.

EVANGELINE 3W-Y
Perianth creamy white, curled in at edges. Segments flat, pointed, barely overlapping. Corona pale soft lemon yellow developing to deep cream greyish base.

EVE DALY 1W-Y
Perianth white. Segments broad, overlapping and inflexed. Corona flared open mouthed, frilled and pale yellow in colour. Neck grows in water. Weak stems.

EVENING 2W-W
Perianth creamy white, a little creased, edges curl in, shovel shape segments. Segments flat, pointed, broad, overlapping. Corona pale cream, neat.

EVERTON 2Y
Perianth yellow. Segments broadly ovate, overlapping with edges turned in. Corona cup shaped, frilled and orange in colour. Neck grows in water.

EVITA PERON 2W-O
Discarded - Short, poor colour, flowers below foliage.

EVOLUTION 11Y-Y
Perianth creamy white, star shape. Segments flat, pointed, separated. Corona light lemon yellow often edged cream, split into six segments, not always evenly, very broad, floppy, top-heavy.

EXPLORER 1Y-Y
Discarded - Short, rough flower.

EXPRESSIVE 1Y-Y
Discarded - Short, hangs head.

EXTOL 4Y-R
Perianth yellow. Segments double layer, broad and overlapping with darker central ridge. Corona deep orange, split and intermixed with inner perianth segments. Neck grows in water.

E.E. MORBEY 2Y-O
Perianth yellow. Segments broad and overlapping. Corona cup shaped, frilled, yellow in centre and orange around rim. Neck grows badly in water.

FAIR COLLEEN 3W-OOY
Perianth poeticus white, slightly creased, circular outline. Segments flat, rounded, well overlapping. Corona creamy apricot orange with wire rim of cream, slightly fluted, lightly frilled, almost flat. Neck grows badly in water.

FAIR GREEN 3W-GYO
Perianth white. Segments rounded and overlapping. Corona small cup, frilled, green eyed, yellow with an orange rim. Neck grows badly in water.

FAIR MAIDEN 3W-YYO
Perianth creamy white, crêpey texture. Segments flat, rounded, overlapping. Corona greenish lemon yellow with shaded pale orange edge.

FAIR TRIAL 1W-Y
Discarded - Short.

FAIRLIGHT 1Y-Y
Perianth pale yellow. Segments broad, overlapping and slightly inflexed. Corona trumpet frilled and darker yellow. Neck grows in water.

FAIRMILE 3W-GYO
Perianth creamy white, base slightly stained yellow, full rounded outline. Segments rounded, overlapping at base, broad, slightly creased. Corona moss green eye changing to creamy yellow with deep orange rim, bowl-shaped, deeply fluted.

FAIRNESS 4W-O
Perianth segments pale sulphur yellow, often greenish on back, twisted, pointed, overlapping. Centre very full and intermixed with very short corona segments of orange red.

FAIRY DREAM 1W-W
Discarded - Short, rough flower.

FAIRY KING 2Y-O
Perianth clear lemon yellow, creased. Segments flat, rounded, overlapping. Corona bright burnt orange, goblet shape. Neat flower. Neck grows in water.

FAIRY TALE 3W-YYO
Perianth creamy white, slightly creased, edges curl in a little. Segments flat, rounded, overlapping. Corona greyish eye, changing to creamy white with narrow rim of apricot, fluted, lightly frilled, bowl shape. Neck grows badly in water.

FAITHFUL 2Y-Y
Perianth clear lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona deeper bright lemon yellow, of trumpet proportions, well balanced, stiff and sturdy.

FALAISE 4W-O
Perianth segments poeticus white stained yellow at base, complete outer row, flat, pointed, broad, overlapping. Corona segments deep orange red, short, frilled and curled up in centre intermixed with 6-8 long pointed perianth segments.

FALSTAFF 2Y-R
Perianth yellow. Segments ovate and overlapping. Corona cup shaped, frilled and deep orange in colour. Neck grows badly in water.

FAMILLE ROSE 2W
No description available.

FANCHEON 2W-YOR
Perianth pale primrose yellow, edges curled in, pointed outline. Segments flat, pointed, overlapping. Corona bright orange red shading to gold base, open mouthed.

FANCY FREE 2W-YYP
Discarded - Rough, flowers below foliage.

FANTEE 1Y-Y
Perianth yellow. Segments broad and overlapping, with ridged texture. Corona flared open mouthed, frilled and same colour as perianth. Neck grows in water.

FAREWELL 2W-Y
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona creamy lemon, almost straight sided. Neat.

FASHION 11Y-YYO
Perianth primrose. Segments round, overlapping and very crêpey and curled in at edges. Corona pale lemon with orange central stripe with slightly frilled edge. Corona 6 segments split to base. Neck grows badly in water.

FASTNET 2W-YYR
Perianth creamy white, a little creased, edges curl in. Segments flat, pointed, broad, overlapping. Corona creamy primrose base shading to primrose with narrow apricot orange edge, slightly fluted, bowl shape. Hangs its head when young. Neck grows in water. Poor poise.

FAVOURITE 2W-Y
Perianth creamy white. Segments broad, overlapping and edges turned in. Corona cup shaped, frilled and yellow. Neck grows in water.

FAWNGLO 2Y-Y
Discarded - Short, flowers below foliage.

FAY MARK 2W-O
Perianth creamy white, creased, notched, top 3 segments often caught in corona, buckled, circular outline. Segments flat, rounded, well overlapping. Corona bright apricot orange with narrow edge of apricot, fluted, heavily frilled in three layers. Rough.

FEBRUARY BICOLOUR 2W-Y
Perianth white. Segments ovate, notched and overlapping - ridged texture. Corona large, open mouthed, frilled, dark yellow centre getting paler towards rim.

FEBRUARY GOLD 6Y-Y
Perianth bright golden yellow, slightly creased, reflexed. Segments flat, pointed, barely overlapping. Corona deep bright gold, almost straight sided.

- FEBRUARY SILVER** 6W-W
Discarded - Short, cyclamineus hybrid.
- FEDORA** 2W
Perianth creamy white, creased, edges curl in. Segments flat, pointed, overlapping. Corona greenish lemon base, shading to orange edge for 1/3 its length, fluted, shallow bowl shape.
- FELINDRE** 9W-GYR
Perianth poeticus white, edges curl in, notched at base, circular outline. Segments flat, rounded, well overlapping. Corona bronze-green base 1/3, changing to gold 1/3, changing to deep orange red edge, fluted, lightly frilled, almost flat. Very long style.
- FERMOY** 2W-YOO
Perianth white, stained yellow at base, smooth, edges curled in. Segments flat, pointed, broad overlapping. Corona gold shading to bright orange edge for 3/5 of its length, fluted. Neck grows in water.
- FERNIE** 3W
Perianth opens primrose, develops creamy white stained yellow for 1/3 at base, creased, sometimes notched, bottom segments reflexed. Segments flat, rounded, well overlapping. Corona orange gold base 2/3 changing to orange red edge following deep cut lobes to centre of corona, fluted, frilled, almost flat. Neck grows badly but stiffly in water.
- FERNISKY** 2W-W
Discarded - Short, rough flower.
- FESTIVITY** 2W-Y
Perianth creamy white. Segments broad and overlapping. Inner 3 segments edges curled in. Corona open mouthed, frilled and yellow in colour. Neck grows badly in water.
- FEU DE JOIE** 4W-O
Perianth segments white, twisted, pointed, separated with 12 long narrow segments intermixed with corona segments which are gold shading to bright red for 1/2 its length. Spidery looking.
- FIELD MARSHAL** 2Y-O
Perianth canary yellow. Segments very broadly ovate and overlapping. Corona cup shaped, frilled and orange. Neck grows in water.
- FIERY CHARIOT** 2Y-R
Perianth deep warm lemon, tinged orange, paler at tips, rounded outline. Segments rounded, overlapping at base, bottom segments broad, slightly creased, curled in at edges. Corona burnt orange with reddish tinge, bowl-shaped, frilled edge. Very good flower substance.
- FIERY KNIGHT** 4Y-O
Perianth segments soft lemon yellow, flat, pointed, separated, in 2 rows with 2 or 3 curled segments in centre. Corona segments orange, intermixed.
- FINCOOL** 1W-W
No description available.
- FINE GOLD** 1Y-Y
Perianth lemon yellow, smooth. Segments flat, pointed, broad, well overlapping, shovel shape. Corona deeper lemon yellow. Stiff and neat.
- FINGLAS** 3W-WWY
Perianth creamy white, edges curl in. Segments slightly twisted, pointed, overlapping. Corona grey eye changing to greyish cream with wire rim of apricot orange, slightly fluted, frilled, lobed, bowl shape. Neck grows stiffly in water.
- FINLAND** 2W-Y
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, broad, overlapping. Corona opens light lemon yellow, develops to creamy primrose, flanged.
- FINMACOOL** 2W-W
Perianth white. Outer segments broad, inner narrower, and 1/2 overlapping. Corona straight flared, slightly frilled and white in colour.
- FINOLA** 1W-W
Perianth white. Segments broad, overlapping and slightly inflexed with yellow stain in centre. Corona open mouthed, rolled, frilled and white in colour. Neck grows in water.
- FINTONA** 2W-P
Perianth white. Segments very broadly ovate, overlapping and inflexed. Corona open mouthed, frilled and pink.
- FIRE CHIEF** 2Y-O
Perianth sulphur, smooth, some segments incurling at edge. Segments very broad but with distinct point, overlapping to half of length. Corona orange, short, funnel shaped, crenate, loosely frilled rim. Hollow stems. Neck grows a little.

FIRE GLEAM Discarded - Small, poor flower, long neck.	2Y-O
FIRE GUARD Discarded - Small, poor flower.	2Y-YOR
FIRE ISLAND Perianth yellow. Segments round and overlapping. Corona cup shaped, frilled and deep orange.	2Y-O
FIREBRAND Perianth pale primrose, deeper at base, a little reflexed. Segments flat, pointed, separated. Corona deep orange throughout, fluted.	3WWY-R
FIRECRACKER Discarded - Small.	2Y-O
FIREGLOW Perianth sulphur yellow, creased, edges curled in. Segments flat, pointed, overlapping. Corona bright orange red, white blotches on edge, fluted, almost flat. Neck grows badly in water.	2Y-O
FIREMASTER Discarded - Small.	2Y-O
FIRENZE No description available.	4W-Y
FIREPROOF Perianth bright lemon, stained orange at base, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona deep orange throughout, fluted, lightly frilled, almost straight sided, small in relation to perianth. Neck grows in water.	2Y-O
FIRETAIL Perianth opens primrose, develops cream stained yellow at base, deeply creased, star shape. Segments flat, pointed, overlapping. Corona burnt orange red, fluted, nearly flat, very small in relation to perianth. Neck grows badly in water.	3W-R
FIREWATER Discarded - Low flower production.	3W-YYO
FIRST FROST Perianth white. Segments ovate and overlapping. Corona funnel shaped, lobed and white in colour.	2W-W
FLAMENCO Perianth cream, stained yellow at base, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona bright solid orange, fluted. Neck grows in water.	2W-O
FLAMING METEOR Perianth yellow. Segments ovate, overlapping, ridged texture. Corona cup shaped, frilled and deep orange. Neck grows badly in water.	2Y-R
FLAMING MORN Perianth yellow. Segments ovate, overlapping and ridged texture. Corona open mouthed, frilled, yellow in centre with orange rim. Neck grows badly in water.	2Y
FLANEUR Perianth pale lemon yellow. Segments flat, pointed, separated. Corona lemon yellow, split into six segments, the top of which is divided into two lobes, often unevenly, three inner segments incurve, very broad.	11Y-Y
FLASH Perianth yellow. Segments very broadly ovate and overlapping, with white tips. Corona cup shaped, frilled and orange in colour. Neck grows badly in water.	2Y-O
FLEURIMONT Perianth primrose yellow, deeply creased but very stiff. Segments flat, pointed, broad, well overlapping. Corona burnt orange throughout, fluted.	2Y-YYR
FLEVO Perianth yellow. Segments broad and overlapping - ridged texture. Corona large, open mouthed, frilled and same colour as perianth.	1Y-Y
FLORA TRISTAN Perianth pale lemon, developing paler with age, with whitish tips, creased, buckled, crêpey texture. Segments flat, rounded, well overlapping. Corona bright orange shading to paler base, very broad, fluted. Rough looking flower.	2Y-O
FLORA'S FAVOURITE Discarded - Short, floppy, flowers below foliage.	2W-Y

- FLORIADE** 2Y-O
Discarded - Untidy flower.
- FLOWER CARPET** 1Y-Y
Perianth clear yellow, a little green at back. Segments flat, pointed, overlapping. Corona deeper clear yellow, neat edge. Neck grows in water.
- FLOWER DRIFT** 4W-OYO
Perianth white. Double segments very broadly ovate and overlapping. Double corona deep orange. Inner perianth segments have light orange streaks. Neck grows in water.
- FLOWER RECORD** 2W-WWO
Perianth opens pale primrose, develops creamy white, deeply creased. Segments flat, pointed, overlapping. Corona deep lemon gold edged bright deep orange for 1/2 its length. Neck grows in water.
- FLOWERSONG** 2Y-YYO
Perianth lemon, slightly creased, some segments cupped, some segments reflexed. Segments acute, overlapping at base. Corona deep lemon at base becoming gold for outer half, short, open mouthed, frilled edge. Neck grows badly.
- FLUSH** 2W-P
Perianth white, slightly creased, star shaped. Segments flat, pointed, overlapping. Corona opens pale pink develops to clear peach with apricot shaded base, ages to buff primrose with pink edge, often with white blotches on edge, deeply frilled.
- FLYING SAUCER** 2W-Y
Perianth white. Segments broad and overlapping with ridged texture. Corona large cup shape, frilled and yellow in colour. Neck grows badly in water. Weak stems.
- FOGGY DEW** 3W-W
Perianth creamy white, creased, edges curl in. Segments flat, rounded, well overlapping. Corona creamy white with moss green eye, fluted, deeply frilled and lobed edge. Neck grows stiffly in water. Hangs head badly.
- FOLLY** 2W-O
Perianth white. Segments broad and overlapping. Corona cup shaped, frilled yellow in centre with orange rim. Neck grows badly in water.
- FONDANT** 2W-Y
Perianth white. Segments broad, overlapping ridged texture and turned in at edges, slightly inflexed. Corona open mouthed, frilled and yellow.
- FORAY** 2W-WWP
Perianth white. Segments ovate and overlapping. Corona large cup, frilled, yellow in centre with orange rim. Neck grows badly in water.
- FORERUNNER** 1Y-Y
Perianth bright golden yellow, with centre line of green on back. Segments twisted, pointed, separated. Corona deep gold, slightly paler at base.
- FORESCATE** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona large open mouthed, frilled, and same yellow as perianth.
- FORESIGHT** 1W-Y
Perianth cream, smooth, slightly twisted. Segments would fit into a double triangle, small and neat. Corona lemon, develops cream, flanged, uneven, slightly frilled rim. Neck grows badly.
- FOREST FIRE** 2Y-O
Perianth pale sulphur yellow, creased, crêpey texture, notched. Segments flat, pointed, overlapping. Corona bright orange red, fluted, bowl shape. Neck grows badly in water. Poor poise.
- FORFAR** 3W-O
Perianth creamy white, stained slight yellow at base. Segments flat, rounded, overlapping. Corona bright orange red, a little paler at base.
- FORTHRIGHT** 2Y-O
Perianth deep lemon, slightly creased, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona mid to deep orange, variable colour, lightly frilled, broad mouth. Stiff. Neck grows in water.
- FORTISSIMO** 2Y-O
Perianth pale yellow. Segments broad and overlapping, ridged texture. Corona open mouthed, frilled and deep orange. Neck grows badly in water.
- FORTUNE** 2Y-O
Perianth clear lemon, creased, sometimes caught in corona. Segments flat, pointed, broad, overlapping. Corona orange shading paler at base, variable colour, frilled, lobed. Neck grows badly in water.

- FORTUNE'S BLAZE** 2Y-O
Perianth pale lemon yellow, segments sometimes creased, edges curl in. Segments flat, rounded, overlapping. Corona deep burnt orange red, goblet shape.
- FORTUNE'S CREST** 2Y-O
Perianth pale sulphur yellow, developing paler with age, smooth. Segments flat, pointed, broad, overlapping. Corona bright orange red, fluted.
- FORTUNE'S SUN** 2Y-O
Perianth primrose yellow, smooth. Segments flat, pointed, broad, overlapping. Corona bright orange, fluted.
- FORTWILLIAM** 1Y-Y
Perianth yellow. Segments very broadly ovate and overlapping. Corona flared, frilled and same colour as perianth. Neck grows in water.
- FOXHUNTER** 2Y-O
Perianth lemon, creased, crêpey, pointed outline. Segments flat, pointed, overlapping. Corona bright orange red, slightly fluted, lightly frilled, bowl shape. Neck grows stiffly in water.
- FOYLE** 2W-P
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona opens creamy primrose, develops pale creamy peach with cream edge, ageing to pale cream, indefinite colour. Neck grows in water.
- FRA DIAVOLO** 1Y-Y
Discarded - Ugly, badly proportioned.
- FRANCISCUS DRAKE** 2W-YYO
Perianth creamy white stained deep yellow at base, creased. Segments twisted, pointed, overlapping. Corona deep burnt orange, fluted. Rough.
- FRANKLIN** 2Y-Y
Perianth light lemon yellow, paler at tips, creased. Segments flat, broad, well overlapping. Corona bright lemon yellow of almost trumpet proportions.
- FRENCH SOL** 8Y-Y
No description available.
- FRIENDLY** 2Y
Perianth sulphur, slightly creased. Segments flat, pointed, broad, overlapping. Corona clear lemon, frilled, flanged, broad mouth.
- FRIGID** 3W-GGW
Perianth poeticus white, slightly crêpey, edges curl in at tips, pointed outline. Segments flat, pointed, overlapping. Corona bright green base 1/3 changing to pure white, fluted, frilled, lobed. Neck grows stiffly in water.
- FRILEUSE** 11W-Y
Discarded - Small, short.
- FRILLED BEAUTY** 2Y-Y
Perianth dirty sulphury-white, creased. Segments slightly twisted, pointed, overlapping. Corona dirty sulphury white with shaded apricot edge and white blotches, unevenly and untidily frilled.
- FROLIC** 1W-Y
Discarded - Short, untidy corona edge, flowers below foliage, very susceptible to basal rot.
- FRONTIER** 1Y-Y
Discarded - Short, weak stem, susceptible to basal rot.
- FROST AND FLAME** 3W-R
Perianth white. Segments very broadly ovate and overlapping. Corona cup shaped, frilled, yellow centre with orange rim.
- FROZEN** 2W-W
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona creamy white, grey-green eye, bowl shape. Stiff and neat.
- FURBELOW** 4Y-O
Perianth canary yellow, paler at tips. Segments broad and round, 1/2 overlapping and slightly buckling on centre line. Inner row segments intermixed with frilled corona. Corona gold/orange colour. Neck grows badly in water.
- FURY** 2Y-O
Perianth mid lemon, creased, pointed outline. Segments flat, pointed, overlapping. Corona deep burnt orange red, fluted, frilled, bowl shape. Hangs its head a little. Neck grows badly in water.

- GADDESBY** 2W
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona pale greenish lemon, overlaid apricot orange in vertical stripes, with pale apricot rim, variable colour, fluted, deep bowl shape. Neck grows in water.
- GAIL O'KAHN** 2W-Y
Discarded - Floppy, badly balanced.
- GALA** 2W-O
Discarded - Hangs head.
- GALASHIELS** 1Y-Y
Perianth lemon, slightly twisted, some segments notched. Segments acute, would fit into a double triangle. Corona deep lemon, large in comparison to perianth, trumpet with open mouth, slightly frilled. Bold flower.
- GALAXY** 2W-P
Discarded - Short, flowers below foliage.
- GALCADOR** 2Y-O
Perianth deep primrose yellow, crêpey substance, edges curl in a little. Segments flat, rounded, well overlapping. Corona deep burnt orange with slight orange edge. Stiff and sturdy.
- GALILEE** 3W-GWO
Perianth white, crêpey, edges curl in slightly. Segments flat, pointed, broad, well overlapping. Corona grey eye changing to greyish cream with narrow rim of apricot orange flecked cream for 1/5 of its length, fluted, frilled, lobed. Neck grows badly in water.
- GALWAY** 2Y-Y
Perianth bright deep yellow. Smooth, perfect double triangle. Segments flat, pointed, broad, well overlapping. Corona deeper bright yellow, of trumpet proportions.
- GARDEN GIANT** 2Y
Perianth sulphur yellow, deeply creased, edges curl in, circular outline. Segments flat, rounded, well overlapping. Corona gold changing abruptly to orange red edge for 2/3 its length, deeply fluted, saucer shape. Neck grows stiffly in water.
- GARDEN PRINCESS** 6Y-Y
Perianth deep clear yellow, creased. Segments twisted, pointed overlapping. Corona deeper yellow, open mouthed. Not a typical cyclamineus hybrid.
- GARDENVALE** 2W-OOY
Perianth poeticus white, edges slightly incurling. Segments broad, rounded and overlapping. Corona flat, orange becoming yellow at very edge. Neat and fresh looking flower. Neck grows badly in water.
- GARLAND** 2Y-YOO
Perianth clear sulphur yellow, a little creased. Segments flat, pointed, broad, overlapping. Corona bright burnt orange-red throughout, almost flat, deeply fluted. Neck grows in water. Stiff and sturdy.
- GARRON** 1Y-Y
Perianth soft sulphur yellow, star shaped. Segments flat, pointed, broad, well overlapping. Corona clear lemon yellow.
- GARTAN** 2Y-O
Perianth deep lemon yellow, creased, crêpey texture, edges curl in. Segments flat, pointed, overlapping. Corona opens yellow, develops orange gold, fluted, stiff.
- GATH-A-BAWN** 2W-W
No description available.
- GAUCHO** 2Y-O
Perianth sulphur yellow, creased. Segments flat, pointed, broad, overlapping. Corona bright burnt orange, paler at base, fluted, bowl shape. Stiff.
- GAY BOY** 2Y-R
Perianth light lemon yellow, edges curl in a little. Segments flat, pointed, broad, well overlapping, shovel shape. Corona gold base for 1/3 its length, shading to orange red edge, fluted, bowl shape. Stiff. Neck grows in water.
- GAY COLOURS** 2W-R
Perianth creamy white, stained yellow base, circular outline. Segments rounded, overlapping at base, fairly smooth, some crinkled at edges, curled in at edges. Corona bright orange, paler at base, fluted with white blotches at edge, bowl-shaped, lobed edge.
- GAY RECORD** 4W-O
Perianth poeticus white, with stained yellow base. Segments round and well overlapping. Inner row buckled/folded and intermixed with short yellow/orange frilled corona segments. Neck grows badly in water.

- GAY TIME** 4W-R
Perianth opens cream, develops white, flat, pointed, broad, overlapping, 1 row complete with 8-10 narrower pointed segments in centre, intermixed evenly with corona segments of orange shading to deep red edges. Neck grows stiffly in water.
- GAYLORD** 2Y-YOO
Perianth clear mid-lemon. Segments well rounded, overlapping, creased, slightly folded and crepey at edges. Corona orange gold changing to mid-orange with reddish tinge at edge; edge lobed. Neck grows badly in water.
- GEHEMA** 2Y-Y
Perianth pale creamy primrose, creased, star-shaped. Segments flat, pointed, overlapping. Corona greenish lemon shading to pale orange edge for 1/3 of its length, open mouthed, fluted, washy colours.
- GEISHA** 2W-P
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and very pale pink. Neck grows in water.
- GEM OF ANTRIM** 2W-P
Perianth creamy white, slightly hooded. Segments outer 3 broad, rounded, inner 3 more acute, overlapping. Corona deep primrose at base, shading to coral at rim, short, straight sided, fluted giving wavy edge. Neck grows in water.
- GEM OF ULSTER** 2W-P
Perianth white. Segments broad and overlapping - ridged texture. Corona cup shaped, frilled yellow with orange rim. Neck grows in water.
- GENERAL PATTON** 1W-Y
Perianth creamy white, creased, sometimes buckled. Segments twisted, pointed, overlapping. Corona pale creamy lemon, broad mouthed.
- GENERAL SMUTS** 2Y-O
Perianth lemon yellow, slightly creased. Segments flat, pointed, broad, overlapping. Corona bright burnt orange, wide mouth, fluted. Neat and stiff.
- GENTILITY** 2W-W
Perianth creamy white, smooth, edges curl in a little. Segments flat, pointed, overlapping. Corona creamy white, trumpet proportion, barely frilled. Hangs head when young.
- GEORGE ARMSTRONG** 2Y-Y
Discarded - Short, poor colour, flowers below foliage.
- GEORGE LEAK** 2W-O
Perianth poeticus white, smooth, edges incurled. Segments would fit into a double triangle. Corona orange, flat with ragged edge, crenate. Neck grows badly in water.
- GEORGE SAND** 2W-O
Discarded - Short, flowers below foliage.
- GEORGIA MOON** 2Y-Y
Perianth pale yellow. Segments broad and overlapping. Corona open mouthed, frilled, and same colour as perianth. Neck grows badly in water. Weak stems.
- GERANIUM** 8W-O
Perianth creamy white, stained slight yellow at base, a little creased. Segments flat, rounded, overlapping. Corona bright deep orange with greenish eye, fluted. Style a little longer than stamens.
- GERTIE MILLAR** 2W-Y
Perianth creamy white, creased, crêpey substance. Segments flat, pointed, broad, overlapping. Corona opens primrose yellow, develops primrose buff, deeply frilled edge.
- GERVO** 2W-W
Perianth cream, creased, crêpey texture, star shape. Segments flat, pointed, overlapping. Corona pale cream with deeper, cream edge for 1/4 its length, green eye, broad mouth, fluted, deeply frilled. Rough.
- GHANA** 2Y-Y
No description available.
- GIANT MUTICUS** 1Y-Y
Discarded - Flowers below foliage, badly proportioned.
- GIBRALTAR** 2Y-O
Perianth lemon yellow, creased, one segment often buckled. Segments flat, pointed, broad, overlapping. Corona deep orange red, shading to gold base, fluted, almost straight sided. Neck grows stiffly in water.
- GILFORD** 3W-O
Perianth creamy white, stained yellow base, notched, back segments reflex, pointed outline. Segments flat, pointed, overlapping. Corona moss green eye changing to deep greenish lemon base 1/2, changing to deep orange red edge, deeply fluted, frilled, lobed, almost flat. Neck grows badly in water.

GIOVANNI	2W-O
Perianth white. Segments outer layer broad, inner layer ovate. Corona large cup, frilled and orange in colour. Neck grows badly in water.	
GIPSY MOTH	2W-W
No description available.	
GLACIER	1W-W
Discarded - Short, flowers below foliage.	
GLANDORE	2W-W
No description available.	
GLEAMING	2Y-O
Perianth deep bright yellow, smooth, edges curl in a little. Segments flat, pointed, overlapping. Corona bright orange red.	
GLEN FIRE	2W-O
Perianth cream, stained deep yellow at base, creased. Segments flat, pointed, overlapping. Corona bright deep orange throughout almost straight sided.	
GLEN SPRINGS	2W-YYO
Perianth creamy white, stained yellow at base, creased. Segments slightly twisted, pointed, broad, overlapping. Corona gold with edge of bright orange for 1/3 of its length, fluted. Stiff.	
GLENBANE	1W-W
Perianth creamy white, a little crêpey, pointed outline. Segments flat, pointed, overlapping. Corona whiter than perianth, neat edge.	
GLENBURVIE	2W-OOR
Perianth white. Segments very broadly ovate and overlapping. Corona cup shaped, frilled yellow with orange rim. Neck grows badly in water.	
GLENBUSH	2W-W
Discarded - Short, hangs head badly, susceptible to basal rot.	
GLENCAIRN	1W-Y
Perianth creamy white, creased, pointed outline. Segments flat, pointed, overlapping. Corona opens creamy lemon, develops buff yellow, rolled back edge.	
GLENDALOUGH	2W-W
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona whiter than perianth, smooth, almost straight edged.	
GLENDERMOTT	2W-W
Perianth white. Segments broad and overlapping, slightly notched. Corona open mouthed, frilled, with green eye and white segments. Neck grows in water.	
GLENGARRIFF	1W-YYW
Perianth creamy white, a little creased, pointed outline. Segments flat, pointed, overlapping. Corona creamy lemon yellow with narrow rim of cream, hangs head badly.	
GLENGORMLEY	2W-OOY
Discarded - Rough, flowers below foliage.	
GLENLESLIE	2W-W
Perianth creamy white, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona creamy white, almost straight sided, almost straight edged. Neck grows in water. Scented.	
GLENMANUS	2W-W
Perianth creamy white, slightly creased and crêpey. Segments flat, pointed, well overlapping. Corona creamy buff developing creamy white, almost straight sided, barely frilled. Hangs head when young. Stiff. Neck grows in water.	
GLENOCUM	2W-W
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and creamy white. Neck grows in water.	
GLENSHANE	2W-W
Perianth white. Segments broad, overlapping and ridged. Corona cup shaped, frilled and lemon yellow. Scented. Neck grows in water.	
GLENSHESK	1W-W
Discarded - Short, flowers below foliage.	
GLENTIES	2W-O
Perianth white. Segments broad, overlapping and reflexed. Corona cup shaped, frilled and orange. Neck grows badly in water.	

- GLENVAR** 3W-R
Perianth poeticus white, stained yellow base, slightly creased, inclined to reflex. Segments flat, pointed, overlapping. Corona deep orange throughout, fluted, frilled, lobed, bowl shape. Neck grows badly though stiffly in water.
- GLENWHERRY** 3W-R
Perianth creamy white, stained yellow at base. Segments flat, pointed, broad, overlapping. Corona deep burnt orange red with greenish gold eye, fluted. Corona edge dies first. Neck grows badly in water.
- GLORIA** 1W-W
Perianth creamy white, slight green on back, a little creased. Segments flat, pointed, broad, overlapping. Corona opens creamy primrose, develops white. Neat.
- GLORIA MUNDI** 2Y-YOR
Perianth very pale lemon, creased. Segments flat, pointed, slightly overlapping. Corona pale orange shading to gold base. Poor looking flower.
- GLORIFICATION** 2W-YOO
Perianth creamy white, creased, notched, often buckled. Segments flat, pointed, overlapping. Corona orange gold base, shading to bright orange with gold edge, fluted, nearly flat.
- GLORIOSUS** 8W-Y
Discarded - Better suited to Isles of Scilly.
- GLORIOUS** 8W-O
Perianth white, stained a little yellow at base, slightly creased. Segments flat, rounded, overlapping. Corona deep burnt red, fluted. Style almost as long as longest stamens. Neck grows in water.
- GLORY OF LISSE** 9W-YYR
Perianth white, stained yellow at base, deeply creased, often buckled, star shape. Segments flat, pointed, barely overlapping. Corona greenish gold eye, shading to lemon with narrow rim of deep red, fluted, almost flat. Neck grows badly in water.
- GLORY OF SASSENHEIM** 1W-Y
Perianth cream, stained deep yellow at base, star shaped. Segments twisted, pointed, barely overlapping. Corona bright deep golden yellow.
- GLOWING PHOENIX** 4Y-O
Perianth segments cream, flat, pointed, separated, 2 complete rows. Corona segments bright orange in whorls with short twisted perianth segments intermixed.
- GLOWING RED** 4W-R
Perianth creamy white, stained base and back. Segments broad, round and 1/2 overlapping with slight buckling, and curled in at edges. Perianth segments intermixed with frilled corona segments. Corona segments shorter than perianth. Neck grows badly in water.
- GLYNVER** 3W-O
Perianth poeticus white, stained yellow base, often buckled, circular outline. Segments flat, rounded overlapping. Corona slight green eye changing to deep orange red, fluted, frilled, lobed. Corona edge dies 5 days before the perianth. Neck grows badly though stiffly in water.
- GOBLET** 1W
Perianth white. Segments broad and overlapping. Corona very large open mouthed, frilled and yellow.
- GODOLPHIN** 1Y-Y
Perianth clear soft sulphur, yellow, very smooth. Segments flat, pointed, broad, overlapping. Corona deep clear sulphur yellow, almost straight.
- GODREVVY** 1Y-Y
Perianth soft sulphur yellow. Segments twisted, pointed, broad, overlapping. Corona deep clear sulphur yellow, very broad. Badly proportioned.
- GOG** 1Y-Y
Perianth yellow. Segments broad and overlapping, twisted and convex. Corona large open mouthed, frilled and yellow. Neck grows in water.
- GOLD BULLION** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and yellow.
- GOLD COLLAR** 11
Perianth deep sulphur yellow. Segments flat, pointed, overlapping. Corona deep lemon yellow, split into six segments, three outer segments lie flat on perianth, three inner segments incurved, deeply frilled. Poor poise, top heavy.
- GOLD CROWN** 2W-Y
Discarded - Short.

- GOLD MEDAL** 1Y-Y
Perianth bright deep clear lemon, slightly creased. Segments flat, rounded, broad, well overlapping. Corona deep lemon yellow. Dumpy appearance.
- GOLD RIM** 2
Discarded - Short, flowers below foliage.
- GOLD RUSH** 11Y-YOY
Perianth white. Segments very broadly ovate and overlapping. Corona split, deep yellow on front, with white back, edges turning in, showing white collar. Neck grows badly in water.
- GOLDBEATER** 1Y-Y
Perianth deep clear maximus gold, green centre line on back. Segments twisted, pointed, barely overlapping. Corona deeper golden yellow, edge lobed.
- GOLDCOURT** 1Y-Y
Perianth deep rich lemon yellow, very smooth. Segments flat, pointed, broad, overlapping. Corona deep, almost orange gold. Good proportion and well balanced.
- GOLDEN ACE** 2Y-Y
Perianth yellow. Segments broad and overlapping. Corona funnel shaped, frilled, and same yellow as perianth.
- GOLDEN ADVANCE** 1Y-Y
Perianth deep lemon yellow, crêpey texture. Segments twisted, pointed, overlapping. Corona deeper intense lemon yellow, open mouthed. Dumpy appearance.
- GOLDEN AGE** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and same yellow as perianth.
- GOLDEN AMBER** 2Y-ORR
Perianth palest cream. Segments stained yellow at lower edge and base, smooth, curled in deeply at edges, rounded outline, overlapping at base. Corona orange/yellow throat developing to blotchy orange middle band developing to deep lemon edge. Flower substance very good.
- GOLDEN BRACELET** 2Y-O
Perianth primrose developing cream, creased, crêpey, sometimes caught in corona, circular outline. Segments flat, rounded, broad, overlapping. Corona orange gold, variable colour, fluted, lobed, bowl shape. Stiff.
- GOLDEN CASTLE** 4Y-O
Perianth segments clear yellow, flat, pointed, overlapping in almost 2 rows. Corona deeper golden yellow, trumpet shaped, almost complete, filled in tightly with segments. Very heavy and clumsy.
- GOLDEN CHEER** 2Y
Perianth yellow. Segments ovate and semi-overlapping, slightly twisted and slightly inflexed. Corona open mouthed, frilled and yellow. Neck grows badly in water.
- GOLDEN CLARION** 1Y-Y
Perianth yellow. Outer segments very broadly ovate, inner segments ovate. Corona open mouthed, frilled and same yellow as perianth. Neck grows in water.
- GOLDEN CLOUD** 1Y-Y
Perianth yellow. Segments as Golden Clarion. Corona flared cylinder, frilled and same colour as perianth.
- GOLDEN COCKEREL** 1Y-Y
Perianth bright lemon, slightly creased. Segments flat, pointed, broad, well overlapping, shovel shape. Corona deep lemon, crenate, flanged.
- GOLDEN CYCLE** 6Y-Y
Discarded - Short, cyclamineus hybrid.
- GOLDEN DAWN** 8Y-O
Approx. 6 florets per stem. Perianth sulphur. Segments slightly wavy edges, incurling. Corona orange, short, wavy edge. Neck grows badly in water, some with hollow stems.
- GOLDEN DAY** 1Y-Y
Perianth clear mid-lemon yellow, paler at edges, smooth, severely twisted. Segments rounded, overlapping. Corona deeper, bright yellow; long, slender; mouth wide, flared.
- GOLDEN DOLLAR** 1Y-Y
Perianth light lemon yellow, smooth. Segments flat, pointed, broad, well overlapping. Corona deeper lemon yellow, neat.
- GOLDEN DUCAT** 4Y-Y
Segments clear deep yellow, flat, pointed, broad, overlapping, in symmetrical rows of segments, a little more pointed in centre. Double camellia flower shape. Outer segments curl back with age.

GOLDEN EMPEROR 1Y-Y
Perianth soft sulphur yellow, some green at back. Segments twisted, pointed, overlapping. Corona lemon yellow, with neat edge.

GOLDEN EMPIRE 1Y-Y
Perianth gold, edges slightly incurling. Segments would fit into a double triangle, overlapping to half length of segments. Corona gold, flanged, deep slightly ragged frill. Neck grows in water, stem weak.

GOLDEN GOBLET 7Y-Y
Perianth deep bright lemon yellow, creased. Segments twisted, pointed, overlapping. Corona deep bright lemon yellow, squat, open mouthed.

GOLDEN GODDESS 1Y-Y
Discarded - Short.

GOLDEN HARBINGER 1Y-Y
Perianth deep clear yellow, slight green back. Segments flat, pointed, overlapping. Corona deep clear golden yellow.

GOLDEN HARVEST 1Y-Y
Perianth clear lemon, slightly green back. Segments slightly twisted, pointed, overlapping. Corona deeper lemon, deeply frilled, flanged. Neck grows badly in water.

GOLDEN HORN 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and same yellow as perianth. Neck grows in water.

GOLDEN INCENCE 7Y-Y
Discarded - Small, flowers below foliage, jonquilla hybrid.

GOLDEN LACQUER 6Y-Y
Perianth bright lemon yellow, creased, barely reflexed. Segments twisted, pointed, overlapping. Corona deep bright yellow, open mouthed. Not typical of cyclamineus hybrids.

GOLDEN MARVEL 1Y-Y
Perianth deep primrose yellow, slight green stripe on back. Segments flat, pointed, broad, overlapping. Corona deep clear yellow.

GOLDEN MELODY 1Y-Y
Perianth sulphur yellow. Segments twisted, pointed, overlapping. Corona pale lemon yellow, rather long in proportion to perianth.

GOLDEN PERFECTION 7Y-Y
Perianth pale lemon yellow, colour fades out from tips of segments with age, crêpey textures. Segments twisted, pointed, broad, overlapping. Corona bright deep lemon yellow. Neck grows in water.

GOLDEN RALLY 1Y-Y
No description available.

GOLDEN RAPTURE 1Y-Y
Perianth yellow. Segments ovate and overlapping. Corona flared funnel, frilled and same yellow as perianth. Neck grows in water.

GOLDEN RAY 1Y-Y
Discarded - Poor looking flower.

GOLDEN RIOT 1Y-Y
Perianth lemon yellow, a little creased, pointed outline. Segments flat, pointed, overlapping. Corona deeper lemon yellow, rather open mouthed.

GOLDEN ROBIN 1Y-Y
Perianth lemon, slightly hooded and edges incurling. Segments would fit into a double triangle. Corona deep lemon, flanged, deeply cut edge. Delicate flower. Weak stems.

GOLDEN SCEPTRE 7Y-Y
Perianth bright canary yellow, creased, with crinkled edges. Segments slightly twisted, pointed, overlapping. Corona deeper yellow, almost trumpet proportion.

GOLDEN SMILE 1Y-Y
Discarded - Flimsy flower.

GOLDEN SPRAY 2Y-Y
Discarded - Flowers below foliage.

GOLDEN SPUR 1Y-Y
Perianth deep yellow, some green on back. Segments twisted, pointed, separated, edges curl back. Corona deeper golden yellow, edge deeply lobed.

- GOLDEN STARLIGHT** 1Y-Y
Perianth lemon, smooth. Segments acute, overlapping at base. Corona deep lemon, long, flanged, gently lobed.
- GOLDEN TOP** 1Y-Y
Discarded - Short, rough flower.
- GOLDEN TORCH** 2Y-Y
Perianth soft clear lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona deep golden yellow, of trumpet proportions.
- GOLDEN TRIUMPHATOR** 2Y-Y
Perianth bright deep yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona bright golden yellow, deeply frilled, of almost trumpet proportions. Rough looking flower.
- GOLDEN VINTAGE** 2Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and deep yellow in colour. Neck grows in water.
- GOLDEN WEALTH** 1Y-Y
Perianth lemon yellow, a little creased, pointed outline. Segments flat, pointed, overlapping. Corona deeper lemon yellow, broad mouth, badly proportioned.
- GOLDFLAKE** 2Y-YYO
Discarded - Low flower production.
- GOLDILOCKS** 7Y-Y
Perianth clear lemon yellow, smooth, edges curl in a little. Segments flat, rounded, overlapping. Corona deeper lemon yellow with greenish eye. Neck grows in water.
- GOLDSBOROUGH** 1Y-Y
Perianth bright lemon, crêpey. Segments flat, pointed, broad, overlapping. Corona deep lemon, crenate, flanged, stiff.
- GOLYER** 1Y-Y
Perianth sulphur, slightly creased. Segments flat, pointed, overlapping. Corona mid lemon, crenate, flanged. Very stiff. Hangs head when young.
- GOORNONG** 1W-Y
Perianth white. Segments ovate, overlapping and inflexed. Corona flared funnel, frilled and yellow. Neck grows in water.
- GOVERT FLINK** 3W-GYO
Perianth creamy white, creased. Segments flat, rounded, well overlapping. Corona bright orange, paler at base with greenish-gold eye, creased.
- GOYESCAS** 3Y-YYO
Perianth sulphur yellow, paler at tips, a little creased. Segments flat, pointed, overlapping. Corona greenish lemon base with bright orange red rim for 1/3 its length, nearly flat, fluted.
- GRACE MOFFAT** 2Y-O
Perianth primrose yellow, a little paler at tips, crêpey texture, segments sometimes flop back. Segments flat, pointed, broad, overlapping. Corona deep burnt orange red, fluted. Neck grows a little in water.
- GRACKLE** 2Y-O
Perianth pale clear primrose yellow, creased. Segments flat, pointed, broad, overlapping. Corona bright orange red shading to orange at base, deeply frilled. Stiff and sturdy.
- GRANADA** 2Y-O
Perianth soft lemon yellow, a little creased, crêpey texture, edges curl back. Segments flat, slightly reflexed, pointed, overlapping. Corona orange gold, developing to deeper colour. Similar to Helios.
- GRAND GALA** 1Y-Y
Perianth yellow. Segments ovate and overlapping. Corona flared funnel, frilled and yellow. Neck grows in water.
- GRAND MONARQUE** 8W-Y
Discarded - Better suited to Isles of Scilly.
- GRAND NATIONAL** 2Y-R
Perianth pale lemon yellow with paler tips, creased, crêpey texture. Segments flat, pointed, overlapping. Corona bright orange, fluted.
- GRAND PRIMO CITRONIERE** 8W-Y
Discarded - Better suited to Isles of Scilly.
- GRAND SOLEIL D'OR** 8Y-O
Perianth bright lemon yellow, reflexed, slightly creased. Segments flat, pointed, overlapping. Corona opens orange, develops deep solid orange red, smooth goblet shape.

- GRANDIS** 1W-Y
Perianth creamy white, stained yellow at base, slightly creased and crêpey texture. Segments slightly twisted, pointed, overlapping. Corona bright lemon yellow, deeply lobed edge, wide mouth.
- GRANVELLE** 2W-O
Perianth creamy white, deeply creased, top three segments incurved. Segments flat, pointed, overlapping. Corona opens primrose, developing creamy apricot with narrow paler edge, fluted, uneven deeply frilled edge.
- GRAPE FRUIT** 1Y-Y
Perianth pale creamy sulphur yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona clear pale creamy sulphur yellow, ageing to apricot buff (in water).
- GRAYLING** 2W-YYW
Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona deep primrose yellow with thin cream edge.
- GREAT LEAP** 4Y-Y
Perianth yellow. Segments double, broad and overlapping. Corona double, deep yellow and intermixed with perianth segments. Scented. Neck grows in water.
- GREAT WARLEY** 2W-Y
Perianth creamy white stained yellow at base, star shaped. Segments slightly twisted, pointed, barely overlapping. Corona bright deep lemon yellow, open mouthed.
- GREEN HOWARD** 3W-GYY
Perianth opens greenish primrose, develops greenish cream, stained yellow at base, slightly creased, circular outline. Segments flat, rounded, broad, overlapping. Corona pale greenish lemon with slightly deeper rim and moss green eye, fluted, shallow. Neck grows stiffly in water.
- GREEN ICE** 2W-GWW
Perianth white. Segments ovate and overlapping with central ridge. Corona cup shaped, frilled, green eyed and white.
- GREEN ISLAND** 2W-GWY
Perianth creamy white, slightly creased, edges curl in a little. Segments flat, rounded, well overlapping. Corona grey green centre for 1/4 of its length, shading to band of greyish primrose for 1/2 length in centre, shading to edge of greenish lemon for 1/4 of its length.
- GREEN MINT** 3W-W
Discarded - Small.
- GREENCASTLE** 3W-GYO
Perianth creamy white, stained yellow base, edges curl in a little, slightly reflexed, pointed outline. Segments flat, pointed, overlapping. Corona deep moss green eye changing to narrow gold band, shading to bright orange edge for 1/2 its length, fluted, frilled, unevenly lobed, almost flat. Neck grows stiffly in water.
- GREENLAND** 2W-W
Discarded - Flowers below foliage.
- GREENMOUNT** 3W-Y
Perianth white. Segments broad, overlapping, turned over at edges. Corona small cup, frilled, white with yellow rim. Neck grows in water.
- GREENORE** 2W-WWY
Perianth greenish cream, slightly creased, edges curl in. Segments flat, pointed, broad, overlapping. Corona greyish cream shading to creamy apricot edge for 1/3 its length, grey green eye, deep bowl shape, stiff.
- GREETING** 2W-Y
Perianth white with slight yellow base, a little creased. Segments flat, pointed, overlapping. Corona clear pale lemon with paler edge, small almost straight sided.
- GREGALACH** 1W-Y
Discarded - Rough flower.
- GRENOBLE** 2Y-Y
Perianth deep lemon, creased. Segments slightly notched, overlapping. Corona gold, flat to open mouthed, slightly frilled at edge. Neat well balanced but small flower. Neck grows badly in water.
- GRETA GARBO** 2Y-O
Discarded - Flowers below foliage, long neck, poor poise.
- GREY LADY** 3W-WWP
Perianth poeticus white, slightly creased, circular outline. Segments flat, rounded, broad, well overlapping. Corona grey green eye shading to band of pale grey, changing to a band of bright orange red for 1/3 its length, slightly fluted, flat. Very prominent style. Neck grows badly in water.

- GRIANAN** 1W-Y
Perianth milky white, stained yellow base and centre line back. Segments pointed, overlapping, slightly creased, curled in at edges. Corona sulphur yellow, paler at base, slender, attractively belled, crenate edge.
- GRULLEMANS' GIANT** 2W-YYO
Discarded - Short, flowers below foliage.
- GUARDIAN** 2W-Y
Perianth creamy white, double triangle, edges curl in a little. Segments flat, pointed, well overlapping. Corona pale clear lemon, develops deep cream with neat rolled back edge, of almost trumpet proportions. Neat and stiff.
- GUIDING STAR** 2W-O
Perianth white. Segments ovate and overlapping. Corona large cup, frilled and orange. Neck grows in water.
- GULLIVER** 3Y-YYO
Perianth soft sulphur yellow, a little paler at the tips. Segments flat, pointed, broad, overlapping. Corona greenish gold shading to bright orange red for 2/3 of its length, fluted. Stiff and sturdy.
- GUSTAV MAHLER** 2W-Y
Discarded - Floppy, rough flower.
- GUVEZNE** 2W-O
Perianth creamy white, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona greenish gold eye shading to bright orange red, with narrow rim of pale apricot, some white blotches on edge, fluted. Neck grows badly in water.
- GWENTER** 2W-W
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona opens creamy primrose, develops to creamy white, of trumpet proportion.
- GWILLS** 3W-GWO
Perianth cream, often with thickened knob of deeper cream on tips, creased, reflexed. Segments flat, rounded, well overlapping. Corona lemon yellow, shading to delicate apricot orange edge for 1/4 of its length and large moss green eye, fluted, flat. Very long style.
- GYRFALCON** 2W-Y
Perianth creamy white, a little creased. Segments flat, pointed, broad, overlapping. Corona creamy primrose, with thick uneven frilling and white blotches on edge. Very heavy and ugly.
- HADES** 2Y-O
Perianth opens primrose yellow, stained yellow at base, develops cream. Segments flat, pointed, overlapping. Corona deep burnt orange red. Neck grows in water.
- HAKA** 2Y
No description available.
- HALF MOON** 5Y-Y
Discarded - Short, small, triandrus hybrid.
- HALSKRAAG** 11W-W
Perianth creamy white, creased, pointed outline. Segments flat, pointed, separated. Corona creamy white, split unevenly into six segments, lower three segments lie flat on perianth, inner three segments curl inwards. Very long style. Neck grows badly in water.
- HALVOSE** 8Y-R
Perianth primrose yellow, suffused apricot. Segments flat, rounded, well overlapping. Corona bright orange red, fluted.
- HALWYN** 1Y-Y
Perianth soft lemon yellow, slightly creased, stiff. Segments slightly twisted, pointed, broad, overlapping. Corona bright lemon yellow.
- HAMZALI** 3W-YYR
Perianth creamy white, creased, edges curl in, often buckled, lower segments reflex, circular outline. Segments flat, rounded, well overlapping. Corona grey eye with narrow primrose band changing to cream with wire rim of apricot orange, fluted, lightly frilled, lobed, almost flat. Neck grows badly in water.
- HAN VAN MEERGEREN** 2Y-R
Perianth primrose yellow, creased, crêpey texture. Segments flat, pointed, broad, overlapping. Corona gold, shading to orange red for 1/2 its length.
- HANDCROSS** 2Y-Y
Perianth yellow. Segments broad, overlapping, with ridged texture. Corona open mouthed, frilled and same yellow as perianth. Neck grows in water.
- HANNIBAL** 11Y-O
Perianth pale yellow. Segments very broadly ovate and overlapping. Corona split, frilled, coloured orange and yellow. Neck grows in water.

- HARBINGER OF SPRING** 1Y-Y
Perianth pale yellow. Segments rounded, overlapping. Corona rich gold, bold, wide flared mouth, deeply crenate edge. Neck grows in water.
- HARDY** 3W-O
Perianth creamy white stained slight yellow at base, creased, badly buckled. Segments flat, pointed, overlapping. Corona bright orange with paler orange base, double frilled.
- HARRIER** 2Y-O
Perianth lemon, creased, slightly crêpey, pointed outline. Segments flat, pointed, overlapping. Corona bright orange paler at base, fluted, lightly frilled, lobed, bowl shape. Neck grows stiffly in water.
- HARROGATE** 2Y-YYR
Discarded - Small, flowers below foliage.
- HARTSDOWN** 1W-W
No description available.
- HARVEST MOON** 5Y-Y
Perianth sulphur yellow, a little creased, pointed outline. Segments flat, pointed, barely overlapping. Corona sulphur yellow, broad mouth, of trumpet proportion, neat edge. Very long style. Neck grows badly in water.
- HARVESTER** 1Y-Y
Perianth soft sulphur yellow, greenish back, inclined to be hooded. Segments twisted, pointed, overlapping. Corona lemon yellow, flanged.
- HATHOR** 7Y-Y
Discarded - Small, flowers below foliage, jonquilla hybrid.
- HAVELOCK** 2Y-Y
Perianth deep primrose yellow, one segment often buckled, crêpey texture. Segments flat, pointed, overlapping. Corona golden yellow.
- HAVEMEIJER** 3W
Perianth white. Segments rounded and overlapping. Corona small cup, frilled, yellow with orange rim. Neck grows badly in water.
- HAWAII** 4Y-O
Perianth yellow. Double segments very broadly ovate and overlapping. Corona double, orange and intermixed with inner perianth segments. Neck grows badly in water.
- HAYLE** 2Y/2Y-YOO
Perianth white. Segments round with slight overlap. Corona cup shaped, frilled and orange. Neck grows in water.
- HAZEL BRILLIANT** 2Y-O
Perianth pale lemon yellow. Segments broad, overlapping, and crêpey. Corona bowl shaped, frilled, orange with small whitish patches on edge.
- HAZELWOOD** 3W-WWY
Discarded - Floppy, flowers below foliage.
- HEART'S DELIGHT** 2Y-Y
Perianth primrose yellow, with paler tips, crêpey texture. Segments flat, rounded, overlapping. Corona deep old gold, wide mouth.
- HEAVEN** 2W-OOY
Discarded - Low flower yield.
- HELEN F.OHMS** 2Y-W
Perianth light lemon yellow, paler at tips, very crêpey texture, creased. Segments flat, pointed, overlapping. Corona orange red, paler at base, fluted. Neck grows badly in water.
- HELFDOR** 1Y-Y
Perianth bright clear gold. Segments slightly twisted, pointed, separated. Corona deep bright gold, almost straight with rolled back edge.
- HELIOS (syn. ABUNDANCE)** 2Y-O
Perianth primrose yellow, edges curled in. Segments flat, rounded, overlapping. Corona deep yellow, edged pale orange with moss green eye, fluted.
- HELIOS** 8Y-Y
Perianth soft lemon yellow, edges curled back, crêpey texture. Segments flat, slightly reflexed, pointed, overlapping. Corona orange gold, developing to deeper orange colour.
- HENRY DETERDING** 1W-Y
Perianth cream, stained deep yellow at base, crêpey texture, stiff. Segments twisted, pointed, broad, overlapping. Corona very deep bright yellow.

- HENRY IRVING** 1Y-Y
Perianth bright clear yellow, a little paler at base. Segments twisted, pointed, overlapping. Corona deeper bright yellow.
- HERBERT BARR** 1Y-Y
Perianth sulphur yellow, some green on back. Segments twisted, pointed, overlapping. Corona lemon yellow.
- HESLA** 7Y-Y
Perianth sulphur yellow, a little deeper at base, smooth, inclined to go paler with age. Segments flat, pointed, overlapping. Corona lemon yellow, crinkled.
- HEXAGON** 2W-W
Perianth creamy white, creased, curled in at edges. Segments flat, pointed, overlapping. Corona opens primrose yellow, develops pale buff then turns pale buff primrose. Similar to Riva.
- HEXAMETER** 9W-GYR
Perianth pure white, smooth. Segments flat, rounded, well overlapping. Corona greenish lemon changing to deep red edge for 1/5 of its length, with a moss green eye, flat, fluted. Style a little longer than longest stamens. Neck grows in water.
- HI KING** 1Y-Y
No description available.
- HIAWASSEE** 8W-W
Perianth white. Segments acute, very little overlap. Corona small cup, green eye and white segments.
- HIGH LIFE** 2W-OOY
Perianth creamy white, creased, a little floppy. Segments flat, rounded, well overlapping. Corona bright apricot orange with thin, paler edge and deeper eye, edge scalloped and frilled. Neck grows in water.
- HIGHFIELD BEAUTY** 8Y-GYO
1-3 florets per stem. Perianth primrose, slightly reflexed. Segments rounded, very well overlapping. Corona orange, darker at rim, cup shaped, wavy edge. Neck grows badly.
- HIGHLAND CASTLE** 1W-Y
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and yellow.
- HILLMOUNT** 2W-R
Perianth opens primrose, develops creamy white stained yellow at base, creased, edges curl in. Segments flat, pointed, overlapping. Corona deep orange red, a little paler at base, fluted. Corona edge dies first. Neck grows badly in water.
- HILLSBOROUGH** 1W-Y
Perianth creamy white, creased, sometimes caught in corona. Segments flat, pointed, broad, well overlapping, shovel shape segments. Corona cream. Hangs head badly.
- HILLSTON** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and same yellow as perianth. Neck grows in water.
- HIMALAYA** 1W-W
Perianth creamy white, slightly creased. Segments slightly twisted, pointed, broad, well overlapping, shovel shape. Corona cream, rather broad mouthed, well proportioned.
- HINDENBURG** 1Y-Y
Perianth bright gold. Segments slightly twisted, pointed, overlapping. Corona deeper gold, wide open-mouth, deeply frilled, dumpy appearance.
- HINDUSTAN** 1W-W
Perianth creamy white, a little creased. Segments flat, pointed, broad, well overlapping. Corona cream. Stiff.
- HIS EXCELLENCY** 1Y-Y
Discarded - Poor quality, susceptible to basal rot.
- HOLLAND** 2Y-O
Perianth deep bright lemon yellow, smooth, one segment sometimes buckled. Segments flat, rounded, well overlapping. Corona bright orange red. Stiff.
- HOLLANDIA** 4Y-O
Perianth segments lemon yellow, flat, pointed, barely overlapping, one row complete. Corona orange, complete, segments tipped yellow in centre. Variations from completely single flowers to those with corona filled in, stamens often present, ugly.
- HOLLY BERRY** 2Y-R
Perianth bright lemon, paler at tips, slightly creased, circular outline. Segments flat, pointed, well overlapping. Corona bright burnt orange paler at base, fluted, frilled, bowl shape. Very stiff.

- HOLLYWOOD** 2Y-O
Perianth pale lemon, developing streaky primrose, creased. Segments flat, pointed, overlapping. Corona light orange, paler at base, fluted, frilled, lobed, bowl shape. Neck grows badly in water.
- HOME FIRES** 2Y-O
Perianth clear lemon, creased, sometimes notched, pointed outline. Segments flat, pointed, overlapping. Corona deep orange red paler at base, fluted, lightly frilled, lobed. Neck grows stiffly in water.
- HOMER** 9W-OOR
Perianth white, deeply creased. Segments flat, pointed, barely overlapping. Corona greenish gold eye for ½ its length, shading to greenish lemon with narrow rim of dark orange red, fluted, uneven edge. Neck grows in water.
- HONEY GOLD** 2Y-Y
Perianth yellow. Segments broad and overlapping. Corona large open mouthed, frilled and same yellow as perianth. Neck grows in water.
- HONEYBIRD** 1Y-W
Perianth yellow. Segments ovate and overlapping. Corona flared funnel, frilled and darker yellow. Neck grows in water.
- HONG KONG** 2Y-O
Perianth light lemon yellow, creased, one segment sometimes buckled. Segments flat, pointed, overlapping. Corona orange gold shading gradually to bright orange red edge, fluted, deep bowl shape.
- HOPESAY** 2W-YYO
Perianth creamy white, slightly creased, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona greenish lemon changing abruptly to bright orange for 1/3 its length, fluted. Neck grows badly in water.
- HORACE** 9W-GOR
Perianth creamy white, creased down centre of segments with edges curled back. Segments twisted, pointed barely overlapping. Corona orange red a little deeper at edge, small greenish gold eye, flat, deeply fluted. Style as long as longest stamens.
- HORIZON** 1Y-Y
No description available.
- HORN OF PLENTY** 5W-W
Discarded - Short, small, triandrus hybrid.
- HORSFIELDII** 1W-Y
Perianth creamy white, stained yellow at base. Segments flat, pointed, barely overlapping. Corona deep lemon yellow. Very long style.
- HOSPODAR** 2Y-O
Perianth primrose yellow, creased, star shaped. Segments flat, pointed, overlapping. Corona orange gold, variable colour, deepens with age, fluted. Neck grows in water.
- HUNTERLEA** 3W-WWR
Perianth white. Segments broad and overlapping. Corona small cup, frilled, yellow with orange rim. Neck grows badly in water.
- HUNTER'S MOON** 1Y-Y
Perianth pale sulphury cream, slightly creased. Segments slightly twisted, pointed, overlapping. Corona clear sulphur yellow, ageing to paler colour with thin lemon rim.
- HUON** 9W-YYO
Perianth pure white stained yellow at base, slightly reflexed, circular outline. Segments flat, rounded, well overlapping. Corona orange changing to orange red edge for 1/4 of its length and slightly greenish eye, almost flat, fluted. Style as long as longest stamens.
- HUSKY** 3Y-YOR
Perianth pale creamy yellow a little deeper at base, creased, spidery looking. Segments twisted, pointed, separated. Corona bright orange red, shading to gold base.
- HUSSAR** 2Y-YYO
Discarded - Small.
- HYPERION** 2Y-Y
Perianth yellow. Segments broad and overlapping. Corona frilled, bowl shaped and yellow. Neck grows in water.
- H. CAMP** 3O-O
Perianth creamy orange. Segments ovate and overlapping. Corona small cup, frilled and orange. Neck grows badly in water.
- ICE CRYSTAL** 11W-W
Discarded - Short, flowers below foliage.

ICE FOLLIES 2W-W
Perianth creamy white, creased, a little floppy. Segments flat, pointed, overlapping. Corona opens creamy lemon, develops paler with cream rim, ageing cream throughout, deeply fluted, deeply frilled, lobed, almost flat. Neck grows stiffly in water.

IDEAL 8W-O
Perianth creamy white, creased and buckled. Segments flat, rounded, overlapping. Corona golden orange, fluted. Style reaches half way between short and long stamens.

ILLUMINATE 2Y-O
Perianth yellow. Segments ovate, overlapping, and ridged texture. Corona cup shaped, frilled and orange. Neck grows badly in water.

IMOGEN 2W-P
Discarded - Short, poor looking flower, variable colour.

IMPERATOR 1W-W
Perianth creamy white, creased. Segments slightly twisted, pointed, overlapping. Corona creamy primrose, develops to pale cream, open mouthed.

IMPROMPTU 2W-GYY
Discarded - Coarse flower, susceptible to basal rot.

INA 3W-O
Perianth cream, paler at tips, deeper at base, creased. Segments flat, rounded, well overlapping. Corona bright burnt orange, sometimes with white blotches on edge, fluted. Neck grows in water.

INCA GOLD 1Y-Y
No description available.

INDIAN CHIEF 4Y-O
Perianth segments deep primrose, outer segments reflexed, flat, pointed, separated, in 4 or 5 rows with pointed, twisted segments in centre all intermixed with corona segments of pale orange. Flower very full and heavy.

INDIAN SUMMER 2Y-O
Perianth deep clear lemon yellow, edges inclined in curl in. Segments flat, pointed, broad, overlapping. Corona bright orange-red, fluted. Neck grows in water.

INDOLINE 1Y-Y
Perianth light lemon, slightly creased. Segments slightly twisted, pointed, broad, overlapping. Corona deeper lemon, lobed, barely full length trumpet.

INDONESIA 2Y-O
Perianth creamy primrose, paler at tips, deeply creased but very stiff. Segments flat, rounded, well overlapping. Corona bright orange red sometimes with white blotches on edge.

INFATUATION 2W-YYP
Perianth white. Segments very broadly ovate and overlapping. Corona cup shaped, frilled, yellow with pink rim.

INFERNO 2Y-O
Perianth yellow. Segments ovate and overlapping. Corona cup shaped, frilled and orange.

INFLAMMABLE 2Y-O
Perianth yellow. Segments broad and overlapping. Corona straight cylinder, frilled and orange. Neck grows in water.

INGLESCOMBE 4Y-Y
Perianth segments clear, pale lemon yellow, flat, pointed, overlapping in 3 or 4 complete rows with smaller twisted segments fairly tight in centre, almost no corona segments, floppy. Neck grows in water.

INGRID VALBERG 11Y-O
Discarded - Floppy, flowers below foliage.

INISHKEEN 1W-Y
Discarded - Short, flowers below foliage.

INNISFALLEN 2W-Y
Perianth creamy white, smooth. Segments flat, pointed, broad, well overlapping. Corona bright lemon yellow, tinged orange. Stiff and sturdy.

INSULINDE 4W-O
Perianth pale primrose yellow, flat, pointed, overlapping, outer segments reflexed, one complete row and up to 6 pointed, narrow twisted segments in centre. Corona segments bright orange, frilled, intermixed in centre. Spidery, ragged look.

INTERIM	2W-YYP
Perianth greenish creamy white, edges curl in, slightly reflexed, pointed outline. Segments flat, pointed, overlapping. Corona pale cream shading to deep coppery peach edge for 1/3 its length, fluted, almost straight sided. Neck grows a little in water.	
INTERLUDE	2W-P
Discarded - Short, hooded.	
INVER	1Y-Y
Discarded - Small, hangs head.	
INVERELL	2W-W
Perianth white. Segments ovate and overlapping. Corona open mouthed, frilled and pale pink.	
INVERGORDON	2Y-O
Perianth pale primrose, creased, edges curled in, crêpey texture. Segments flat, pointed, broad, overlapping. Corona bright orange, red, shading to slight gold base, fluted.	
INVERNESS	2Y-YOO
Perianth yellow. Segments broad and overlapping. Corona frilled cup, orange with a yellow centre. Neck grows badly in water.	
INVESTMENT	1Y-Y
Perianth lemon, slightly twisted. Segments small and overlapping at base. Corona deep lemon, flanged, deeply cut rim. Weak stems. Neck grows in water.	
IRANI	4W-Y
Perianth creamy white, fairly smooth. Perianth broad, rounded, overlapping, curled in at edges; one complete row with six or more buckled segments intermixed with creamy yellow pale-edged corona segments in compact centre; frilled edge. Weak stems. Neck grows in water.	
IRENE COPELAND	4W-Y
Perianth segments creamy white, often opening greenish white, flat pointed, separated. Many narrow perianth segments intermixed evenly with primrose coloured corona segments throughout. Flat head, stiff.	
IRISH CHARM	2W-O
Discarded - Small, flowers below foliage.	
IRISH LEGEND	2W-O
Perianth creamy white, stained yellow base. Segments rounded, well overlapping. Corona rich deep orange with slight reddish tinge, expanded, bowl-shaped, frilled edge. Flower downwards facing. Neck grows badly in water.	
IRISH LUCK	1Y-Y
Perianth bright deep lemon yellow. Segments, flat, pointed, broad, overlapping. Corona deeper yellow. Neck grows in water.	
IRISH MINSTREL	2W-Y
Perianth white. Outer segments very broadly ovate, inner segments ovate and notched, and overlapping. Corona frilled cup, yellow in colour. Neck grows in water.	
IRISH PRINCE	1Y-Y
Perianth yellow. Segments ovate, slightly twisted with very slight overlap. Corona large wide mouthed, frilled and same yellow as perianth. Neck grows badly in water.	
IRISH RANGER	3W-O
No description available.	
IRISH ROSE	2W-P
Discarded - Short.	
IRISH SPLENDOUR	3W-R
Perianth white. Segments ovate, overlapping and slightly inflexed. Corona cup shaped, frilled and deep orange. Neck grows badly in water.	
IRMELIN	8W-R
Perianth creamy white, deeply creased and buckled, pointed outline. Segments twisted, pointed, barely overlapping. Corona bright orange red with moss green eye, fluted, almost flat. Style as long as longest stamens.	
ISLANDREAGH	2W-Y
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and yellow in colour.	
ISOBELINE	2W-P
Perianth white. Segments broad and overlapping with ridged texture. Corona open mouthed, frilled and pink. Neck grows in water.	

ISOLA 3W-YYO
 Perianth opens primrose, develops creamy white, stained yellow at base, edges curl in, creased, often buckled. Segments flat, pointed, broad, overlapping. Corona gold changing to deep orange for 1/2 its length, fluted, almost flat. Corona edge dies first. Neck grows badly in water.

JACINTA MARTO 2W-O
 Perianth white, creased and a little floppy. Segments flat, rounded, well overlapping. Corona bright apricot orange, extra frilled edge.

JACK SNIPE 6W-Y
 Perianth opens cream, develops white stained slight yellow at base, reflexed. Segments flat, pointed, overlapping. Corona bright yellow with hint of orange, almost straight sided. Neat and wiry.

JACKPOT 2Y-R
 Perianth sulphur, slightly hooded, crepey, edges incurling. Segments ovate, overlapping. Corona deep orange, short, open mouthed, crenate, edge lightly frilled. Neck grows stiffly in water.

JAGUAR 2Y-O
 Perianth lemon, tinged slight orange base, crêpey, circular outline. Segments flat, rounded, broad, overlapping. Corona bright burnt orange, fluted, barely frilled, deeply cut lobed edge, shallow bowl shape. Neck grows stiffly in water.

JALNA 2Y-R
 Perianth soft lemon yellow, creased, odd segments caught in corona or buckled. Segments flat, rounded, overlapping. Corona bright orange red. Neck grows in water.

JAMAICA 3W-O
 Perianth creamy white, stained yellow base, creased, sometimes notched, edges curl in a little. Segments flat, pointed, overlapping. Corona deep bright orange fluted, frilled, lobed, bowl shape. Neck grows stiffly in water.

JANA 6Y-Y
 Discarded - Short, small, cyclamineus hybrid.

JANUS 2Y-Y
 Perianth yellow. Segments broad and overlapping with ridged texture. Corona frilled and yellow. Neck grows badly in water.

JANZ 2Y-R
 No description available.

JEANNE DESOR 2W-Y
 Discarded - Flowers below foliage, untidy flower.

JENNIFER RUTH 1W-W
 Perianth white, creased, pointed outline. Segments flat, pointed, overlapping. Corona creamy white.

JENNY 6W-W
 Discarded - Short, cyclamineus hybrid.

JERPOINT 2Y-YYO
 Perianth lemon yellow, creased, edges curled in. Segments flat, pointed, broad, overlapping. Corona gold shading to bright orange red for 1/2 its length, fluted, straight sided. Stiff and erect.

JERSEY BEAUTY 2Y-O
 Discarded - Flowers below foliage, long neck, poor poise.

JERSEY CREAM 1W-Y
 Perianth creamy white. Segments flat, pointed, overlapping. Corona buff primrose, open mouthed, barely trumpet length.

JESSONDA 2W-YYP
 Perianth white, slightly creased. Segments flat, pointed, broad, well overlapping. Corona apricot primrose with peach pink edge, developing to deep buff primrose with cream edge. Stiff and sturdy.

JEST 2Y-Y
 Discarded - Short, rough flower.

JEZEBEL 3Y-R
 Discarded - Small.

JILL 2W-YYO
 Perianth creamy white, creased, circular outline. Segments flat, rounded, overlapping. Corona pale greenish lemon with narrow edge of orange, fluted. Neck grows in water.

JINDALEE 2W-Y
 Perianth cream, creased, edges incurling, and slightly hooded. Segments triangular, overlapping. Corona clotted cream, open mouthed with frilled edge.

- JINGELLIC 2W-P
No description available.
- JOBI 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona funnel shaped, frilled and yellow in colour.
- JOHANNESBURG 2W-O
Perianth creamy white, deeply creased, buckled, often caught in corona. Segments flat, pointed, well overlapping. Corona bright orange edged pale apricot, thick frilled edge in three or four rows. Rough. Neck grows in water.
- JOHN EVELYN 2W-O
Perianth creamy white, creased. Segments flat, rounded, well overlapping. Corona opens lemon yellow quickly developing to apricot, extra frilled edge.
- JOHN GREENWAY 2W-WWY
Perianth white. Outer segments very broadly ovate, inner segments ovate and overlapping. Corona open mouthed, frilled and yellow. Neck grows in water. Soft hollow stems.
- JOHN OF SALISBURY 2W-Y
Perianth white. Segments very broadly ovate and overlapping. Corona flared cup, frilled and white in colour. Neck grows badly in water.
- JOKE FULMER 2W-YOO
Perianth poeticus white, deeply creased, edges curl in a little, circular outline. Segments flat, rounded, broad, well overlapping. Corona bright apricot orange, shade paler at edge, fluted, deeply lobed, double frilled. Stiff.
- JOLI-COEUR 2W-YYO
Discarded - Floppy, untidy flower.
- JOSEPH MACLEOD 1Y-Y
Perianth sulphur, slightly creased, crêpey. Segments slightly twisted, pointed, overlapping. Corona clear mid lemon, frilled, crenate. Neck grows in water.
- JOYANT 2W-O
No description available.
- JOYOUS 2W-Y
Perianth white. Outer segments very broadly ovate, inner whorl ovate. Corona funnel shaped, flared, frilled and lemon colour. Neck grows in water.
- JUANITA 2Y-O
Perianth lemon yellow, deeply creased, top three segments often buckled and twisted. Segments slightly twisted, pointed, overlapping. Corona orange red shading paler at base, white blotches on edge, fluted, bowl shape. Neck grows in water. Rough.
- JUBILANT 2Y-Y
Perianth bright lemon yellow, double triangle, hooded. Segments flat, rounded, overlapping. Corona deeper bright lemon yellow.
- JUBILIATION 2W-Y
Perianth white. Segments broad and overlapping. Corona bowl shaped, frilled and yellow. Neck grows badly in water.
- JULES VERNE 2W-Y
Perianth creamy white stained pale yellow at base, and green on back, edges notched and often caught in corona. Segments twisted, pointed, broad, overlapping. Corona bright lemon yellow, developing a little paler. Coarse, rough flower.
- JULIET 9W-YYR
Perianth poeticus white, stained slight yellow base, a little creased, slightly reflexed. Segments flat, rounded, overlapping. Corona yellow green base 2/5 changing to greenish primrose 2/5, changing to deep red edge, fluted, lightly frilled, flat. Style slightly longer than longest stamens, neck grows stiffly in water.
- JULIUS CAESAR 1Y-Y
Discarded - Poor, ugly flower.
- JUNCO 1W-W
Perianth white. Segments very broadly ovate and overlapping. Corona large open mouthed, frilled and same white as perianth. Neck grows in water.
- JUNE ALLYSON 2W
Perianth opens pale primrose, quickly developing creamy white, stained cream at base, slightly creased, edges curl in. Segments flat, rounded, well overlapping. Corona bright orange red, a little paler at base, fluted, almost flat. Neat and stiff.

- JUVENTUS** 1W-Y
Perianth lemon yellow. Segments broad and overlapping. Corona large open mouthed, frilled and canary yellow. Neck grows in water.
- KALGOORLIE** 1Y-Y
Perianth clear pale lemon yellow, a little creased. Segments flat, pointed, broad, overlapping. Corona deeper lemon yellow, ageing to paler colour with lemon rim. Barely trumpet length.
- KANCHENJUNGA** 1W-W
Perianth creamy white, crêpey texture, perfect double triangle, inclined to be hooded. Segments flat, pointed, very broad, well overlapping. Corona creamy white, very wide mouthed.
- KANGA** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona flared, frilled funnel, same yellow as perianth.
- KANSAS** 3W-GYO
Perianth creamy white, smooth. Segments flat, pointed, broad, overlapping. Corona greenish primrose changing to deep apricot edge for 1/3 its length, with greenish grey eye and sometimes white blotches on the edge. Neck grows in water.
- KARACHI** 2Y-YYR
Perianth lemon yellow, slightly creased. Segments flat, pointed, well overlapping. Corona gold base changing to deep orange red edge for 1/3 its length, almost straight sided. Stiff and neat. Neck grows a little in water.
- KARAMUDLI** 1W-Y
Perianth white. Segments very broadly ovate and overlapping. Corona open mouthed, frilled and lemon yellow. Neck grows in water. Weak stems.
- KARANJA** 1W-P
Discarded - Small.
- KARELIA** 1W-Y
Discarded - Poor colour and shape.
- KATHLEEN FERRIER** 2W-Y
Perianth white. Segments broad and overlapping. Corona bowl shaped, frilled and orange in colour. Neck grows in water.
- KEN FARMER** 2Y-YYR
Perianth pale lemon, creased. Segments curled in at edges, ovoid, well rounded, overlapping. Corona gold lower half developing to orange upper, paler at edges; expanded, lobed edge. Weak stems.
- KENBANE** 1W-W
Perianth creamy white, creased. Segments slightly twisted, pointed, well overlapping. Corona opens deep primrose yellow, develops to deep cream, inside base of corona yellow, large open-mouth, badly balanced.
- KENMARE** 2W-P
Perianth creamy white, edges curl in, slightly creased. Segments flat, pointed, overlapping. Corona opens pale lemon, developing creamy peach, paler at base, variable colour.
- KERRY PIPER** 1Y-Y
Perianth bright lemon yellow, crêpey texture. Segments flat, pointed, broad, well overlapping, shovel shape. Corona deeper lemon yellow. Neat.
- KESTREL** 9W-GOR
Perianth white, creased, edges curl in a little. Segments flat, pointed, barely overlapping. Corona deep orange changing to deep red edge for 1/3 of its length with golden-green eye. Style as long as longest stamens.
- KHARTUM** 3W-R
Perianth cream, a little deeper at the base, odd segments sometimes creased. Segments flat, rounded, overlapping. Corona deep sealing wax red, deeply fluted. Stiff and sturdy.
- KIBO** 2W-W
Perianth creamy white, creased, crêpey, top segments incurve a little, pointed outline. Segments flat, pointed, overlapping. Corona cream, frilled, crenate, flanged, broad mouth. Neck grows in water.
- KILBRIDE** 2Y-Y
Perianth pale greenish lemon, pointed outline. Segments flat, pointed, overlapping. Corona deeper greenish lemon, fluted, bowl shape. Stiff and neat.
- KILCOOL** 2W-YYO
Perianth creamy white. Segments very broadly ovate and overlapping. Corona bowl shaped, frilled and orange in colour. Neck grows badly in water.
- KILCORAN** 2W-YYO
Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona pale apricot, shading to apricot orange edge for 1/2 its length, fluted, shallow bowl shape. Neck grows in water.

- KILDRUM** 3W-R
Perianth creamy white, stained yellow base, creased, top 3 segments incurve, pointed outline. Segments flat, pointed, overlapping. Corona bright orange red shading deeper at edge, fluted, frilled. Poor poise. Neck grows very badly in water.
- KILIMANJARO** 2W-YOO
Discarded - Flowers below foliage, floppy, susceptible to basal rot.
- KILKENNY** 1Y-Y
Perianth sulphur yellow. Segments slightly twisted, pointed, broad, well overlapping, shovel shape. Corona lemon yellow. Neat.
- KILLALA** 2W-YYO
Perianth creamy white, slightly creased, edges curl in. Segments flat, pointed, broad, overlapping. Corona light lemon yellow, changing to deep orange edge for 1/4 its length, of trumpet proportion.
- KILLALOE** 2W-W
Perianth creamy white, creased, notched. Segments flat, pointed, well overlapping. Corona cream, broad mouth.
- KILLARA** 8W-Y
Perianth white. Segments rounded with slight overlap. Corona small wavy cup, lobed and yellow in colour. Neck grows badly in water.
- KILLIFRETH** 3W
Perianth opens primrose, develops creamy white stained yellow base, slightly creased, notched, edges curl in, inclined to reflex. Segments flat, rounded, overlapping. Corona bright orange throughout, fluted, frilled, lobed, nearly flat, long prominent stigma. Neck grows in water.
- KILLIGREW** 2Y-O
Perianth pale primrose yellow, deeper at base, fading with age. Segments flat, pointed, overlapping. Corona bright orange with slightly paler edge. Neck grows in water.
- KILLYMOON** 2W-W
No description available.
- KILLYNURE** 1W-Y
Discarded - Flowers below foliage, hangs head badly.
- KILMORACK** 2Y-O
Perianth bright lemon, stained slight orange at base, slightly crêpey, very flat. Segments flat, rounded, broad, well overlapping. Corona deep burnt orange red with paler edge, slightly frilled, almost straight sided. Neck grows stiffly in water.
- KILMORE** 3W-O
Perianth white with yellow stain at base, smooth, slightly reflexed. Segments acute, edges incurving, overlapping. Corona bright orange, short, straight sided, slightly ribbed. Neck grows stiffly in water.
- KILMORNA** 2W-Y
Perianth creamy white, one segment deeply creased and buckled. Segments flat, pointed, broad, well overlapping. Corona opens primrose yellow, develops to deep cream, rolled back edge.
- KILREA** 2W-W
Perianth creamy white with slight green base, a little creased and crêpey, pointed outline. Segments flat, pointed, overlapping. Corona opens primrose, develops pale creamy buff with deeper edge, ageing to cream, lightly frilled. Neck grows in water.
- KILTER** 3W-R
Perianth opens creamy primrose, develops to pale cream, creased. Segments flat, pointed, overlapping. Corona deep burnt orange, almost flat, deeply fluted and creased.
- KILWORTH** 2W-YOO
Perianth creamy white, stained yellow at the base, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona brilliant deep red, fluted.
- KIMANGE** 1Y-Y
No description available.
- KINARD** 2W-Y
Perianth white. Segments very broadly ovate and overlapping. Corona frilled cup, white with yellow rim. Neck grows badly in water.
- KINDLED** 2Y-R
Discarded - Small.
- KING ALBERT** 1Y-Y
Perianth bright clear lemon yellow, slightly creased. Segments slightly twisted, pointed, overlapping. Corona deep golden yellow, long but open-mouthed.

- KING ALFRED** 1Y-Y
Perianth bright clear yellow, a little green on back. Segments twisted, pointed, overlapping. Corona deep bright yellow, neat edge.
- KING CARDINAL** 2W-R
Perianth opens primrose, quickly develops creamy white, stained yellow base, smooth, circular outline. Segments flat, pointed, overlapping. Corona greenish gold base 1/3 shading to deep burnt orange red, deeply fluted, deeply cut lobed edge, bowl shape. Neck grows stiffly in water.
- KING KI** 1Y-Y
No description available.
- KING LUD** 1Y-Y
Discarded - Short.
- KING OF HEARTS** 2Y-O
Perianth soft primrose yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona bright burnt orange throughout. Neat and stiff.
- KINGCRAFT** 8W-O
Perianth deep cream stained yellow at base, creased. Segments flat, rounded, well overlapping. Corona burnt orange with greenish eye, fluted. Style longer than stamens. Neck grows in water, and corona edge dies first.
- KINGDOM** 2W-WWY
Perianth creamy white, stained yellow at base, creased. Segments flat, pointed, broad, overlapping. Corona deep lemon yellow, ageing to creamy lemon.
- KINGFISHER** 3W-GYR
Perianth creamy white, stained yellow at base, circular outline. Segments flat, pointed, broad, well overlapping. Corona small moss green eye, changing to narrow band of gold, changing to rim of deep orange red for 1/2 its length, deeply fluted. Stiff.
- KING'S RANSOM** 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled and same yellow as perianth.
- KINGSCOURT** 1Y-Y
Perianth soft lemon yellow, very smooth, forming perfect double triangle. Segments flat, pointed, broad, well overlapping. Corona deeper lemon yellow. Very well balanced. Inclined to remain hooded when opened in water from bud.
- KINGSLEY FAIRBRIDGE** 1W-Y
Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona opens primrose, develops cream, long, rather out of proportion.
- KINGSTON** 2W-Y
Discarded - Flowers below foliage, very susceptible to basal rot.
- KLAMATH** 2W-Y
Discarded - Poor colour, untidy, flowers below foliage.
- KNIPP** 2Y-YOO
Perianth pale primrose yellow, paler at tips, washy. Segments flat, pointed, overlapping. corona gold shading to orange edge for 1/2 its length.
- KNOCKSTACKEN** 1Y-Y
Perianth buttercup yellow. Segments pointed, overlapping, very slightly creased, top segments a little twisted. Corona deep yellow, paler at base, fluted, even crenate edge.
- KNOWEHEAD** 2W-W
Perianth white. Segments broad and overlapping. Corona open mouthed, frilled and same colour as perianth.
- KRAKATOA** 2Y-O
Perianth clear bright yellow, slightly creased, crêpey texture, sometimes catching in corona. Segments flat, pointed, overlapping. Corona bright orange red.
- KWANNON** 2W-Y
Perianth creamy white, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona cream, paler at base, deep bowl shape. Floppy, poor poise. Neck grows in water.
- LA ARGENTINA** 2W-WOR
Perianth white, creased, crêpey texture, notched, often caught in corona. Segments flat, rounded, broad, well overlapping. Corona pale creamy orange shading to deeper orange at edge with deep cut lobes edged with white. Very variable colour.
- LA BEAUTE** 3W-R
Perianth white, creased, reflexed. Segments flat, rounded, overlapping. Corona bright flame red, flat, deeply fluted. Neck grows in water, corona edge dies first.

- LA BELLE** 7Y-O
Perianth pale primrose yellow, segments curl back at edges, sometimes creased. Segments flat, rounded, overlapping. Corona deep gold changing to bright orange edge for half of its length, with greenish gold eye, flat, fluted, unevenly shaped edge.
- LA FIANCEE** 8W-Y
Perianth white, stained slight yellow at base, edges curl in, slightly reflexed. Segments flat, pointed, overlapping. Corona orange gold, with slight green eye, fluted. Style as long as longest stamens.
- LA RIANTE** 3W-YOO
Perianth opens creamy white, develops white, slightly creased. Segments flat, rounded, well overlapping. Corona greenish primrose with bright orange red edge 1/3 of its length and small greenish gold eye, almost flat, fluted. Neck grows in water.
- LABRADOR** 1W-W
Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona creamy white. Stiff and neat, but small.
- LADY BEE** 2W-P
Perianth creamy white, thin, star shape. Segments twisted, pointed, separated. Corona primrose shading to bright peach for 1/2 its length, fluted, straight sided, stiff. Neck grows a little in water.
- LADY BIRD** 2W-O
Perianth creamy white, deep creased, notched, often caught in corona. Segments flat, pointed, overlapping. Corona opens buff pink, develops creamy peach, paler at base ageing to cream, open-mouthed, often split, badly proportioned. Neck grows in water.
- LADY DIANA MANNERS** 3W-YYR
Perianth opens primrose yellow, develops creamy white stained yellow at base, creased. Segments flat, rounded, overlapping. Corona orange shading to bright orange red edge for 1/2 its length, deeply fluted and creased.
- LADY GODIVA** 3W-?YYO
Perianth cream stained yellow at base. Segments slightly twisted, pointed, separated. Corona yellow with bright orange edge for 1/3 of its length, fluted.
- LADY IVOR** 2W-P
Perianth white, a little creased. Segments flat, pointed, well overlapping. Corona creamy apricot, flecked with deeper apricot, a little paler at edge, fluted, bowl shape.
- LADY JEAN RANKIN** 2W-W
Perianth creamy white, star shape. Segments flat, pointed, overlapping. Corona cream, slender trumpet shape.
- LADY JOWITT** 2W-P
Perianth white. Outer segments very broadly ovate, inner whorl ovate, and overlapping. Corona open mouthed, frilled and pale pink in colour. Neck grows badly in water.
- LADY KESTEVEN** 3W-O
Perianth white, stained deep yellow at base, creased, floppy. Segments flat, pointed, overlapping. Corona brilliant orange red, deeply fluted.
- LADY LUCK** 2Y-O
Discarded - Poor colour.
- LADY MARGARET BOSCAWEN** 2W-Y
Perianth creamy white, stained yellow at base, creased, star-shaped. Segments flat, pointed, overlapping. corona bright lemon yellow, almost straight sided.
- LADY MOORE** 3W-OOR
Perianth cream, deeper at base. Segments flat, rounded, overlapping. Corona bright lemon changing to bright orange red edge for 1/5 of its length, fluted.
- LADY PENRHYN** 2Y-YYO
Perianth bright pale lemon yellow. Segments flat, rounded, overlapping. Corona gold with shaded orange edge, fluted, stiff.
- LADY SERENA** 9W-GYR
Perianth poeticus white, stained yellow base, full rounded outline. Segments rounded, overlapping, broad, slightly creased, curled in at edges. Corona moss green throat, changing to citrus yellow with faint white ring before very narrow brick red rim, expanded, frilled edge. Neck grows in water.
- LAETITIA** 8Y-YYO
Perianth yellow. Segments rounded and overlapping. Corona small cup and dark yellow in colour.
- LAGONDA** 2W-YYW
Perianth creamy white, creased, crêpey texture. Segments flat, pointed, broad, well overlapping. Corona pale lemon, with paler edge, developing to deep cream, of almost trumpet proportions.

- LAMINGTON** 2Y-O
Perianth deep lemon yellow, suffused orange at base, sometimes split, a little creased, pointed outline. Segments flat, pointed, overlapping. Corona deep burnt orange shading to orange base, fluted, broad bowl shape. Stiff. Neck grows a little in water.
- LANARTH** 7Y-O
Perianth deep clear yellow, slightly creased, crêpey texture. Segments flat, rounded, overlapping. Corona orange-gold, colour varies a little, irregularly lobed. Neck grows in water.
- LANCASTER** 3W-GYO
Perianth creamy white, stained yellow base, double triangle outline. Segments rounded, overlapping at base, bottom segments very broad, top segments with raised centre line, smooth, curled deeply at edges. Corona moss green throat, changing to citrus yellow with bright orange edge; expanded, deeply lobed edge.
- LANDITHY** 2Y-Y
Perianth bright gold, creased, slightly hooded, crêpey texture, pointed outline. Segments flat, pointed, overlapping. Corona orange-gold.
- LANDULPH** 2Y-Y
Perianth pale lemon yellow, creased, crêpey texture, rough. Segments flat, pointed, broad, well overlapping. Corona bright lemon yellow, of trumpet proportions.
- LANGLO** 1Y-Y
Perianth pale sulphur, smooth, slightly hooded. Segments broad, overlapping to half the length. Corona lemon, long with obvious flange, deeply cut frilled edge. Hollow stems.
- LANTANA** 1W-Y
Perianth creamy white, stained yellow base, top 3 segments inclined to incurve. Segments flat, pointed, overlapping. Corona deep lemon, crenate, flanged, broad mouth. Badly proportioned.
- LANTEGLOS** 2
Perianth opens sulphur, developing paler tips, creased, crêpey, pointed outline. Segments flat, pointed, overlapping. Corona light orange, paler at base, variable colour, fluted, lightly frilled, bowl shape. Neck grows in water.
- LAPFORD** 1W-Y
Discarded - Short, hangs head.
- LAPLAND** 1W-W
Perianth white, slightly creased, crêpey texture. Segments slightly twisted, pointed, broad, well overlapping. Corona cream, flanged, broad mouth but well proportioned.
- LARGO** 2Y-YYO
Discarded - Very frilly corona with white blotches on edge.
- LARKELLY** 6Y-O
Perianth deep primrose, paler at tips, reflexed. Segments flat, pointed, overlapping. Corona deep gold, shading to orange edge for 1/2 of its length.
- LATIFA** 2W-O
Perianth opens primrose, develops cream, creased. Segments flat, pointed, overlapping. Corona bright orange, deeply fluted, uneven deep cut edge.
- LATONIA** 3W-R
Perianth opens cream, develops creamy white, stained yellow at base, reflexed. Segments flat, pointed, overlapping. Corona burnt orange red with small greenish gold eye, flat, deeply fluted, creased.
- LATROBE** 1Y-Y
Perianth yellow. Segments broad, overlapping and slightly twisted. Corona large open mouthed, frilled and same colour as perianth. Neck grows in water. Weak stems.
- LAURENS KOSTER** 8W-Y
Perianth creamy white, creased, edges curled in. Segments flat, rounded, overlapping. Corona golden orange, fluted. Style reaches only as far as shortest stamens.
- LAVINO** 2W-P
Perianth creamy white, creased, edges curl in, pointed outline. Segments twisted, pointed, barely overlapping. Corona greenish cream shading to bright apricot orange for 1/3 its length, some white blotches on edge. Poor poise. Neck grows badly in water.
- LAWALI** 4W-P
Perianth white. Segments very broadly ovate and overlapping. Corona frilled, open mouthed and pink. Neck grows badly in water.
- LE BEAU** 6Y-Y
Perianth bright yellow, a little floppy. Segments slightly twisted, pointed, barely overlapping. Corona deeper yellow, slightly waisted.

- LEANDER** 2Y-O
Perianth yellow. Segments ovate and overlapping. Corona frilled cup and orange in colour. Neck grows badly in water.
- LEBANON** 2W-Y
Perianth cream, slightly reflexed. Segments blunt, some notched near end, overlapping. Corona primrose at base with cream rim, open mouthed, ribbed, frilled rim. Neck grows stiffly in water. Hollow stems.
- LEEUWENHORST** 2W-YYO
Perianth creamy white stained yellow at base, creased. Segments flat, pointed, broad overlapping. Corona deep apricot orange.
- LEINSTER** 1Y-Y
Perianth soft lemon yellow, crêpey texture, forming double triangle. Segments flat, pointed, broad, overlapping, shovel shaped. Corona deeper lemon yellow.
- LEITRIM** 2Y-Y
Perianth lemon, slightly creased. Segments flat, pointed, overlapping. Corona deeper lemon, crenate, slightly flanged, trumpet proportion. Stiff.
- LEMBERG** 1Y-Y
Perianth soft sulphur yellow, green centre line on back. Segments flat, pointed, overlapping. Corona deeper lemon yellow, rather long and slender, deeply crenate.
- LEMNOS** 2Y-Y
Perianth opens primrose, develops cream, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona opens pale lemon, developing to buff apricot, paler at base.
- LEMON BEAUTY** 11W-Y
Perianth poeticus white, creased, crêpey, edges curl in slightly, circular outline. Segments flat, rounded, broad, overlapping. Corona split into 6 segments, each white with central lemon stripe for 3/4 of its length, merging into greenish lemon base, frilled. Remains hooded in water.
- LEMON CUP** 2W-Y
Perianth creamy white, deeply creased, often buckled. Segments flat, pointed, overlapping. Corona opens gold, develops pale apricot with narrow rim of primrose, deeply fluted. Rough and ugly.
- LEMON DORIC** 2Y-W
Perianth very pale lemon. Segments broad and ovate. Corona open mouthed, frilled, same colour as perianth with an intermittent yellow rim. Neck grows in water.
- LEMON FANCY** 1Y-Y
Perianth opens greenish primrose, developing cream, creased, crêpey texture. Segments flat, pointed, overlapping. Corona opens primrose yellow, develops cream with deeper edge. Perianth sometimes catches in corona.
- LEMONADE** 3Y-Y
Perianth creamy white, stained yellow base, rounded outline. Bottom segments broad, narrower top segments creased, some buckled, all curled in at edges. Corona lemon base fading to creamy white edge, expanded, lobed edge.
- LEMSTAR** 1Y-Y
Perianth deep bright yellow, some green on back. Segments flat, pointed, overlapping. Corona deep golden yellow.
- LENORE NETHERCOTE** 2Y-YYO
Perianth yellow. Segments broad and overlapping. Corona open mouthed, frilled, yellow with an orange rim. Neck grows badly in water.
- LEONAINÉ** 2W-P
Perianth white. Segments rounded/very broadly ovate and overlapping. Corona large frilled cup and pink in colour. Neck grows in water.
- LEPRECHAUN** 2Y-R
Perianth lemon yellow. Segments flat, pointed, overlapping. Corona deep orange red, straight sided, burns easily. Corona edge dies first. Neck grows in water.
- LETTERKENNY** 1Y-Y
Perianth lemon, creased. Segments flat, pointed, overlapping. Corona deeper lemon, crenate, slightly flanged. Barely full length trumpet. Stiff.
- LEVIATHAN** 2Y-Y
Perianth pale sulphur yellow, deeper at base, creased, floppy. Segments twisted, slightly reflexed, rounded, overlapping. Corona deep orange-gold shading to deep bright orange edge for 1/4 of its length.
- LHASA** 1W-W
No description available.

- LIBBY HOLMAN 1W-W
Perianth white. Outer segments broad, inner ovate, and overlapping. Corona frilled open mouthed, same colour as perianth. Neck grows in water.
- LIBERTY 1Y-Y
Perianth deep primrose yellow, green on back. Segments twisted, pointed, separated, curled back at edges. Corona deep lemon yellow, deeply frilled and creased down length. Badly balanced.
- LIBERTY LIGHT 2W-YYO
No description available.
- LIBYA 2W-O
Perianth white. Segments broad and overlapping. Corona large frilled cup, deep orange in colour with a darker rim. Neck grows badly in water.
- LIGHTHEART 1W-Y
Discarded - Short, flowers below foliage, hangs head badly.
- LIKOVAN 1Y-Y
Perianth yellow. Segments broad, 1/2 overlapping and twisted. Corona large frilled open mouthed and same colour as perianth. Neck grows in water.
- LILY RONALDS 2W-GYO
Perianth white. Segments very broadly ovate and overlapping. Corona large frilled cup, green in centre, then yellow, with an orange rim. Neck grows in water.
- LILYDALE 2W-W
Perianth white. Outer segments broadly ovate, inner ovate, and overlapping. Corona large open mouthed, frilled and yellow.
- LIMELIGHT 1Y-Y
Perianth creamy sulphur, paler at base, deeper on back, crêpey. Segments twisted, pointed, overlapping. Corona pale creamy lemon with rim of lemon, frilled. Hangs head.
- LIMERICK 3W-O
Perianth opens cream, develops white stained cream at base, smooth. Segments flat, pointed, broad, overlapping. Corona deep sealing wax red, almost flat, fluted, lobed edge. Neck grows in water.
- LIMONE 1Y-Y
Discarded - Poor shape and colour.
- LINGERING LIGHT 2W-P
Discarded - Poor colour.
- LINN 2W-Y
Discarded - Short, poor colour.
- L'INNOCENCE 8W-Y
Perianth white, creased, curled in at edges. Segments flat, rounded, overlapping. Corona bright gold edged light orange, fluted. Style reaches a little longer than shortest stamens.
- LINTIE 7Y-O
Perianth opens primrose, developing deep cream, often buckled. Segments flat, rounded, overlapping. Corona moss green eye, shading to gold for 1/2 its length, with bright orange edge, almost flat, fluted, unevenly lobed edge. Neck grows badly in water.
- LISANORE 2W-P
Perianth white, stained slight green back base, double triangle outline. Segments rounded, overlapping, creased particularly at edge, narrower top segments curled deeply forward. Corona deep shell pink, fluted, lobed edge.
- LISBANE 3W-GYR
Perianth creamy white, stained yellow base, full rounded outline. Segments rounded, overlapping, broad, generally smooth, top segments some wrinkled lower edge. Corona moss green throat, developing to citrus yellow mid-third and changing to light brick red top third; expanded, irregular deeply lobed edge. Neck grows badly in water.
- LISBREEN 2W-GOO
Perianth creamy white, slightly creased, inclined to incurve, pointed outline. Segments flat, pointed, well overlapping. Corona opens cream tinted peach, develops peach with deeper edge, a little fluted. Neck grows badly in water.
- LISCARTON 3Y-O
Perianth soft sulphur yellow, deeper at base, crêpey texture, a little creased. Segments flat, pointed, broad, overlapping. Corona burnt orange.
- LISDILLON 1W-P
Discarded - Short, small, flowers below foliage.

- LITTLE DIAMOND** 3Y-O
Perianth sulphur yellow, paler at tips, deeply creased. Segments flat, pointed, overlapping. Corona bright orange red, deeply fluted and frilled, almost flat. Neck grows badly in water.
- LITTLE WITCH** 6Y-Y
Perianth lemon yellow, creased, reflexed. Segments flat, pointed, barely overlapping. Corona bright gold, fluted, pinched in.
- LIVIA** 4Y-O
Perianth segments pale lemon yellow, flat, rounded, overlapping, with one or two complete outer rows, reflexed. Corona orange gold, almost complete, packed tightly with incurving perianth segments intermixed with corona segments. Neck grows in water.
- LIZARD LIGHT** 2Y-O
Perianth pale lemon yellow, a little creased. Segments flat, pointed, overlapping. Corona bright orange red, a little paler at base, fluted, bowl shape, deeply frilled. Neck grows in water.
- LOCH HOPE** 2Y-R
Perianth bright rich lemon. Segments rounded, overlapping at base, leathery appearance. Corona burnt orange, goblet shaped, frilled edge. Neck grows badly in water.
- LOCH MAREE** 2W-P
Perianth creamy white, creased, often caught in corona. Segments flat, pointed, overlapping. Corona opens creamy lemon, develops creamy peach, ages to cream, of trumpet proportion. Rough. Neck grows badly in water.
- LOCHINCH** 2Y-O
Perianth yellow. Segments very broadly ovate, overlapping, with ridged texture. Corona small, frilled orange cup. Neck grows badly in water.
- LOCHINVAR** 2W-O
Perianth white. Segments very broadly ovate and overlapping, and slightly inflexed. Corona small frilled orange cup. Neck grows badly in water.
- LOGAN ROCK** 7
Perianth yellow. Outer segments very broad, inner segments ovate, and overlapping. Corona medium sized frilled cup, same yellow as perianth. Neck grows badly in water.
- LOMBARD** 2Y-Y
Perianth clear lemon yellow, a little creased. Segments flat, pointed, broad, overlapping. Corona deep lemon gold.
- LONGFORD** 1W-W
Perianth white. Segments broad and overlapping, ridged texture, inner segments slightly twisted, and slightly inflexed. Corona flared, frilled funnel, same colour as perianth.
- LORD ANTRIM** 1Y-Y
Perianth deep, soft, sulphur yellow. Segments slightly twisted, pointed, broad, overlapping. Corona bright deep lemon yellow. Neck grows in water.
- LORD CHANCELLOR** 1W-Y
Discarded - Short.
- LORD KITCHENER** 2W-Y
Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona pale creamy primrose yellow, developing paler with age.
- LORD NELSON** 1Y-Y
Perianth sulphur yellow, a little creased, crêpey texture. Segments slightly twisted, pointed, overlapping. Corona lemon yellow, open-mouth. Dumpy appearance.
- LORD TEDDER** 2W-GYY
Perianth creamy white, creased, edges curl in a little, circular outline. Segments flat, rounded, well overlapping. Corona greenish primrose, cut into 6 lobes striped vertically with cream pale orange edges, variable colour, fluted. Stiff. Neck grows in water.
- LORD WELLINGTON** 1Y-Y
Perianth bright lemon yellow, crêpey texture. Segments slightly twisted, pointed, barely overlapping. Corona deeper lemon yellow.
- LORELEI** 2W-P
Perianth muddy white, 3 top segments a little hooded, 3 lower segments reflexed. Segments flat, rounded, overlapping. Corona deep apricot shading paler at base, fluted.
- LORENZO** 2W-O
Perianth creamy white, stained yellow base. Segments rounded, overlapping at base. Corona burnt orange, paler at edge, with six white blotches, expanded, frilled edge.

- LOTHARIO** 2W-Y
Perianth yellow. Segments very broadly ovate and overlapping. Corona medium sized frilled cup, orange in colour. Neck grows badly in water.
- LOTTA SVARD** 1W-Y
Perianth cream, stained yellow, for 1/3 of the base. Segments flat, pointed, overlapping. Corona very deep lemon yellow, long trumpet.
- LOUGH AREEMA** 3W-GWO
Perianth poeticus white, edges curl in a little, smooth. Segments flat, pointed, overlapping. Corona bright green eye, changing to greyish cream with narrow edge of bright orange, fluted, almost flat. Neck grows in water.
- LOUGH ERNE** 3W-YYR
Perianth creamy white, slightly creased, back segments reflex, circular outline. Segments flat, rounded, well overlapping. Corona small yellow green eye, changing to greenish lemon, then to deep orange band for ½ its length, fluted, frilled, lobed, almost flat. Neck grows badly in water.
- LOUGH NEAGH** 3W-Y
Perianth creamy white, crêpey, slightly creased, top segments curl in a little, circular outline. Segments flat, rounded, broad, overlapping. Corona grey green eye changing to buff, developing to creamy white, deeply fluted, frilled, deeply lobed, almost straight sided. Neck grows in water.
- LOUGHGALL** 2W-P
Perianth white. Segments broad and overlapping, and inflexed. Corona frilled, lemon yellow in colour. Neck grows badly in water.
- LOUIS D'OR** 2Y-Y
Discarded - Short, flowers below foliage, poor poise.
- LOUIS PASTEUR** 2W-GYO
Perianth cream stained yellow at base, creased. Segments flat, pointed, broad, overlapping. Corona gold with bright deep orange edge for 1/2 of its length and a greenish gold eye, fairly flat, fluted.
- LOUISE DE COLIGNY** 2W-O
Perianth creamy white, edges curl back, creased and notched. Segments twisted, pointed, overlapping. Corona salmon pink shading to pale cream tinted apricot. Untidy looking flower.
- LOVE DREAM** 3W-O
Perianth creamy white, very creased. Segments flat, rounded, well overlapping. Corona bright orange red throughout, deeply fluted, creased, uneven shape.
- LOVE SONG** 2W-YOO
Perianth creamy white, slightly hooded. Segments very broad, papery, creased, overlapping very extensively. Corona pale orange with cream rim, flat to open mouthed, fluted with uneven gently frilled rim. Overall flower looks untidy and crumpled. Neck grows stiffly in water.
- LOVENEST** 2W-YP
Perianth white, slightly creased. Segments flat, pointed, broad, overlapping. Corona opens primrose yellow, developing to apricot then ageing to paler colour. Similar to Suda.
- LUCCOMBE** 2Y-O
Perianth bright clear lemon yellow, creased, star shaped. Segments flat, pointed, overlapping. Corona bright orange red shading to slight gold base, fluted.
- LUCIFER** 2W-YOO
Perianth creamy primrose yellow, stained yellow at base, star shaped, floppy. Segments twisted, pointed, separated. Corona bright orange red, paler at base.
- LUDLOW** 2W-W
Perianth white, slightly creased, often caught in corona, double triangle. Segments flat, pointed, broad, well overlapping. Corona opens pale lemon, quickly develops pure white. Well proportioned.
- LUNAR MOTH** 1Y-Y
Perianth pale yellow. Segments ovate and overlapping, slightly notched. Corona frilled flared, funnel, yellow (as perianth) with a darker yellow rim. Neck grows in water.
- LUNAR RAINBOW** 2W-Y
Perianth creamy white, creased, pointed outline. Segments flat, pointed, overlapping. Corona opens pale primrose, develops pale creamy peach, open-mouthed, fluted. Poor poise. Neck grows in water.
- LUNAR SEA** 1Y-W
Perianth yellow. Segments broad and overlapping and slightly twisted. Corona frilled, flared cylinder, same colour as perianth. Neck grows in water.
- LUNAR SPELL** 2Y-WWY
Perianth pale creamy yellow. Segments broad overlapping, and crêpey. Corona frilled open mouthed, same colour as perianth with a yellow rim. Neck grows in water.

- LUNE DE MIEL** 4W-Y
Perianth segments white, flat, pointed, overlapping, two complete rows. Corona segments open lemon yellow, develop apricot, intermixed in centre with a few long pointed perianth segments.
- LURGAIN** 1Y-Y
Perianth yellow. Segments broad, overlapping, turned in at edges and inflexed. Corona large frilled open mouthed, and same colour as perianth.
- LUXURY** 1Y-Y
Perianth bright deep lemon yellow, crêpey texture, stiff. Segments slightly twisted, pointed, broad, overlapping. Corona deeper bright lemon yellow.
- LYN PARKER** 1Y-Y
Perianth lemon yellow. Segments broad, overlapping crêpey and inflexed. Corona large frilled open mouthed, and darker yellow than perianth.
- LYON** 2W-OOY
Perianth creamy white, a little creased. Segments flat, pointed, well overlapping. Corona apricot with narrow rim of pale creamy apricot, fluted.
- LYSANDER** 2W-YYO
Perianth white. Segments very broadly ovate, overlapping, and edges turned in. Corona frilled cup, pale yellow with a darker yellow rim. Neck grows in water.
- M.J. BERKELEY** 1Y-Y
Perianth deep lemon yellow, much green on back, thin, spidery appearance. Segments twisted, pointed, separated. Corona deeper lemon yellow, slender base, broad mouth, very deeply lobed and frilled edge.
- MABEL TAYLOR** 2W-WPP
Perianth white. Segments broad and overlapping, with ridged texture. Corona medium sized frilled cup yellow with an orange rim. Neck grows badly in water.
- MACEBEARER** 2W-YYO
Perianth creamy primrose yellow, stained yellow at base, creased. Segments twisted, rounded, overlapping. Corona deep gold shading to bright orange edge for 1/3 its length, fluted.
- MADEIRA** 2Y-O
Perianth yellow. Segments very broadly ovate and overlapping. Corona frilled trumpet, orange in colour. Neck grows in water.
- MADOLON** 2Y-O
Perianth cream stained yellow at base, smooth, circular outline. Segments flat, rounded, well overlapping. Corona bright orange with thin gold rim, fluted, almost flat. Stems split and curl in water. Neck grows in water.
- MADRIGAL** 2W-WWY
Discarded - Hangs head, flowers below foliage.
- MAGHERALLY** 2Y-O
Perianth deep bright lemon yellow, slightly creased. Segments flat, rounded, well overlapping. Corona deep orange red shading to slight orange base, straight sided.
- MAGIC CIRCLE** 2W-YYR
Discarded - White blotches and rough corona edge.
- MAGIC DAWN** 2W-P
Perianth white. Segments very broadly ovate, overlapping, and crêpey. Corona large frilled cup, pink in colour. Neck grows badly in water.
- MAGISTRATE** 1Y-Y
Perianth yellow. Segments broad, overlapping, and twisted. Corona large frilled, flared cylinder, same colour as perianth. Neck grows badly in water.
- MAGNET** 1W-Y
Perianth cream, stained yellow at base, crêpey texture. Segments twisted, pointed, overlapping. Corona deep bright lemon yellow.
- MAGNIFICENCE** 1Y-Y
Perianth deep maximus gold, crêpey texture, greenish back. Segments twisted, pointed, overlapping. Corona deep gold slightly paler at base, deeply frilled.
- MAHEE** 1Y-Y
Discarded - Small.
- MAHMOUD** 3W-R
Perianth creamy white, smooth, circular outline. Segments flat, rounded, well overlapping. Corona bright deep red, fluted, almost flat. Neck grows in water.

- MAID OF ULSTER** 2W-YYR
Perianth white, stained yellow base, circular outline. Segments broad, rounded, overlapping at base, smooth, curled in at edges. Lower two-thirds corona deep citrus yellow changing to bright orange edge, expanded, deeply lobed edge.
- MAIDEN'S BLUSH** 2W-P
Perianth creamy white, crêpey texture, often caught in corona. Segments flat, pointed, broad, overlapping. Corona cream base shading to pale coppery peach for 1/3 its length, sometimes split, of trumpet proportions. Neck grows in water.
- MAIVEROE** 1W-W
Perianth white, smooth. Segments flat, pointed, broad, overlapping, shovel shape. Corona creamy white, slender, neat.
- MAJARDE** 2Y-O
Perianth sulphur yellow, a little paler at tips, creased and crêpey texture. Segments slightly twisted, pointed, broad, overlapping. Corona bright orange, flattened, deeply fluted.
- MAJESTIC** 1Y-Y
No description available.
- MAJOLICA** 2Y-Y
Discarded - Poor colour, short, flowers below foliage.
- MAJORCA** 2Y-YYO
Perianth yellow. Segments very broadly ovate and overlapping. Corona frilled cup, yellow with orange rim. Neck grows badly in water.
- MAJORETTE** 1Y-Y
Perianth pale yellow. Segments broad and overlapping, and slightly reflexed. Corona large frilled open mouthed, same colour as perianth.
- MAKASSAR** 2Y-Y
Perianth lemon yellow, creased, crêpey texture, perianth often catches in corona. Segments twisted, pointed, broad, well overlapping. Corona bright gold.
- MALHEUR** 2W-Y
Perianth white. Segments broad and overlapping. Corona frilled open mouthed and yellow in colour. Neck grows in water.
- MALONE** 1Y-Y
Discarded - Short, hangs head, flowers below foliage.
- MALTA** 2Y-O
Perianth lemon yellow, odd segments sometimes creased. Segments flat, rounded, overlapping. Corona orange red shading to gold base. Stiff and sturdy. Neck grows in water.
- MALVERN CITY** 1Y-Y
Perianth lemon, edges curl in a little, slightly hooded. Segments flat, pointed, overlapping. Corona dark lemon, crenate, flanged, neat edge. Neck grows in water.
- MALVERN GOLD** 2Y-Y
Perianth sulphur, ribbed, twisted. Segments narrow, pointed, only overlapping at base, prominently mucronate. Corona deep sulphur, short, flanged, deeply frilled edge. Neck grows badly in water.
- MAMOUTH** 3W-R
Perianth white. Segments rounded and overlapping. Corona small frilled cup, orange/red in colour. Neck grows badly in water.
- MANCHU** 2W-Y
Perianth creamy white. Segments flat, pointed, broad, well overlapping. Corona opens bright lemon develops to buff-orange. Corona too large for the small perianth.
- MANCO** 2W-YYO
Perianth opens primrose, develops cream stained yellow base, sometimes catches in corona. Segments flat, pointed, overlapping. Corona orange shading to deeper orange edge, deeply fluted. Badly proportioned.
- MANDATE** 1W-W
Perianth white, slightly creased. Segments flat, pointed, broad, well overlapping. Corona creamy white, slender, almost straight sided.
- MANDO** 1Y-Y
Perianth yellow. Segments ovate, slightly overlapped and twisted. Corona large frilled open mouthed, and same colour as perianth. Neck grows in water.
- MANDOLIN** 2W-GPP
Discarded - Short, floppy flower.

- MANGOSTEEN** 3Y-R
Perianth deep primrose yellow, paler at tips, a little creased, crêpey texture, perianth catches in corona. Segments flat, pointed, broad, overlapping. Corona burnt orange, fluted.
- MARA** 9W-YYO
Perianth poeticus white. Segments very broadly ovate and overlapping. Corona very small frilled cup, yellow with orange rim. Neck grows badly in water.
- MARAVEL** 1Y-Y
Perianth opens pale greenish primrose, developing greenish cream, deeper at base. Segments twisted, pointed, broad, well overlapping. Corona opens greenish primrose, developing cream with narrow rim of lemon.
- MARCH MADNESS** 2Y-R
Perianth bright mid lemon. Segments pointed, overlapping, bottom segments broad, creased, some buckled, curled in at edges. Corona burnt orange, tinged red, fluted frilled edge. Neck grows in water.
- MARCH SUNSHINE** 6Y-Y
Perianth clear yellow, slightly reflexed. Segments flat, pointed, barely overlapping. Corona orange gold, pinched in.
- MARCOA** 2Y-O
Perianth yellow. Segments ovate, overlapping, and very slightly inflexed. Corona frilled cup, orange in colour. Neck grows in water.
- MARCURA** 2Y-R
Perianth yellow. Segments broad, overlapping, with ridged texture. Corona small frilled cup, orange in colour. Neck grows badly in water.
- MARFORT** 2
No description available.
- MARGARET MITCHELL** 3W-YYR
Perianth poeticus white, stained yellow base. Segments broad, rounded, overlapping, edges curling slightly forwards. Corona short and flat, orange becoming red at edge. Neck grows badly in water.
- MARGRIET** 2W
Perianth creamy white, creased, notched, often caught in corona, edges curl in a little, pointed outline. Segments flat, pointed, overlapping. Corona cream base, shading to primrose with narrow cream edge, fluted, frilled, deeply lobed, deep bowl shape. Neck grows in water.
- MARIMBA** 2Y-YYO
Perianth lemon. Inner segments narrower than outer segments. Segments pointed, overlapping with some deeply creased. Corona lobed, frilled, slightly flanged edge, burnt orange with a paler centre.
- MARION CRAN** 2Y-O
Perianth primrose yellow, creased, edges curl in. Segments flat, pointed, overlapping. Corona deep orange with slight gold edge, fluted. Neck grows in water.
- MARKANT** 3W-R
Perianth opens cream, develops pure white stained yellow at base, curled in at edges. Segments flat, rounded, well overlapping. Corona deep orange-red, paler at base, fluted.
- MARKET GEM** 2W-Y
Perianth dirty creamy white stained yellow at base and on the back. Segments flat, pointed, overlapping. Corona deep bright lemon yellow.
- MARKET MERRY** 3Y-O
Perianth soft sulphur yellow ageing to cream, a little creased, crêpey texture. Segments flat, rounded, well overlapping. Corona light orange, fluted. Neck grows in water.
- MARKETHILL** 1Y-Y
Perianth lemon, creased. Segments flat, pointed, overlapping. Corona deeper lemon, lightly frilled, neat edge, almost straight sided.
- MARKSMAN** 2Y-O
Perianth pale lemon yellow, crêpey texture. Segments flat, pointed, overlapping. Corona deep burnt orange, shading to orange base. Very stiff and sturdy.
- MARS** 2Y-R
Perianth deep lemon, inward curling. Segments would fit into a double triangle, forward curling. Corona orange, small and straight sided. Neck grows stiffly in water.
- MARSHAL TSJOEKOF** 2W-OOR
Perianth creamy white, creased, floppy, often buckled. Segments slightly twisted, pointed, overlapping. Corona gold base shading to bright apricot orange with paler edge, fluted, almost flat.

- MARTHA WASHINGTON** 8W-O
Perianth creamy white stained yellow at base, a little creased. Segments flat, pointed, broad, overlapping. Corona orange gold shading to deep orange edge, and greenish gold eye, almost flat, fluted. Neck grows in water.
- MARY ANN** 2W-W
Perianth white. Segments broad and overlapping. Corona large frilled, flared funnel, same colour as perianth.
- MARY BLONK** 2W-YOO
Discarded - Floppy flower, white blotches on corona edge.
- MARY BOHANNON** 2Y-O
Perianth yellow. Segments broadly ovate and overlapping. Corona large frilled cup, orange in colour. Neck grows badly in water.
- MARY COPELAND** 4W-O
Perianth segments cream, stained yellow at base, flat, pointed, broad, overlapping, one complete row. Corona segments bright red shading to yellow base intermixed in centre with 6-8 pointed perianth segments.
- MARY HOUSLEY** 2W-YYO
Perianth cream, stained slight yellow at base, curled in at edges, deeply creased. Segments flat, rounded, overlapping. Corona bright orange red, deeper at base, untidy edge.
- MARY ROOZEN** 2Y-O
Perianth light lemon yellow, a little creased. Segments flat, pointed, broad, overlapping. Corona bright burnt orange shading to orange gold base. Neck grows in water.
- MASAI KING** 2Y-O
Perianth yellow. Segments broad and overlapping. Corona small frilled cup, and orange in colour. Neck grows in water.
- MASAKA** 3W-R
Perianth creamy white, stained yellow base, creased, circular outline. Segments flat, rounded, overlapping. Corona sealing wax red, deeply fluted, frilled, unevenly lobed edge, almost flat. Neck grows badly though stiffly in water.
- MASKED LIGHT** 2Y-O
Perianth very deep lemon, overlaid gold at base, slightly creased, inclined to reflex. Segments flat, pointed, overlapping, inclined to curl back. Corona deep bright orange red, slightly fluted, frilled, bowl shape. Neck grows stiffly in water.
- MASTER FLORIS** 1W-Y
Perianth creamy white, stained yellow at base, slightly creased. Segments flat, pointed, overlapping. Corona creamy lemon. Squat appearance.
- MASTER ROBERT** 1Y-Y
Perianth soft lemon yellow, pointed outline. Segments flat, pointed overlapping. Corona bright lemon yellow.
- MASTERPIECE** 3W-OOR
Perianth cream, deeper at base, reflexed. Segments flat, pointed, overlapping. Corona deep orange red, almost flat, fluted, uneven edge.
- MATA HARI** 2W-O
Perianth creamy white, creased, one segment often buckled, edges curled back. Segments flat, pointed, broad, overlapping. Corona opens bright orange, develops to deep apricot with paler edge, fluted, almost flat. Corona colour variable.
- MATADOR** 8Y-O
Perianth pale primrose yellow, with darker base and edges. Segments round and overlapping. Corona lobed and frilled with moss green centre and orange remainder. 3 florets/stem. Neck grows badly in water.
- MATAPAN** 3W-R
Perianth poeticus white, stained slight yellow base, smooth, circular outline. Segments flat, rounded, broad, overlapping. Corona deep burnt orange red, deeply fluted, frilled, lobed, nearly flat. Neck grows stiffly in water.
- MATCHLESS** 2W-O
Discarded - Floppy flower.
- MATLOCK** 2Y-R
Perianth sulphur yellow, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona deep burnt orange, some white blotches on edge, deeply frilled. Poor poise.
- MAVISTON** 2Y-Y
Perianth light lemon yellow, creased, crêpey texture. Segments flat, pointed, broad, well overlapping. Corona deeper lemon yellow, of trumpet proportions. Neck grows in water.

- MAXIMUS SUPERBUS** 1Y-Y
Perianth deep bright gold. Segments flat, pointed, broad, overlapping. Corona deeper gold a little paler at base, deeply lobed. Squat looking flower.
- MAY MOLONY** 2W-Y
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona pale lemon yellow, trumpet shape, neat.
- MAY MURIEL** 2W-Y
Perianth creamy white, creased, slightly twisted. Segments would fit into a double triangle, broad based, overlapping. Corona flat to open mouthed, pale primrose, crenate, deeply frilled. Neck grows stiffly in water. Hollow stems.
- MAY QUEEN** 2W-GYP
Perianth creamy white, green back base. Segments overlapping, bottom Segments broad, rounded, top narrower, pointed, creased, curled in at edges. Corona clear pale lemon with faintest apricot tinge, expanded, frilled edge.
- MAYMYO** 2Y-O
Perianth opens mid lemon, develops pale tips, crêpey, deeply creased, top 3 segments incurve, buckled, sometimes caught in the corona. Segments flat, pointed, overlapping. Corona bright gold base ½, shading to deep orange, fluted, frilled, lobed, bowl shape. Neck grows in water.
- MEDAILLON** 2Y-O
Discarded - Small.
- MEDALIST** 2W-YPP
Perianth creamy white. Segments broad and overlapping. Corona large frilled cup, yellow in colour. Neck grows badly in water.
- MEDINA** 4W-W
Perianth poeticus white, flat, rounded, overlapping, one complete row, circular outline. Centre made up of 6-8 narrow, creased, pointed perianth segments intermixed with very short cream corona segments. Neck grows very badly in water.
- MEDUSA** 8W-O
Perianth white, stained slight yellow at base. Segments flat, pointed, broad, overlapping. Corona deep red with slight green-gold base, fluted. Style much below the shortest stamens.
- MEGA** 9W-GYR
Perianth white, deeply creased, notched, often buckled. Segments flat, rounded, overlapping. Corona greenish lemon, changing to band of deep orange red for 1/4 its length, fluted, almost flat. Neck grows in water.
- MELANITE** 3W-YOO
Perianth white, stained, yellow at base, deeply creased, edges curl in, star shape. Segments flat, pointed, overlapping. Corona greenish gold eye changing to gold with rim of bright orange red for 1/3 its length, fluted, almost flat. Neck grows in water.
- MELISSA** 1Y-Y
No description available.
- MELODY LANE** 2W-P
Perianth white. Segments ovate and overlapping. Corona flared cylinder, very pale pink. Neck grows badly in water.
- MELROSE** 3W
Perianth opens primrose, develops cream stained yellow base, paler at tips, slightly creased, reflexed, circular outline. Segments flat, rounded, overlapping. Corona small greenish gold eye changing to bright orange red, fluted, frilled, uneven lobed edge, shallow bowl shape. Neck grows stiffly in water.
- MENTON** 2W-P
Perianth creamy white, creased, edges curled in, crêpey texture. Segments twisted, pointed, overlapping. Corona opens apricot with paler base, developing to salmon pink with pale salmon base, ages to cream with pale salmon edge. Rough.
- MERCATO** 2W-YYO
Perianth opens primrose, develops to creamy white stained slight yellow at base, deeply creased. Segments flat, pointed, broad, well overlapping. Corona gold, shading to bright orange red edge for 1/2 its length, deeply fluted. Stiff and sturdy.
- MERIDIAN** 2Y-YYO
Perianth bright lemon, slightly creased and crêpey, pointed outline. Segments flat, pointed, overlapping, edges curl in. Corona deep lemon base 1/2 changing to orange red, fluted, lightly frilled, shallow bowl shape. Neck grows badly though stiffly in water.
- MERIVALE** 2W-W
Perianth creamy white, creased, crêpey texture. Segments flat, pointed, overlapping. Corona opens primrose, develops cream, neat edge. Hangs head.

- MERLIN** 3W-YYR
Perianth white. Segments rounded and overlapping. Corona medium sized frilled cup yellow with orange rim. Neck grows badly in water.
- MESSENGER** 2Y-O
Perianth sulphur yellow, paler at tips, creased, crêpey texture. Segments flat, pointed, well overlapping. Corona shading paler at base, variable streaky colour, fluted, untidy edge. Rough. Neck grows stiffly in water.
- METROPOLITAN** 2W-O
Perianth creamy white, slightly creased, circular outline. Segments flat, pointed, broad, well overlapping. Corona bright orange, a shade paler a edge, deeply fluted, almost flat.
- MEXICO** 2Y-O
Perianth clear bright lemon yellow, smooth, star-shaped. Segments flat, pointed, overlapping. Corona deep burnt orange red, goblet shape, fluted.
- MEXICO CITY** 2Y-O
Perianth yellow. Segments rounded and overlapping, and lighter in centre. Corona small frilled cup, orange in colour. Neck grows badly in water.
- MIKADO** 2W-O
No description available.
- MILAN** 9W-GYR
Perianth pure white, creased, edges curl in, circular outline. Segments flat, rounded, broad, well overlapping. Corona moss green eye for 1/3 its length, changing to pale greenish primrose, with narrow rim of deep red, deeply fluted, almost flat, long style, stiff. Neck grows a little in water.
- MILANION** 1Y-Y
Perianth clear lemon yellow, smooth, stiff. Segments flat, pointed, broad, overlapping. Corona bright, deeper lemon yellow.
- MILK AND CREAM** 2W-Y
Perianth creamy white, creased, segments catch in corona, hooded, floppy. Segments flat, pointed, well overlapping. Corona opens primrose yellow, develops creamy buff. Ugly, badly proportioned.
- MILK AND HONEY** 4W-Y
Perianth segments creamy white, twisted, pointed, separated 2-3 rows. Corona segments open primrose yellow and fade to pale apricot cream, very frilled edges, intermixed loosely with perianth segments.
- MILKMAID** 2W-W
Perianth creamy white, stained primrose at base, pointed outline. Segments flat, pointed, overlapping. Corona primrose yellow, straight sided.
- MILLISLE** 3W-W
Discarded - Hangs head, flowers below foliage.
- MINISTER TALMA** 1Y-Y
Perianth lemon yellow, thin, spidery appearance. Segments twisted, curled back, pointed, separated. Corona deep lemon yellow, very deep lobed edge, broad mouth, badly proportioned.
- MIRABELLE** 3W-YYO
Perianth white. Segments rounded, overlapping and slightly reflexed. Corona small frilled cup, yellow with an orange rim. Neck grows badly in water.
- MIRAGLO** 2Y-O
No description available.
- MISSOURI** 2Y-O
Perianth bright lemon yellow, touch of red at base, paler tips. Segments flat, pointed, overlapping. Corona deep orange red, fluted.
- MISTRAL** 11W-Y
Perianth creamy white. Segments ovate with very little overlap. Corona split, very large, frilled and laid back over yellow perianth segments. Neck grows badly in water.
- MISTY MOON** 3W-WOO
Perianth poeticus white, slightly creased, edges curled in a little, reflexed. Segments flat, rounded, well overlapping. Corona greyish primrose with bright salmon edge for 1/3 of its length and grey green eye, fluted, neck grows in water, corona edge dies first.
- MITE** 6Y-Y
Discarded - Short, small, cyclamineus hybrid.
- MITYLENE** 2W-Y
Perianth creamy white, smooth. Segments flat, pointed, overlapping. Corona primrose yellow develops to pale buff-primrose, sometimes with white blotches on edge, fluted. Neck grows in water.

- MODESTA** 11Y-Y
Discarded - Short, flowers below foliage.
- MOINA** 3W-WWO
Perianth white. Outer segments very broadly ovate, inner ovate, and overlapping. Corona small frilled cup, white with orange rim. Neck grows badly in water.
- MOIRA O'NEILL** 1W-Y
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona pale creamy lemon yellow. Well balanced but small.
- MONACO** 2W-O
Perianth opens pale primrose, develops creamy white. Segments flat, pointed, overlapping. Corona bright mid-orange, paler at base, fluted, bowl shape. Very stiff but brittle neck.
- MONARCH** 1Y-Y
Perianth primrose yellow, creased, crêpey. Segments twisted, pointed, overlapping. Corona lemon yellow, paler at base, long in proportion to perianth.
- MONEYMORE** 2Y-R
No description available.
- MONIQUE** 2W-OOR
Perianth opens primrose, develops cream stained yellow at base, creased. Segments flat, pointed, well overlapping. Corona greenish lemon shading to rim of light orange for 1/4 its length, fluted, shallow bowl shape.
- MONKSTOWN** 2Y
Perianth yellow. Segments overlapping, outer ovate, inner more acute. Corona frilled open mouthed and same yellow as perianth. Neck grows in water.
- MONTE CASSINO** 2W-GRR
Perianth opens creamy primrose with paler tips, develops to creamy white stained yellow at base, creased. Segments flat, pointed, broad, overlapping. Corona brilliant red, with greenish eye, almost flat, deeply fluted. Stiff and sturdy.
- MONTERRICO** 4W-O
Perianth white. Double segments, rounded and overlapping. Corona orange and intermixed with perianth segments. Neck grows badly in water.
- MOON BOY** 2Y-Y
Perianth opens pale lemon yellow, developing cream with deeper edge. Segments flat, pointed overlapping. Corona opens pale lemon yellow, developing cream with narrow edge of lemon yellow, of trumpet proportions.
- MOON GODDESS** 1Y-Y
Perianth opens greenish primrose yellow, developing greenish cream, paler at base. Segments flat, pointed, broad, well overlapping, shovel shape. Corona opens sulphur yellow, developing greenish cream with narrow rim of lemon yellow.
- MOON ORBIT** 1/2Y-Y
Perianth pale yellow, fading to white at base. Segments broadly ovate and overlapping. Corona large flared frilled funnel, white in centre, fading to pale lemon. Corona edge wire rimmed in a slightly deeper yellow.
- MOONDANCE** 3W-W
Perianth creamy white, slightly creased, sometimes notched, edges curl in, circular outline. Segments flat, rounded, overlapping. Corona greyish eye changing to cream with narrow creamy buff edge, slightly fluted, lightly frilled, bowl shape. Hangs head with young. Neck grows in water.
- MOONGATE** 6W-W
No description available.
- MOONGOLD** 1Y-Y
Perianth sulphur yellow, slightly creased, pointed outline. Segments flat, pointed, broad, overlapping. Corona pale lemon yellow, barely frilled, rolled back on itself when fully developed.
- MOONIE** 2W-GWW
No description available.
- MOONMIST** 1Y-Y
Perianth pale yellow. Segments broad, overlapping with slight twisting. Corona open mouthed, frilled, darker yellow than perianth. Weak stems. Neck grows in water.
- MOONRISE** 1Y-Y
Perianth pale greenish sulphur yellow, some green on back. Segments flat, pointed, broad, overlapping. Corona sulphur yellow.
- MOONSHINE** 5W-W
Perianth creamy white. Segments flat, pointed, barely overlapping. Corona creamy white, almost straight sided, corrugated horizontally. Long style.

- MOONSPELL** 2Y-W
Perianth yellow. Segments slightly overlapping, outer are ovate, inner more acute. Corona large frilled and darker yellow than perianth. Neck grows badly in water.
- MOONSTRUCK** 1Y-Y
Perianth opens greenish primrose developing cream. Segments flat, pointed, broad, well overlapping. Corona opens pale lemon yellow, developing cream with narrow edge of pale yellow. Well proportioned.
- MORAY** 1W-W
Perianth white. Segments ovate and overlapping. Corona frilled funnel, creamy white in colour. Neck grows badly in water.
- MOTHER CATHERINE GRULLEMANS** 11W
Discarded - Hooded perianth, short, flowers below foliage.
- MOUNT HOOD** 1W-W
Perianth cream, slightly twisted. Segments edges incurling, some notched, broad, acute. Corona deep cream, large trumpet, flanged, edge raggedly lobed. Neck grows in water.
- MOUNT PLEASANT** 2W-Y
Perianth white, twisted, smooth. Segments broad, ovate, overlapping at base, edges incurling. Corona egg yolk yellow, open mouthed to flat, fluted with lightly frilled edge. Neck grows in water.
- MOUNT TACOMA** 2W
Perianth creamy white, creased, curled in at edges, segments catch in corona, floppy. Segments flat, pointed, barely overlapping. Corona pale apricot gold developing to pale buff primrose, broad and shallow.
- MOUNT VERNON** 2W-P
Perianth white. Segments very broadly ovate and overlapping, with a ridged texture. Corona frilled large cup, white with a pink rim. Neck grows badly in water.
- MOUNTJOY** 7Y-Y
Discarded - Small, flowers below foliage, jonquilla hybrid.
- MOUNTNORRIS** 2W-Y
Perianth white, smooth. Segments broad with blunt tip, overlapping extensively, edges slightly incurling. Corona lemon, flat, ragged, frilled edge. Neck grows badly in water. Weak, hollow stems.
- MOVILLE** 2W-W
Perianth creamy white, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona opens primrose, develops cream with paler edge. Neck grows very badly in water.
- MOYLENA** 2W-P
Discarded - Very susceptible to basal rot.
- MOYLOUGH** 3W-YR
Perianth creamy white, stained yellow at base, a little creased. Segments flat, pointed, overlapping. Corona gold shading to orange red for 3/4 its length, small moss green eye, fluted, almost flat. Neck grows badly in water.
- MR JINKS** 3W-YOO
Perianth white, creased, circular outline. Segments flat, rounded, broad, overlapping. Corona greyish orange eye, changing to buff primrose 2/3 with band of bright apricot orange at edge, fluted, frilled, deeply lobed. Neck grows badly though stiffly in water.
- MRS DAVID CALVERT** 3W-GRR
Perianth white. Segments rounded and overlapping, smooth with curled edges. Corona orange. Flat fluted, lobed and frilled, clean flower. Poor arching neck, grows badly in water.
- MRS E.C.MUDGE** 1W-Y
Perianth creamy white, stained slight yellow at base. Segments flat, pointed, overlapping. Corona clear creamy lemon with slightly paler edge.
- MRS ERNST H.KRELAGE** 1W-W
Perianth creamy white. Segments almost flat, pointed, overlapping. Corona pale creamy lemon, slender, edge rolled back. Neat but small.
- MRS LANGTRY** 3W-W
Perianth creamy white. Segments flat, pointed, separated. Corona opens cream shading to deep cream edge, developing white throughout, fluted, pinched in. Neck grows in water.
- MRS NETTE O'MELVENY** 3W-GYY
Perianth creamy white, some segments creased. Segments flat, pointed, broad, overlapping. Corona bright pale greenish lemon with thin edge of deeper colour, fluted.
- MRS OSCAR RONALDS** 2W-P
Perianth creamy white, slightly creased and crêpey, circular outline. Segments flat, rounded, broad, well overlapping. Corona opens pink, develops cream base 1/3 shading to pale salmon pink with some cream blotches on edge, fluted, frilled, deep bowl shape. Stiff, hangs head when young.

- MRS R.O.BACKHOUSE** 2W-P
Perianth creamy white stained slight apricot at base. Segments twisted, pointed, barely overlapping. Corona apricot primrose shading to salmon pink edge, slender.
- MRS WILLIAM COPELAND** 4W-W
Perianth segments creamy white, flat, pointed, overlapping, outside row complete with some loose pointed segments in centre intermixed with quite large corona segments of pale primrose.
- MULATTO** 1Y-WWY
Perianth pale sulphur yellow, a little creased. Segments flat, pointed, overlapping. Corona pale lemon yellow ageing to paler yellow with rim of lemon yellow.
- MULRANY** 2Y-Y
Perianth bright lemon yellow, edges curl in a little. Segments flat, pointed, broad, well overlapping, shovel shape. Corona deeper lemon yellow, flanged edge, of trumpet proportions. Stem soft and hollow, usually collapses in water.
- MURMANSK** 2W-W
Perianth creamy white, smooth and very flat. Segments flat, pointed, broad, well overlapping, shovel shape. Corona white, fluted, neat. Hangs head badly.
- MUSCADET** 2W-Y
Perianth creamy white, smooth. Segments broad, rounded, overlapping. Corona opens lemon fades to cream, open mouthed, fluted giving frilled rim. Sweetly scented. Neck grows in water.
- MUSIC HALL** 1W-Y
Perianth cream, stained yellow at base. Segments twisted, pointed, broad, overlapping. Corona deep clear yellow.
- MUSTANG** 2Y-YOO
Perianth pale lemon yellow, crêpey substance. Segments flat, rounded, overlapping. Corona pale orange shading to gold base. Rather washy colours.
- MY LOVE** 2W-Y
Perianth creamy white, smooth, edges curl in slightly. Segments flat, pointed, broad, well overlapping, shovel shape. Corona opens primrose develops cream, paler at base, occasional white blotches on edge, lightly frilled, almost straight sided. Trumpet proportion. Hangs head when young. Neck grows in water.
- MY WORD** 2W-P
Perianth cream. Segments notched, edges incurving, acute, overlapping, good star shape. Corona salmon pink, straight, gently lobed. Weak, hollow stems. Neck grows badly in water.
- MYRIANTHA** 3W-WWR
Perianth creamy white, a little creased, circular outline. Segments flat, rounded, well overlapping. Corona greenish primrose, changing to band of bright orange for 1/3 its length, with some white blotches on edge, fluted, almost flat. Neck grows badly in water.
- MYSTERY** 2Y-O
Perianth yellow. Segments broad and overlapping. Corona large frilled cup, orange in colour.
- MYSTIC** 3W-GWO
Perianth greenish-cream, slightly creased. Segments flat, pointed, broad, overlapping. Corona greyish cream, with thin pencil line of apricot orange and a moss-green eye, almost flat, fluted.
- NAIROBI** 2W-O
Perianth opens primrose, quickly develops creamy white stained yellow at base, smooth, edges curl in. Segments flat, pointed, overlapping. Corona bright orange throughout, fluted, frilled, lobed, bowl shape. Stiff.
- NAMPA** 1Y-W
Perianth primrose, paler at centre, slightly twisted. Segments small, overlapping only at base. Corona long and narrow, almost straight, pale primrose at base, becoming darker at rim, gently lobed. Unusual colouring.
- NAMSOS** 2W-W
Perianth creamy white, creased. Segments slightly twisted, broad, well overlapping, shovel shape. Corona opens buff primrose, develops pale cream, broad mouth, finely frilled edge. Hangs head badly when young.
- NANCEGOLLAN** 7W-W
Discarded - Small, flowers below foliage, jonquilla hybrid.
- NANCHISEL** 2Y-O
Discarded - Flowers below foliage, very long neck.
- NANNY NUNN** 3Y-OOR
Perianth pale primrose shading to cream tips, creased, edges curled in. Segments flat, rounded, overlapping. Corona deep clear orange throughout.

- NANTUCKET** 2W-O
Perianth white. Segments broad, overlapping and turned in at edges. Corona large frilled cup and orange in colour. Neck grows badly in water.
- NANTWICH** 2W
Perianth white, stained yellow at base, sometimes notched, circular outline. Segments flat, rounded, broad, well overlapping. Corona gold base, shading to orange red for 1/2 its length, goblet shape. Hangs head badly when young. Neck grows stiffly in water.
- NAOMI** 2W-YYO
Perianth opens primrose, develops to cream stained primrose at base. Segments flat, pointed, broad, overlapping. Corona deep gold with broad edge of deep orange red for 2/3 of its length. Neat and stiff.
- NARVIK** 2Y-O
Perianth light lemon yellow, slightly creased. Segments flat, pointed, overlapping. Corona deep burnt orange red throughout, straight sided, fluted. Neck grows in water.
- NAZARETH** 2Y-W
Perianth primrose with pale primrose at base, slightly twisted. Segments broad, overlapping, fairly rounded. Corona primrose, crenate, unevenly folded, slightly flanged, only lightly frilled. Neck grows in water.
- NEARULA** 3W-R
Discarded - Low flower yield.
- NELLY** 3W-Y
Perianth creamy white, smooth, edges curled in a little. Segments flat, pointed, broad, overlapping. Corona bright greenish lemon with thin pencil edge of light apricot and greenish eye, fluted. Neck grows in water.
- NEMO** 3W-YYO
Perianth opens primrose, develops cream, edges curl in, creased. Segments flat, pointed, broad, well overlapping. Corona pale lemon changing to band of orange red for 1/3 its length, deeply fluted, almost flat, lobed edge. Neck grows in water.
- NEVANA** 2W-P
No description available.
- NEW GOLD** 2Y-Y
Perianth sulphur yellow, creased a little. Segments flat, pointed, broad, well overlapping. Corona bright deep lemon yellow, suffused gold. Stiff. Neck grows in water.
- NEW HORIZON** 1Y-Y
Perianth opens primrose, developing cream, crêpey texture, pointed outline. Segments slightly twisted, pointed, overlapping. Corona opens creamy lemon yellow developing cream with rolled back edge.
- NEWCASTLE** 1W-Y
Perianth white. Segments overlapping, outer broad, inner more ovate. Corona frilled, flared cylinder, yellow in colour.
- NEWTON** 8Y-Y
Discarded - Better suited to Isles of Scilly.
- NIMROD** 2Y-Y
Perianth pale lemon, slightly creased, edges curl in a little, pointed outline. Segments flat, pointed, overlapping. Corona gold, fluted, frilled, deep bowl shape.
- NINTH LANCER** 2Y-O
Perianth bright lemon yellow, creased, notched, pointed outline. Segments flat, pointed, overlapping. Corona deep orange red throughout, slightly fluted, deep bowl shape. Neck grows badly in water.
- NIPHETOS** 2W-W
Perianth creamy white, slightly creased, perfect double triangle. Segments flat, pointed, broad, overlapping. Corona opens soft lemon yellow, develops to pale cream.
- NIRVANA** 7W-W
Perianth creamy-white, curled in at edges. Segments flat, rounded, overlapping. Corona opens creamy sulphur, develops to cream. Neck grows in water.
- NISSA** 2W-Y
Perianth creamy white, slightly creased. Segments flat, pointed, broad, overlapping. Corona deep bright lemon yellow. Neat but small.
- NIVETH** 5W-W
Perianth white. Segments flat, pointed, barely overlapping. Corona white (whiter than perianth) deep bowl shape. Long style. Neck grows in water.
- NOBLE DIAMOND** 3W-GYR
Perianth opens cream, develops white stained yellow at base, creased. Segments flat, rounded, overlapping. Corona deep sealing wax red shading to slight greenish gold eye, fluted.

NOBLESSE Discarded - Rough flower, poor colour.	2Y-O
NODDY Discarded - Short, susceptible to basal rot.	2W
NOR-NOR Perianth opens creamy sulphur yellow, ages to creamy white, stained yellow at base for 1/2 length of segments, crêpey texture. Segments slightly twisted, pointed, barely overlapping. Corona opens orange gold, develops to deep orange, slightly waisted.	2Y-Y
NORTH CAPE Perianth creamy white, slightly creased, one segment sometimes buckled. Segments flat, pointed, broad, well overlapping. Corona opens primrose, develops buff and ages to cream with deeper edge, deep bowl shape.	2W-W
NORTHERN LIGHT Perianth white. Segments broad and overlapping. Corona medium sized cup, frilled orange, slightly paler in centre. Neck grows badly in water.	2W-O
NORVAL Perianth white, stained yellow base, double triangle outline. Segments rounded, overlapping, bottom segments much broader, creased, sometimes buckled, curled in at edges. Corona rich bright orange, bowl-shaped, fluted, deeply lobed edge.	2W-O
NORWAY Perianth pale yellow. Segments broad and overlapping and slightly inflexed. Corona large open mouthed, frilled trumpet and yellow in colour.	1W-Y
NUAGE Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona opens primrose, develops creamy white, smooth, rolled back edge, of trumpet proportions.	2W-W
NURSEMAID Perianth primrose yellow with white tips, creased. Segments flat, rounded, overlapping. Corona deep orange, a little paler at base, deeply fluted.	3W-O
NYORA Perianth white. Segments very broadly ovate and overlapping. Corona small, frilled cup, yellow in centre with orange rim. Neck grows badly in water.	3W-YYR
N.CYPRI No description available.	10
N. JONQUILLA Perianth brilliant lemon yellow. Segments flat, pointed, separated. Corona deeper lemon yellow, smooth.	10Y-Y
N. OBVALLARIS Perianth deep bright gold. Segments flat, pointed, broad, overlapping. Corona deeper gold a little paler at base, deeply lobed. Squat looking flower.	10
N. OCHROLEUCUS (syn. TAZETTA OCHROLEUCUS) Discarded - better suited to Isles of Scilly.	10
N. POETICUS POETICUS RECURVUS Perianth pure white, stained slight yellow at base, edges curled in a little, reflexed. Segments flat, pointed, overlapping. Corona greenish lemon changing to thin rim of deep red, with deep moss green eye. Very stiff and sturdy.	10W-YYR
N. POETICUS RADIIFLORUS POETARUM Perianth white stained slight yellow at base, creased, edges curled in. Segments twisted, pointed, separated. Corona bright orange red, a little deeper at edge, with small greenish eye, finely fluted.	10W-GYR
N. PSEUDONARCISSUS GAYI Perianth pale primrose, crêpey texture. Segments twisted, pointed, separated. Corona pale lemon yellow, paler at base, very slender trumpet. Soft stem.	10Y-Y
N. TAZETTA LACTICOLOR (syn. CANALICULATUS) Perianth white, star shaped. Segments flat, pointed, barely overlapping. Corona deep gold, smooth, goblet shape.	10W-Y
N. x MEDIOLUTEUS (syn. BIFLORUS) Perianth creamy white, stained yellow in base. Segments broad, well rounded, well overlapping, creased, buckled, curled in at edges. Corona citrus yellow, fine silver line near narrow rim, expanded, frilled edge. Unattractive scent.	10W-Y

OCARINO	4Y-Y
Perianth pale primrose darkening at edges. Segments smooth, rounded, overlapping at base, curled in at edges; one and a half complete rows with three or more segments pointed forwards and intermixed with bright orange-yellow Corona segments, open centre, frilled edge. Weak stems. Neck grows in water.	
OCEAN MIST	1W-W
Perianth white. Segments broad and overlapping. Corona large flared, frilled trumpet, white in colour. Neck grows in water.	
ODENSE	2W-YYO
Perianth poeticus white, smooth, some segments reflexed. Segments broad, acute, overlapping. Corona opens sulphur becomes pale orange streaked with orange, still sulphur at edge, fluted, open mouthed to flat giving lobed, frilled edge. Neck grows badly in water.	
OKLAHOMA	1W-Y
Perianth creamy white, stained yellow base, creased. Segments slightly twisted, pointed, overlapping. Corona deep chrome yellow, deeply crenate, flanged, too large in relation to perianth. Neck grows in water.	
OLDERFLEET	2W-P
No description available.	
OLIVER CROMWELL	1W-Y
Perianth creamy white stained yellow at base. Segments twisted, pointed, separated, giving thin spidery appearance. Corona bright clear lemon yellow, too large for size of perianth.	
OLIVET	2W-W
Perianth white. Segments broad and overlapping, slightly inflexed. Corona frilled large cup, same white as perianth. Weak stems. Neck grows in water.	
OLYMPIA	1Y-Y
Perianth primrose yellow, creased, crêpey, often caught in corona. Segments twisted, pointed, overlapping. Corona lemon yellow, deep cut lobes, untidy edge. Badly proportioned.	
OLYMPIC FIRE	2Y-R
Discarded - Small.	
OLYMPIC GOLD	1Y-Y
Perianth bright lemon, double-triangle outline. Segments rounded, overlapping at base, bottom segments broader, top segments tending to twist. Corona slightly darker, straight, fluted, crenate edges. Flower substance very good. Scented.	
OLYMPIC TORCH	2W-YYO
Discarded - Small, short, untidy corona.	
OMAROE	2W-Y
Perianth white. Segments broad and overlapping with ridged texture. Corona large frilled cup, yellow in colour. Neck grows in water.	
ORANGE BELL	2Y-O
Perianth sulphur yellow, creased, notched, one segment often caught in corona. Segments flat, pointed, broad, well overlapping. Corona orange throughout often with white blotches on edge, untidy deep cut lobed edge, rough. Neck grows badly in water.	
ORANGE BIRD	2Y-R
Perianth pale sulphur yellow, creased, crêpey texture. Segments flat, pointed, overlapping. Corona bright orange red, fluted. Neck grows in water.	
ORANGE COCKADE	3W-O
Discarded - Poor colour.	
ORANGE CREST	2Y-YYO
Perianth yellow. Segments broad, overlapping with ridged texture. Corona large frilled cup, yellow centre with orange rim. Neck grows badly in water.	
ORANGE CUP	8Y-YYR
Perianth creamy primrose, creased, edges curl in. Segments flat, pointed, overlapping. Corona green eye shading to orange with deep orange edge, often with cream blotches, finely cut edge, untidy.	
ORANGE FAVOURITE	2
Perianth sulphur yellow, creased. Segments flat, pointed, overlapping. Corona bright orange red, neatly frilled edge, bowl shape. Neck grows in water.	
ORANGE FIRE	2W-O
Discarded - Short, low flower yield.	
ORANGE FLAG	2W-O
Discarded - Poor shape.	

- ORANGE FRILLED** 2Y-O
Perianth sulphur yellow, creased, top three segments incurved. Segments flat, pointed, well overlapping. Corona gold base, shading to orange edge, some white blotches. Edge so heavily and broadly frilled as to make the flower top heavy.
- ORANGE GLORY** 6Y-O
Perianth soft lemon yellow, slightly reflexed. Segments twisted, pointed, separated. Corona deep orange gold shading to slightly paler base, waisted.
- ORANGE GLOW** 2Y-O
Perianth pale sulphur yellow, star shaped, floppy. Segments flat, pointed, barely overlapping. Corona pale golden orange shading to gold 1/2 way down, deeply frilled.
- ORANGE MASTER** 2Y-O
Perianth pale lemon yellow, crêpey texture, creased. Segments slightly twisted, pointed, overlapping. Corona orange, a little paler at base, deep cut lobed edge, almost trumpet length. Rough. Neck grows badly in water.
- ORANGE MONARCH** 2W-YOO
Perianth creamy white, stained yellow base, creased, often buckled. Segments flat, rounded, well overlapping. Corona bright solid orange throughout. Some white blotches on edge, fluted, shallow. Neck grows in water.
- ORANGE QUEEN** 7Y-Y
Perianth golden orange, star shaped. Segments twisted, pointed, barely overlapping. Corona, deeper golden orange.
- ORANGE SHERBERT** 2Y-YOY
Perianth creamy white, stained at base. Segments pointed, creased, curled in at edges and overlapping. Corona pale orange, yellow wire edge with a shallow bowl shape, fluted frilled small flower.
- ORANGE STANDARD** 2Y-O
Perianth deep lemon overlaid gold, slightly creased, star shape. Segments flat, pointed, barely overlapping. Corona dark orange red, fluted, frilled almost straight sided. Floppy. Neck grows in water.
- ORANGERY** 11W-POY
Perianth white. Segments rounded and overlapping, crêpey. Corona split and some segments interspersed with perianth segments. Corona orange in colour. Neck grows in water.
- ORANIQUE** 2W-YYO
Perianth creamy white, a little creased, often caught in corona. Segments flat, pointed, overlapping. Corona bright apricot orange with narrow rim of apricot, fluted, shallow bowl shape. Neck grows badly in water.
- ORANJE BRUID** 2W-GOO
Perianth poeticus white, sometimes caught in corona, circular shape. Segments flat, rounded, well overlapping. Corona brilliant orange with paler edge and greyish green eye, almost flat, double frilled.
- ORANJE VAAN** 2Y-O
Perianth primrose yellow, creased. Segments flat, pointed, overlapping. Corona orange-red throughout, some white blotches on edge. Neck grows in water.
- ORATORIO** 2W-Y
Perianth cream, creased, outer 3 segments reflexed. Segments broad, rounded, incurving edges, overlapping. Corona opens pale lemon, fades at base to cream, straight sided, wider at mouth, slightly fluted, rim deeply lobed. Neck grows in water.
- ORIENTAL RAY** 2Y-YYO
Perianth yellow. Segments broad, overlapping and slightly inflexed. Corona large frilled cup, yellow with orange rim. Neck grows in water.
- ORISSA** 3Y-YYO
Perianth sulphur yellow, deeply creased, often buckled. Segments flat, pointed, broad, overlapping. Corona greenish gold shading to bright orange red for 1/2 its length, deeply fluted, shallow bowl shape.
- ORMEAU** 2Y-Y
Perianth lemon, slightly creased and crêpey. Segments flat, pointed, broad, overlapping. Corona deeper lemon, lightly frilled, almost straight sided. Trumpet proportion. Stiff.
- ORNATUS (syn. POETICUS ORNATUS)** 10W-YYR
Perianth white, creased, buckled. Segments flat, pointed, separated. Corona pale greenish primrose with narrow edge of dark red, fluted, flat. Short style half way between long and short stamens. Neck grows in water.
- ORNATUS MAXIMUS** 9W-GYR
Perianth white stained slightly yellow at base, reflexed, circular outline. Segments flat, rounded, overlapping. Corona moss green eye changing to pale greenish lemon with narrow rim of deep red, fluted, almost flat. Style long and protruding. Neck grows in water.
- ORONTES** 2Y-O
Perianth light lemon yellow, developing pale tips, slightly creased, circular outline, a little reflexed. Segments flat, rounded, overlapping. Corona bright orange red, a little paler at base, fluted, bowl shape.

ORPHEUS 2W-PPY
 Perianth white. Segments very broadly ovate and overlapping. Corona medium sized frilled cup, with yellow eye and pink edges.

OSIRIS 1Y-Y
 Perianth rich mid-gold, creased, crepey. Segments very twisted, pointed, overlapping at base. Corona darker, bright gold, expanded with wide belled mouth, deeply crenate edge. Neck grows in water.

OSLO 2W-W
 Perianth creamy white, creased, edges curl in. Segments flat, pointed, well overlapping. Corona cream, fluted, deep bowl shape. Hangs head badly.

OSPREY 2W-Y
 Perianth pure white, creased, floppy. Segments twisted, pointed, separated. Corona pale apricot lemon, shading to deeper apricot, lemon edge, with greenish lemon eye.

OSWEGA 2W-Y
 Discarded - Long neck, poor poise.

OTRANTO 3W-YYO
 Perianth white. Outer segments rounded, inner very broadly ovate, overlapping. Corona small frilled cup, yellow with orange rim. Neck grows badly in water.

PAINTED LADY 2W-O
 Perianth opens cream, develops creamy white stained yellow at base. Segments flat, rounded, well overlapping. Corona brilliant solid red. Neat and stiff.

PALAVERER 1W-Y
 Perianth white, slightly creased, slightly twisted. Segments would fit into a double triangle, acute, some notched. Corona lemon becoming very pale at edge, narrow trumpet, slightly flanged, frilled. Neck grows stiffly in water.

PALESTINE 2Y-YOO
 Perianth bright lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona orange shading to deep orange red edge, straight sided, fluted. Very stiff. Neck grows stiffly in water.

PALMY DAYS 2Y-YOO
 Perianth yellow. Segments ovate, very little overlap and twisting. Corona medium sized, frilled cup, orange with yellow centre. Neck grows badly in water.

PAPILLON BLANC 11W-W
 Perianth creamy white, creased. Segments flat, rounded, well overlapping. Corona opens primrose developing cream with thin pencil edge of apricot orange and a grey-green eye, split evenly into six overlapping segments. Neck grows in water.

PAPRIKA 3W-R
 Perianth creamy white, stained yellow base, creased, edges curl in a little. Segments flat, pointed, overlapping. Corona bright orange throughout, fluted, frilled, lobed, shallow bowl shape. Neck grows stiffly in water.

PAPUA 4Y-Y
 Perianth canary yellow, paler at tips. Segments round and 1/2 overlapping, slightly crêpey and curled in at edges. Segments intermixed with pale orange frilled corona segments. Neck grows badly in water.

PARACUTIN 2Y-R
 Perianth yellow. Segments broad, overlapping and inflexed. Corona medium sized, frilled orange trumpet. Neck grows in water.

PARAMARIBO 2Y-R
 Perianth lemon yellow, creased, sometimes notched. Segments flat, pointed, overlapping. Corona deep solid orange red, some white blotches on edge, deeply fluted and frilled, deep bowl shape. Neck grows in water.

PARCPAT 7Y-O
 Perianth opens lemon yellow, quickly fades to pale primrose, stained yellow at base, smooth. Segments flat, rounded, overlapping. Corona golden orange. Neck grows in water.

PARISIENNE 11W-O
 Discarded - Short, flowers below foliage, rough.

PARKMORE 2W-W
 Perianth creamy white, slightly creased, crêpey texture. Segments flat, pointed, broad, well overlapping, shovel shape. Corona pale cream, of trumpet proportions.

PARSIFAL 2W-R
 Perianth creamy white, stained yellow base. Segments rounded, overlapping at base, rolled in at edges to give pointed effect. Corona deep rich orange with faint reddish tinge, shallow, bowl-shaped, fluted, deeply lobed edge.

PASSIONALE	2W-P
Perianth white, smooth. Segments broad, rounded, overlapping only slightly. Corona pale pink, open mouthed, gently wavy edge.	
PATAGONIA	2Y-O
Perianth deep lemon, smooth. Segments rounded but with edges incurling, would fit into a double triangle, overlapping at base. Corona deep orange, short, open mouthed, ribbed, ragged, lobed edge. Neck grows badly in water.	
PATHAN	2Y-O
Perianth sulphur yellow, developing paler at tips, creased, edges curl in. Segments flat, pointed, overlapping. Corona deep burnt orange red, deeply fluted, bowl shape. Neck grows in water.	
PATRIA	1W-Y
Perianth creamy white, stained yellow at base and on the back, crêpey texture. Segments slightly twisted, pointed, overlapping. Corona pale lemon yellow.	
PATRIARCH	2W-O
Perianth creamy white, slightly creased, edges notched. Segments flat, rounded, well overlapping. Corona deep apricot with paler edge and centre, sometimes white blotches on edge.	
PATRICIA	4W-R
No description available.	
PAUL BUNYON	1Y-Y
Discarded - Poor colour, badly proportioned.	
PEARL SCHUBERT	2W-Y
Perianth creamy white. Segments broad and overlapping. Corona medium sized, frilled, open mouthed and pale yellow in colour.	
PEDRO	1Y-Y
Perianth deep clear greenish yellow, green on back. Segments twisted, pointed, overlapping. Corona deep golden yellow. Too small for height.	
PEEPING TOM	6Y-Y
Perianth clear lemon yellow, crêpey texture, reflexed. Segments flat, pointed, barely overlapping. Corona deep solid gold, very long and slender with flanged edge.	
PEERLESS GOLD	1Y-Y
Discarded - Badly proportioned.	
PEKING	1Y-Y
Perianth cream, a little green and yellow on back, slightly creased. Segments, flat, pointed, broad, overlapping. Corona creamy lemon yellow, open-mouthed. Squat and too large for stem length.	
PEMBRAZE	1Y-Y
Perianth bright clear lemon yellow, creased. Segments slightly twisted, pointed, broad, overlapping. Corona deeper lemon yellow, almost straight with untidy uneven frilled edge.	
PENBEAGLE	2Y-Y
Perianth clear lemon yellow, slightly creased. Segments flat, pointed, broad, overlapping. Corona bright, deep lemon yellow, flanged.	
PENBEARTH	1Y-Y
Perianth yellow. Segments broad and overlapping. Corona large open mouthed, frilled, same yellow as perianth. Neck grows badly in water.	
PENCOYS	2Y-YOO
Perianth pale creamy primrose, deep primrose at base, crêpey texture. Segments flat, rounded, overlapping. Corona gold, shading to burnt orange for 1/2 its length.	
PENCREBAR	4Y-Y
Perianth bright lemon yellow, crêpey texture, flat, pointed, overlapping, two or three complete rows, continuing in centre, evenly intermixed with deeper lemon yellow corona segments. Sweet jonquil scent.	
PENGORNIN	2W-Y
Perianth creamy white. Segments broad and overlapping. Corona large frilled cup, pale yellow in colour. Neck grows badly in water.	
PENPOL	7Y-Y
Perianth soft lemon yellow, creased, inclined to reflex. Segments flat, pointed, broad, overlapping. Corona deeper lemon yellow. Neck grows in water.	
PENQUITE	2Y-O
Perianth soft lemon yellow, segments curled in at edges, crêpey texture. Segments flat, rounded, overlapping. Corona orange, fluted, bowl shaped. Stiff and sturdy.	

PENRYN 2W-W
 Perianth creamy white, slightly creased and crêpey, top segments incurve slightly, edges curl in, circular outline. Segments flat, rounded, broad, well overlapping. Corona opens buff cream with moss green eye, develops cream with grey green eye, fluted, frilled, deeply lobed, hangs head when young. Neck grows in water.

PENSIVE 2W-P
 Discarded - Poor colour, flowers below foliage, susceptible to basal rot.

PENTEWAN (?syn. CALIFORNIA) 2Y-Y
 Perianth lemon, slightly twisted. Segments acute, overlapping to 1/2 their length. Corona deep lemon, long, straight sided then flanged, deeply indented edge. Well balanced flower. Neck grows in water.

PENTREATH 3Y-O
 Perianth opens lemon yellow, develops white tips, creased. Segments slightly twisted, pointed, overlapping. Corona bright orange, fluted, bowl shape.

PENVER 2Y
 Discarded - Floppy, flowers below foliage, susceptible to basal rot.

PENVOSE 2Y-Y
 Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona opens deep lemon yellow, develops deep buff-orange, of trumpet proportions. Stiff and sturdy.

PEPPER 2Y-O
 Perianth bright lemon yellow, pointed, star shaped. Segments slightly twisted, pointed, separated. Corona bright orange-red fluted.

PER LAGERQUIST 2Y-YYO
 Perianth yellow. Segments very broadly ovate, overlapping with a ridged texture. Corona medium sized frilled cup, yellow with orange rim. Neck grows badly in water.

PERA 3W-R
 Perianth cream white, smooth, inclined to reflex. Segments flat, pointed, broad, well overlapping. Corona sealing wax red with greenish gold eye, fluted, nearly flat, lobed edge. Neck grows stiffly in water.

PERDITA 9W-GYR
 Perianth poeticus white. Segments rounded and overlapping, slightly reflexed. Corona small frilled cup, green eye, yellow segments with an orange/red rim. Neck grows badly in water.

PERICLES 1W-W
 Discarded - Short, hangs head.

PERIDOT 2W-GYY
 Perianth white. Segments broad and overlapping. Corona small frilled cup, green eye, very pale yellow segments fading to white. Neck grows badly in water.

PERIMETER 3Y-YYO
 Perianth yellow. Segments very broadly ovate and overlapping. Corona small frilled yellow cup with orange rim. Neck grows badly in water.

PERSONALITY 2W-Y
 Discarded - Hangs head, very susceptible to basal rot.

PERSUASION 2W-OOY
 Perianth white. Segments broad and overlapping. Corona medium sized, frilled cup, yellow with orange centre. Neck grows badly in water.

PERU 2Y-O
 Perianth yellow. Segments broad and overlapping, slightly ridged texture. Corona medium sized orange frilled cup. Neck grows in water.

PETER PIPER 2Y-O
 Perianth pale lemon yellow, creased. Segments flat, pointed, overlapping. Corona bright orange red throughout, fluted, bowl shape. Neck grows in water.

PETERBOROUGH 2Y-Y
 Perianth light lemon yellow, sometimes catches in corona, creased. Segments flat, rounded, well overlapping. Corona gold base shading to pale orange edge for 1/3 its length, fluted, deeply frilled. Neck grows in water.

PETILLANT 3W-YYO
 Discarded - Small, poor perianth, flowers below foliage.

PETSAMO 1W-W
 Discarded - Perianth catches in corona, hangs head.

PHILALETUS 2W-WWY
 Perianth creamy white, notched, creased, circular outline. Segments flat, rounded, well overlapping. Corona cream with narrow rim of lemon yellow, developing to pale apricot, fluted. Neck grows in water.

- PHILLUS CHIDGEY** 2Y-Y
Perianth yellow. Outer segments very broadly ovate, inner more blunt and overlapping. Corona frilled, funnel shape, deeper yellow than perianth. Neck grows in water.
- PICADOR** 3W-GOO
Perianth creamy white, stained yellow at base, smooth. Segments flat, rounded, well overlapping. Corona orange gold shading to orange red edge for 2/3 of its length and moss green eye, almost flat, fluted. Neat and stiff.
- PICK ME** 2Y-O
Perianth yellow. Segments ovate and overlapping. Corona medium sized, frilled cup, orange in colour. Neck grows badly in water.
- PICK UP** 11Y-O
Perianth white. Outer segments rounded, inner very broadly ovate. Corona is split, frilled and orange in colour, a segment of which is attached to each perianth segment. Neck grows badly in water.
- PICKWICK** 2W-YYO
Perianth poeticus white, slightly twisted, smooth. Segments broad, rounded with prominent mucronate tip, well overlapping. Corona lemon at base with pale orange band at edge, short, open mouthed, creased, slightly frilled at edge. Neck grows badly in water.
- PIERROT** 3W-R
Perianth white, stained yellow base, creased, edges curl in at tips, reflexed, pointed outline. Segments flat, pointed, overlapping. Corona greenish gold base 1/4 shading to bright orange, slightly fluted, lightly frilled, bowl shape. Neck grows in water.
- PIGEON** 2W-W
Perianth white, circular outline. Segments flat, rounded, broad, well overlapping. Corona cream with grey eye, fluted, bowl shape. Neat and stiff.
- PILGRIM** 2W-Y
Perianth opens primrose, develops cream stained yellow base, edges curl in, star shape. Segments flat, pointed, separated. Corona pale greenish lemon, fluted, pinched in. Neck grows badly in water.
- PILGRIMAGE** 2Y-Y
Perianth clear sulphur yellow, edges curl in a little, smooth. Segments flat, pointed, broad, well overlapping. Corona bright lemon yellow.
- PILOT LIGHT** 2Y-O
Perianth sulphur shading paler at tips, creased, crêpey, sometimes buckled. Segments flat, pointed, overlapping, edges curl in. Corona deep orange red, paler at base, deeply fluted, frilled, deeply cut lobed edge, bowl shape. Poor poise. Neck grows badly in water.
- PINDAR** 3W-GWP
Perianth poeticus white, edges curl in. Segments flat, pointed, well overlapping. Corona moss green eye shading to greenish cream with pencil rim of deep red, fluted, almost flat. Neck grows badly but stiffly in water.
- PINK BEAUTY** 2W-P
Perianth creamy white, deeply creased, pointed outline, sometimes caught in corona. Segments flat, pointed, overlapping. Corona bright pink developing paler, slender. Rough. Neck grows in water.
- PINK CHIFFON** 4W-P
Perianth creamy white, stained yellow at back and base. Segments round, overlapping with creased centre line. Segments intermixed with corona segments. Corona untidy and sometimes a mix of undifferentiated anther and perianth segments. Corona rose pink, with a lemon frilled edge. Neck grows badly in water.
- PINK FANCY** 2W-P
Perianth creamy white, creased, stiff. Segments slightly twisted, pointed, overlapping. Corona apricot gold shading to deep salmon edge for 1/5 of its length, developing paler with age.
- PINK GIN** 2W-GPP
Perianth white. Segments broad and overlapping, inner whorl more ovate. Corona medium sized, frilled cup, pink, with yellow centre. Weak stems. Neck grows badly in water.
- PINK GLORY** 2W-P
Perianth creamy white, slight pink at base. Segments twisted, pointed, barely overlapping. Corona pale peach shading to creamy peach base, developing creamy buff with age.
- PINK ISLE** 2W-P
Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona bright pure pink, shade paler at base, ageing a little paler, some white blotches on edge, straight sided.
- PINK PACER** 2W-P
Perianth pale cream, slightly hooded, head hangs down. Segments overlapping at base only. Corona very pale pink at base, becoming pale pink at rim, colour fades to cream, straight sided with slightly frilled rim. Neck grows in water.

- PINK PROMISE** 2W-P
Perianth white. Segments ovate and overlapping. Corona medium sized, frilled cup, pastel pink in colour. Neck grows in water.
- PINK RIM** 2W-YYP
Perianth creamy white, slightly creased. Segments flat, rounded, overlapping. Corona pale buff primrose shading to thin primrose line, with narrow salmon pink edge.
- PINK SELECT** 2W-OOP
Perianth creamy white, creased, deeply notched, often caught in corona. Segments flat, pointed, overlapping. Corona opens primrose, develops peach, paler at base, often split in two or three places. Badly proportioned, ugly.
- PINK SMILES** 2W-P
Perianth creamy white, slightly creased, sometimes notched. Segments flat, pointed, overlapping. Corona opens primrose, develops primrose flushed salmon with some white blotches on edge, frilled, lobed. Trumpet proportion.
- PINK SPRITE** 2W-P
Discarded - Small, short, flowers below foliage.
- PINKEEN** 2W-P
Perianth creamy white, double triangle. Segments flat, rounded, well overlapping. Corona opens primrose yellow, develops buff primrose with white blotches on the edge, almost straight sided.
- PINOCCHIO** 2W-GOO
Perianth pure white, creased, double triangle, circular shape. Segments flat, rounded, well overlapping. Corona opens primrose yellow, develops to apricot with greenish eye, later ageing to bright orange gold - varying from season to season, double frilled. Stiff and sturdy.
- PINWHEEL** 2Y-Y
Discarded - Poor colour, badly proportioned, soft stem.
- PINZA** 2Y-YOR
Perianth lemon, smooth. Segments rounded and overlapping, slightly reflexed. Corona orange becoming more intense at edge, straight sided, small frills at edge. Neck grows very badly in water.
- PIONEER** 1W-Y
Perianth white. Segments ovate, twisted, with slight overlapping. Corona large flared, frilled and bright yellow. Neck grows in water.
- PIRATE KING** 2W-O
Perianth white becoming cream at base, smooth, edges slightly incurled. Segments broad, slightly rounded, overlapping to half of length. Corona orange, short, open mouthed, slightly lobed at rim. Neck grows in water. Perianth catches in corona. Neck grows in water.
- PIXIE'S PARLOUR** 3W-Y
Perianth poeticus white, slightly twisted. Segments edges incurling, rounded and overlapping. Corona cream, small and short, slightly flared, deeply notched rim, slightly frilled. Neck grows badly in water.
- PLANET** 4Y-Y
Perianth clear lemon. Segments pointed, two outer whorls distinct, overlapping, inner Segments becoming progressively smaller and less distinct and intermixed with irregular gold corona segments. Weak stems. Neck grows badly in water.
- PLANTINAL** 3W-GWW
Discarded - Small.
- PLAYBOY** 2Y-O
Discarded - Poor colour.
- PLEASANT** 2Y-R
Perianth deep clear yellow, slightly creased, crêpey texture. Segments flat, pointed, broad, overlapping. Corona orange shading to gold base. Stiff.
- PLEASING** 2Y-Y
Perianth lemon yellow, perfect double triangle, smooth, nice sheen. Segments flat, rounded, well overlapping. Corona deeper lemon yellow, neat edge, well proportioned.
- PLENUS (syn. POETICUS FLORE PLENO)** 4W-W
Perianth segments pure white, flat, pointed, broad, overlapping, three complete rows with some narrower pointed segments in centre. Neat and symmetrical.
- PLUVIUS** 2Y-O
Perianth sulphur yellow, paler at tips, creased, crêpey texture. Segments twisted, pointed, overlapping. Corona light orange throughout, long and slender, deep cut, untidy edge. Untidy, ugly flower. Neck grows in water.

POINT BARROW 1W-W
Discarded - Short, flowers below foliage, susceptible to basal rot.

POLAR ICE 3W-W
Perianth poeticus white, edges curled in, slightly reflexed. Segments flat, pointed, overlapping. Corona white with a faint line of buff on edge and a bright green eye.

POLAR SEA 3W-GWW
Perianth poeticus white, smooth, star shape. Segments flat, pointed, separated. Corona bright green, changing to pure white for 3/4 its length, fluted, almost flat. Neck grows stiffly in water.

POLGLASE 8W-R
Perianth white, stained yellow at base, creased, curled in at edges, crêpey texture. Segments flat, pointed, overlapping. Corona deep red. Style longer than longest stamens. Neck grows in water. Rough looking flower.

POLINDRA 2W-Y
Perianth creamy white, perfect double triangle. Segments flat, pointed, broad, overlapping, shovel shaped. Corona lemon yellow developing to deep cream. Well proportioned.

POLNESK 7Y-Y
Perianth pale greenish lemon, very smooth. Segments flat, pointed, overlapping. Corona bright lemon yellow, well balanced. Neck grows in water.

POMONA 3W-GYO
Perianth poeticus white, a little creased, reflexed. Segments flat, rounded, well overlapping. Corona greenish yellow with bright orange edge for 1/4 of its length.

PONDEROSA 1Y-Y
Perianth lemon, slightly creased. Segments acute, forming a double triangle, some notched. Corona deep lemon, flanged, deeply lobed, slightly frilled edge. Well balanced but small flower. Neck grows in water.

PONTRESINA 2W-Y
Perianth white. Segments rounded and overlapping. Corona large frilled bowl, creamy white. Neck grows badly in water.

PORTADOWN 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona large frilled funnel shape, same yellow as perianth. Neck grows in water.

PORTAL 2W-WPP
Perianth creamy white. Segments broad and overlapping. Corona large frilled cup, pink, with yellow centre. Neck grows in water.

PORTARLINGTON 1Y-Y
Perianth lemon yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona deeper lemon yellow, neat.

PORTHCHAPEL 7Y-O
Perianth yellow. Segments ovate and slightly overlapping. Corona medium sized frilled cup, orange in colour. Neck grows badly in water.

PORTHGWARRA 2W-Y
Perianth white. Segments broad and 1/2 overlapping, slightly twisted and inflexed. Corona large frilled, flared, and yellow in colour. Neck grows in water.

PORTHILLY 2Y-O
Perianth pale lemon, white tips and lighter streak down centre of segments, creased. Segments flat, pointed, broad, overlapping. Corona bright orange red, fluted. Neck grows in water.

PORTMARNOCK 2Y-Y
Perianth lemon, creased, notched, sometimes buckled, pointed outline. Segments flat, pointed, overlapping. Corona deeper lemon, uneven deep cut edge, broad bowl shape. Neck grows in water.

PORTRUSH 3W-W
Perianth greenish creamy white, smooth, circular outline. Segments flat, rounded, broad, well overlapping. Corona moss green eye, changing to cream for 2/3 its length, pinched in, deeply fluted. Very stiff. Neck grows in water.

POSTMISTRESS 2W-R
Perianth creamy white, smooth. Segments broad, rounded, overlapping. Corona orange, short, cup shaped, fluted with slightly frilled rim. Neck grows badly in water.

PREAMBLE 1W-Y
Perianth white, a little creased. Segments flat, pointed, broad, overlapping, shovel shape. Corona opens chrome yellow, develops creamy primrose with deeper rim, lightly frilled. Hangs head when young.

PRECENTOR 1Y-Y
Perianth yellow. Segments ovate 1/2 overlapping, and twisted. Corona large, flared, frilled cylinder, same yellow as perianth. Neck grows in water.

PREFIX 6Y-Y
Perianth yellow. Segments acute, slight overlap and reflexed. Corona large flared, frilled cylinder, same yellow as perianth.

PRESIDENT LEBRUN 1W
Perianth creamy white, stained cream at base. Segments rounded, broad, overlapping. Corona soft buff yellow, very wide mouth, giving squat appearance.

PRESTIGE 1W-W
Discarded - Corona often splits, hangs head, susceptible to basal rot.

PRETORIA 1Y-Y
Perianth deep clear yellow, smooth. Segments flat, pointed, broad, well overlapping. Corona deeper lemon yellow. Neck grows in water.

PRIDE OF CORNWALL 8W-YYR
Perianth creamy white stained slightly yellow at base, creased and buckled. Segments flat, pointed, overlapping. Corona bright red, shading to slight gold base, almost flat. Style reaches almost as long as longest stamens. Neck grows in water.

PRIDE OF ERIN 3W-YYR
Perianth creamy white, stained yellow at base, edges curl in, creased, top three segments incurve. Segments flat, rounded, broad, overlapping. Corona greenish lemon, changing to bright orange for 1/2 its length, deeply fluted. Neck grows badly in water.

PRIMROSE BEAUTY 4Y
Perianth segments primrose yellow, smooth, flat, rounded, overlapping, complete. Corona deeper yellow with pencil line of orange, complete, centre filled in with twisted corona and perianth segments. Neat.

PRIMROSE PHOENIX 4Y-Y
Perianth segments cream, stained yellow at base, flat, pointed, overlapping, complete outer row. Corona almost complete, orange red filled with twisted perianth and corona segments. Neck grows in water, ugly.

PRINCE REGENT 2Y-O
Perianth bright clear lemon yellow, creased, edges curl in. Segments flat, pointed, overlapping. Corona bright orange, fluted, deeply lobed edge, bowl shape. Neck grows badly in water.

PRINCEPS 1W-Y
Perianth creamy sulphur, deeper yellow on back, crêpey texture. Segments twisted, pointed, separated. Corona clear yellow, a little paler at base, almost straight with lobed edge.

PRINCESS BEATRIX 2W-R
Perianth poeticus white, deeply creased, notched, sometimes buckled, bottom 3 segments inclined to reflex. Segments flat, rounded, well overlapping. Corona bright apricot orange with narrow edge of creamy apricot, deeply fluted, double frilled and lobed edge. Stiff.

PRINCESS OF WALES 3W-WWO
Perianth cream, circular shape. Segments flat, rounded, overlapping. Corona greenish lemon with thin red line at edge, very flat, fluted. Stiff and sturdy.

PRINCESS VICTORIA 2Y
Perianth pale sulphur yellow, edges curl in, slightly creased, star shape. Segments flat, pointed, separated. Corona pale greenish lemon, shading to light orange edge for 1/3 its length, fluted, bowl shape. Neck grows in water.

PRINCIPAL 1Y-Y
Perianth soft sulphur yellow, crêpey texture, creased. Segments flat, pointed, broad, overlapping. Corona bright lemon yellow.

PRINS CARNAVAL 11Y-OOY
No description available.

PRIORITY 1Y-Y
Perianth deep rich golden yellow, stiff. Segments flat, pointed, broad, overlapping. Corona deeper golden yellow. Well proportioned.

PRISK 7W
Discarded - Small, flowers below foliage, jonquilla hybrid.

PRIVATEER 3W-O
Perianth yellow/white. Segments broad and overlapping, very slightly inflexed. Corona small, frilled orange cup. Neck grows badly in water.

PROFESSOR EINSTEIN 2W-R
 Perianth poeticus white, creased, back 3 segments inclined to reflex. Segments flat, rounded, well overlapping. Corona bright solid orange-red, deeply fluted lobed edge, flat. Rather rough and floppy. Neck grows in water.

PROFILE 2W-Y
 Perianth white, slightly twisted, edges uneven. Segments broad, would fit into a double triangle, some notched at tip. Corona pale primrose, quite short, flanged, loosely frilled at edge. Neck grows badly in water.

PROLIFIC 1Y-Y
 No description available.

PROLOGUE 1W-Y
 Perianth cream, slightly creased. Segments would fit into a double triangle, acute. Corona lemon, long and narrow then rolled back, gently lobed. Neck grows in water.

PROMPTITUDE 2W-Y
 Perianth creamy white, slightly creased. Segments flat, pointed, overlapping. Corona clear light lemon yellow, neat edge.

PROSPERO 3W-R
 Perianth creamy white, stained yellow base, slightly creased, notched, reflexed, circular outline. Segments flat, rounded, broad, well overlapping. Corona bright orange throughout, fluted, frilled, lobed, almost flat.

PROVERB 2Y-Y
 Perianth yellow. Outer segments broad, inner ovate and overlapping. Corona medium sized, open mouthed, and same yellow as perianth.

PROWESS 2W-Y
 Perianth white. Segments rounded and well overlapping. Corona large frilled, bowl shaped, yellow, but paler towards centre. Neck grows badly in water.

PTOLEMY 1W-Y
 Perianth lemony white. Segments broad and overlapping, twisted and slightly inflexed. Corona large frilled, flared, yellow in colour. Neck grows in water.

PURITY 2W-W
 Perianth white. Segments broad, overlapping and slightly inflexed. Corona medium sized, lobed and white in colour. Neck grows badly in water.

PYTCHLEY 3W-R
 Perianth creamy white, stained yellow at base, deeply creased, top 3 segments buckled. Segments flat, pointed, well overlapping. Corona deep orange red throughout, fluted. Neck grows stiffly in water.

QUEEN JULIANA 1W-W
 Perianth white. Segments broad, overlapping and twisted. Corona medium sized flared, frilled and lemony/white in colour.

QUEEN OF BICOLORS 1W-Y
 Perianth creamy white, stained yellow at base. Segments slightly twisted, pointed, broad, overlapping, edges curled in a little. Corona bright yellow, wide mouthed.

QUEEN OF HEARTS 3W-O
 Perianth cream, stained yellow at base. Segments flat, pointed, overlapping. Corona burnt orange throughout, fluted.

QUEEN OF NARCISSI 3W-YYR
 Perianth creamy white, stained yellow at base, creased. Segments flat, pointed, broad, well overlapping. Corona deep clear greenish lemon with bright orange red edge for 1/3 of its length, fluted. Neck grows in water.

QUEEN OF STARS 3W-YYR
 Perianth cream, stained yellow at base. Segments flat, rounded, overlapping. Corona deep gold with bright orange red edge for 1/3 of its length, fluted.

QUEEN OF THE NORTH 3W-Y
 Perianth creamy white. Segments flat, pointed, barely overlapping. Corona pale lemon yellow, fluted, pinched in.

QUEEN OF ULSTER 1W-Y
 Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona creamy lemon yellow.

QUEEN'S COUNSEL 1W-W
 Discarded - Flowers below foliage, susceptible to basal rot.

QUEEN'S LADY 1W
 Perianth creamy white, crêpey texture. Segments flat, pointed, broad, overlapping. Corona pale creamy lemon, paler at base.

- QUEENSLAND** 2W-P
Perianth creamy white, a little creased, top segments inclined to incurve. Segments slightly twisted, pointed, overlapping. Corona opens coppery salmon, developing cream flushed salmon base 1/2, shading to salmon edge, occasional white blotches on edge, frilled, lobed, trumpet proportion. Neck grows in water.
- QUETTA** 3W-O
Perianth creamy white, edges curl in, a little reflexed. Segments flat, pointed, overlapping. Corona greenish gold eye, changing to primrose and shading through apricot orange to deep orange red edge, fluted, bowl shape.
- QUIRINUS** 2Y-O
Perianth yellow. Segments broad, smooth, rounded, and 1/2 overlapping. Corona burnt orange with deep lobed edge. Poor neck. Neck grows badly in water. Weak stems.
- RADIATION** 2W-P
Discarded - Short, poor colour.
- RAFFREY** 2W-WWY
Perianth white. Segments very broadly ovate and overlapping. Corona large frilled cup, yellow with white centre.
- RAINBOW** 2W-WWP
Perianth creamy white. Segments rounded, overlapping at base, bottom Segments broad, well rounded, slightly creased, curled in at edges. Corona moss green throat, edged by band of cream fading to paler cream, developing to salmon top edge; deep bowl-shaped, lobed edge. Neck grows in water.
- RAMESES** 2W-R
Perianth poeticus white, stained yellow base, broad circular outline. Segments rounded, overlapping at base, slightly creased, curled in at edges. Corona bright burnt orange, bowl-shaped, fluted, lobed edge. Neck grows in water.
- RAMILLIES** 2Y-O
Perianth soft lemon yellow with whitish tips, creased, crêpey texture. Segments flat, pointed, overlapping. Corona bright orange red, fluted.
- RAMOAN** 3W-GYO
Perianth white. Segments very broadly ovate and overlapping, inner whorl more ovate than outer. Corona small, frilled cup, yellow with orange rim. Neck grows in water.
- RANGATIKEI** 2
Perianth yellow. Segments broad and overlapping, slightly inflexed. Corona medium sized frilled cup, darker yellow than perianth. Neck grows in water.
- RAPALLO** 3Y-YYR
Perianth pale lemon yellow, creased, often notched, circular outline. Segments flat, rounded, overlapping. Corona greenish lemon changing to bright orange red rim for 1/4 its length, fluted, almost flat. Stiff. Neck grows in water.
- RASHEE** 1W-W
Perianth pale creamy white, smooth, slightly hooded. Segments would fit into a double triangle, pointed. Corona fairly long, flanged, pale creamy white, edge slightly frilled. Weak, hollow stems. Neck grows in water.
- RATHCAVAN** 2W-GWW
Discarded - Small.
- RATHGAR** 2W-WWY
Perianth white. Segments broad and overlapping. Corona flared, frilled, open mouthed, creamy white, with yellow rim. Neck grows badly in water.
- RATHKENNY** 1W-Y
Perianth creamy white, much yellow on back and stained yellow at base. Segments flat, pointed, overlapping. Corona deep clear lemon yellow.
- RATHOWEN FLAME** 2Y-O
No description available.
- REALM** 2Y-Y
Perianth yellow. Outer segments very broadly ovate, inner more ovate and overlapping. Corona large frilled cup. Same yellow as perianth. Neck grows badly in water.
- RED ABBOT** 2W-O
Perianth creamy white, stained yellow at base, floppy. Segments flat, pointed, overlapping. Corona bright clear orange red.
- RED APRIL** 2W-R
Perianth opens primrose, develops cream stained yellow base, slightly creased, edges curl in. Segments flat, pointed, overlapping. Corona bright orange red throughout, fluted. Neck grows in water.

- RED BAY** 2Y-R
Perianth yellow. Segments very broadly ovate, inner more ovate and overlapping. Corona small, frilled cup, orange/red.
- RED BEACON** 3W-O
Perianth cream, deeper at base, a little creased. Segments flat, pointed, overlapping. Corona orange red shading to orange gold base, fluted.
- RED BIRD** 2W-O
Perianth creamy white stained deep yellow at base. Segments flat, pointed, well overlapping. Corona brilliant orange throughout, fluted.
- RED BUTTON** 2Y-O
Perianth sulphur with paler tips, creased, circular outline. Segments flat, rounded, well overlapping. Corona orange red throughout, fluted, lightly frilled, bowl shape. Stiff. Neck grows in water.
- RED COTTAGE** 2W-YYR
Perianth poeticus white, slightly twisted. Segments edges incurling, broad, rounded, overlapping, some notched near tip. Corona lemon at base becoming red at edge in two distinct bands, short, bowl shaped, creased, ragged at edge. Neck grows badly in water.
- RED CROSS** 2-YOO
Perianth primrose yellow, stained deep yellow at base, creased, edges curled back. Segments twisted, pointed, overlapping. Corona bright clear orange, straight sided.
- RED DEVON** 2Y-O
Perianth clear lemon yellow, creased, odd segments catch in corona, pointed outline. Segments flat, pointed, broad, overlapping. Corona bright orange-red shading to slight orange base, deeply lobed edge. Neck grows badly in water.
- RED GAUNTLET** 3W-O
Perianth pale primrose yellow, deeper at base, paler at tips. Segments flat, pointed, barely overlapping. Corona bright orange, deeply fluted, pinched in.
- RED GOBLET** 2Y-O
Perianth bright lemon yellow suffused orange, a little paler at base, with whitish tips. Segments flat, rounded, overlapping. Corona bright orange-red, a little paler at base, goblet shape.
- RED GUARD** 8YO-O
Perianth deep yellow, suffused red, paler at tips and centre streaks. Segments flat, pointed, overlapping. Corona deep orange red, fluted. Neck grows in water.
- RED HACKLE** 2W-YOO
Perianth creamy white, stained a little yellow at base, pointed outline, reflexed. Segments flat, pointed, overlapping. Corona deep burnt orange red, fluted. Neat.
- RED IDOL** 2Y-R
Perianth bright lemon overlaid orange for half its length, develops paler tips, slightly creased, inclined to reflex. Segments flat, pointed, broad, overlapping. Corona deep orange red throughout, fluted, lightly frilled, straight sided. Stiff. Neck grows in water.
- RED LAKE** 3W-OOR
Perianth white. Segments very broadly ovate and overlapping. Corona very small frilled cup, mainly orange with a red rim. Neck grows badly in water.
- RED RANGER** 2Y-O
Perianth lemon yellow developing pale tips, edges curl in. Segments flat, pointed, well overlapping. Corona bright orange red throughout, fluted, bowl shape. Neck grows in water.
- RED RASCAL** 2Y-R
Discarded - Small, flimsy.
- RED RIBAND** 2Y-YYR
Perianth sulphur, creased, pointed outline. Segments flat, pointed, overlapping. Corona greenish lemon base, changing to deep orange edge for 1/3 of its length, occasional white blotches on edge, fluted, lightly frilled, bowl shape. Neck grows in water.
- RED RIM** 9W-YYR
Perianth white, circular outline, reflexed. Segments flat, rounded, overlapping. Corona lemon yellow changing to deep red edge for 1/4 its length with slight greenish eye. Style very long.
- RED SATIN** 2W-R
Perianth white. Segments very broadly ovate and overlapping. Corona small frilled cup, yellow with orange rim. Neck grows in water.

RED SQUIRREL 2Y-O
 Perianth lemon, paler at tips, stained slight orange base, crêpey, a little creased, edges curl in slightly. Segments flat, rounded, overlapping. Corona bright orange throughout, slightly fluted, lightly frilled, almost straight sided. Neck grows stiffly in water.

RED SUNRISE 2Y-O
 Discarded - Long neck, floppy flower.

REDMAN 2Y-R
 Perianth lemon, outer segments rounded, inner segments pointed, half overlapping, some longitudinal ridges. Corona orange, funnel shaped, edge frilled.

REDMARLEY 2Y-O
 Perianth light sulphur yellow, edges notched, often caught in corona. Segments twisted, pointed, overlapping. Corona deep orange with some white blotches on edge, straight sided, deeply frilled.

REDSTART 3W-GWP
 Perianth white. Outer segments very broadly ovate, inner more ovate, and overlapping. Corona small, frilled cup, yellow with an orange rim. Neck grows badly in water.

REGALITY 3W-YOO
 Perianth poeticus white, stained yellow base, creased, sometimes notched, edges curl in, circular outline. Segments flat, rounded, broad, well overlapping. Corona gold 2/3 shading to deep orange edge, fluted, frilled, deeply lobed. Neck grows badly though stiffly in water.

REGINALD DIXON 2Y-O
 Perianth soft yellow flushed slight orange, creased, crêpey texture, inclined to be hooded. Segments flat, rounded well overlapping. Corona bright orange-red, shading to orange-gold, fluted.

REMBRANT 1Y-Y
 Perianth bright deep lemon yellow, crêpey texture. Segments twisted, pointed, overlapping. Corona deep canary yellow, open mouthed. Neck grows in water.

RENE DE CHALONS 2Y-YOO
 Perianth soft sulphur yellow, paler at tips, a little deeper at base, creased. Segments flat, pointed, broad, overlapping. Corona opens lemon yellow, develops to light orange shading to deep gold base. Neck grows in water.

RENEE DONALDSON 2Y-O
 Perianth very pale yellow. Segments broad and overlapping. Corona medium sized frilled orange cup.

REPRIEVE 3W-GWY
 Discarded - Small.

REPROOF 2Y-R
 Perianth yellow. Segments broad and overlapping. Corona small frilled cup, pale orange in colour. Neck grows badly in water.

RESERVE 2Y-O
 Perianth deep bright lemon, paler at tips, creased, a little crêpey, edges curl in. Segments flat, pointed, well overlapping. Corona deep bright orange, fluted, lightly frilled, almost straight sided. Neck grows stiffly in water.

RETREAT 2Y-R
 Discarded - Severe virus in stock.

REVELRY 2Y-O
 Perianth clear lemon yellow, smooth. Segments flat, rounded, well overlapping. Corona deep solid orange red, almost straight sided. Very stiff and sturdy. Neck grows in water.

REWA 2W-O
 Perianth white, stained slight yellow at base, circular outline. Segments flat, rounded, overlapping. Corona deep apricot shading to deep apricot orange edge. Too small for height.

REYNOLDSTOWN 2Y-O
 Perianth bright deep lemon yellow, suffused red at base, pointed outline. Segments flat, pointed, overlapping. Corona solid orange red, pinched in. Stiff.

RIBER 1W-W
 Discarded - Short, hangs head.

RICHARD TAUBER 8W-Y
 Perianth white stained slight yellow at base, creased, edges curled in. Segments flat, pointed, broad, overlapping. Corona deep orange gold with bright orange edge and greenish eye, fluted. Style reaches almost as long as longest stamens.

RICHHILL 2Y-YYR
 Discarded - Hangs head, flowers below foliage.

- RIDEAU HALL** 3W-YYO
Perianth pure white, stained yellow at base, creased. Segments flat, rounded, well overlapping. Corona deep lemon gold shading to bright orange red for 1/3 its length, extra frilled edge. Neck grows in water.
- RIESENBECK** 2W-YYO
Perianth white, stained yellow at base, creased, curled in at edges, perianth catches in corona. Segments flat, pointed, well overlapping. Corona deep lemon yellow changing to bright orange red edge for 1/5 of its length, extra frilled, fluted. Neck grows in water.
- RIJNVELD'S EARLY SENSATION** 1Y-Y
Perianth pale greenish sulphur yellow, paler tips, green on back. Segments flat, pointed, overlapping. Corona clear-medium lemon yellow, a little paler at base.
- RIMA** 1W-P
Perianth creamy white. Segments smooth, slightly twisted, would fit into double triangle. Corona pale pink fading to cream at the edge, slightly flanged with frilled edge. Neck grows in water.
- RING O' ROSES** 2Y-YYR
Perianth pale lemon yellow with paler tips, crêpey texture. Segments flat, pointed, overlapping. Corona gold shading to bright orange edge for 1/4 its length.
- RINGMASTER** 2Y-YYO
Perianth yellow. Segments very broadly ovate and overlapping. Corona medium sized frilled cup, deeper yellow than perianth with an orange rim. Neck grows in water.
- RINSEY** 3W-W
Perianth creamy white, edges curl in. Segments flat, pointed, broad overlapping. Corona cream with slight greyish cream eye, fluted, stiff. Neck grows in water.
- RIOTOUS** 4Y-Y
Perianth yellow. Double segments ovate and slightly overlapping. Corona double, frilled and same yellow as perianth. Neck grows in water. Weak stems.
- RIPPLING WATERS** 5W-W
Perianth creamy white. Segments flat, pointed, overlapping. Corona creamy white, corrugated horizontally, yellow base inside. Long style.
- RISTIN** 1Y-Y
Perianth yellow. Segments very broadly ovate and overlapping. Corona frilled funnel, medium sized, same yellow as perianth. Neck grows badly in water.
- RIVA** 2W-W
Perianth creamy white, edges curled in. Segments slightly twisted, pointed, barely overlapping. Corona opens creamy primrose, develops buff-cream, ageing to almost white. Almost trumpet proportion.
- ROB ROY** 3W-R
Perianth white. Outer segments very broadly ovate, inner ovate and overlapping. Corona small frilled deep orange cup. Neck grows badly in water.
- ROBERTA** 1W
Perianth creamy white, stained yellow base, crêpey. Segments flat, pointed, overlapping. Corona opens lemon, develops deep cream, deeply lobed, flanged.
- ROBESPIERRE** 2W-O
Perianth creamy primrose stained a little yellow at base, smooth, star shaped. Segments flat, pointed, barely overlapping. Corona bright orange red.
- ROBEY** 1Y-Y
Perianth clear lemon yellow, slightly crêpey texture. Segments twisted, pointed, overlapping. Corona deeper bright lemon yellow, wide-mouthed.
- ROCKET** 2W
Perianth poeticus white, stained yellow base, creased, inclined to reflex, circular outline. Segments flat, rounded, overlapping. Corona gold 2/3 changing to bright orange red edge, deeply fluted, frilled, lobed. Neck grows badly in water.
- ROCOCO** 2W
Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona deep apricot shading to paler apricot with deep greeny lemon centre, deeply cut double frilled edge, almost flat, covering over half the perianth.
- RODOMONT** 2Y-O
Perianth clear lemon yellow, crêpey texture. Segments flat, rounded, well overlapping. Corona orange, shading to gold base. Neat. Neck grows in water.
- ROGER** 6Y-O
Discarded - Small, short, cyclamineus hybrid.

ROIMOND Discarded - Hangs head, poor poise.	2W-O
ROMAINE Discarded - Rough flower.	1W-W
ROMAN CANDLE Perianth white. Segments broad overlapping, curled in at edges, and slightly inflexed. Corona frilled open mouthed, creamy yellow in colour. Neck grows in water.	2W-P
ROMAN TILE Perianth white. Segments broad, overlapping and slightly inflexed. Corona medium sized frilled cup pink with red rim. Neck grows badly in water.	2W-WYP
ROMANTICA Discarded - Small, rough flower.	2O-O
ROMANUS (syn. ? DOUBLE ROMAN) Perianth yellow. Double segments, slightly star shaped. Corona, double, yellow in colour.	4W-O
ROMEIO Perianth sulphur, slightly hooded. Segments broad, rounded, overlapping. 4-7 flowers per stem. Corona pale orange, small, flat, fluted giving frilled edge. Neck grows badly in water.	8Y-O
ROSA VAN LIMA Perianth creamy white, creased, star shape. Segments flat, pointed, overlapping. Corona opens primrose, develops coppersy peach, paler at base, ageing to creamy peach, fluted. Neck grows in water.	2W-P
ROSARIO Perianth creamy white, creased, edges curl in a little. Segments flat, pointed, broad, well overlapping. Corona cream base, shading to soft salmon pink for 1/2 its length, of trumpet proportions. Neck grows in water.	2W-P
ROSDEW Perianth creamy white. Segments very broadly ovate and overlapping, slightly inflexed. Corona medium sized frilled funnel, yellow in colour. Neck grows in water.	2W-P
ROSE BROCADE No description available.	2W-P
ROSE CAPRICE Perianth white. Segments broad and overlapping. Corona medium sized frilled cup, pale pink, with a deeper pink rim.	2W-GPP
ROSE NOBLE Perianth white. Segments ovate and 1/2 overlapping. Corona medium sized frilled cup pink in colour. Neck grows badly in water.	2W-P
ROSE OF CUAN Discarded - Small.	2W
ROSE OF MAY Perianth creamy white, flat, rounded, broad, well overlapping, circular outline. Made up of 4 or 5 regular whorls of perianth segments, gradually becoming more pointed in centre. Neat and solid flower. Neck grows in water.	4W-W
ROSE OF TRALEE Perianth creamy white, slight pink staining at base, perianth often catches in corona, edges notched. Segments flat, pointed, overlapping. Corona opens apricot salmon throughout, developing to paler colour, of trumpet proportion.	2W-P
ROSE RIBBON Perianth creamy white. Segments broad, overlapping and slightly inflexed. Corona medium sized frilled cup - yellow with orange rim. Neck grows badly in water.	2W-YYP
ROSE ROYALE Perianth white, stained yellow/green/pink back base, pointed outline. Segments pointed, creased, crepey, slightly curled at edges. Corona moss green throat, salmon, shallow scalloped edge.	2W-P
ROSEANNA Perianth creamy white, creased, sometimes notched, pointed outline. Segments flat, pointed, overlapping. Corona pure pink, paler at base, ageing to creamy pink throughout, almost straight sided.	2W-P
ROSEATE HUES Perianth cream, creased, top 3 segments often buckled and incurved. Segments flat, pointed, overlapping. Corona creamy primrose with narrow coppersy salmon rim, frilled, deeply lobed, broad mouth. Variable colour development when opened in water.	2W-YYP

ROSEBUD 3W-O
Perianth opens primrose yellow, develops creamy white, stained yellow at base, edges curled in. Segments flat, rounded, overlapping. Corona orange with deeper orange rim, fluted, uneven edge.

ROSEDEW 2W-P
Perianth white. Segments broad and overlapping. Corona frilled open mouthed, and pink in colour.

ROSEWELL 2W-P
Perianth creamy white, slightly creased, edges curl in, pointed outline. Segments flat, pointed, well overlapping. Corona bright peach pink, a little paler at base, neat. Neck grows in water.

ROSEWORTHY 2W-P
Discarded - Short, flowers below foliage.

ROSEYARDS 2W-P
Discarded - Small, short, poor colour.

ROSIE MCNABB 2W-P
Perianth white. Segments broad and overlapping. Corona medium sized, frilled, pink cup. Neck grows badly in water.

ROSSINI 2W-O
Perianth creamy white, stained yellow base. Segments rounded, overlapping at base, bottom segments broad, top segments narrower, buckled, all curled in at edges. Corona rich burnt orange, expanded, deeply fluted irregular edges. Head faces downwards. Neck grows in water.

ROSTOV 2W-W
No description available.

ROSY DIAMOND 2W-WWP
Perianth white, creased, edges curl in and a little waved. Segments slightly twisted, pointed, overlapping. Corona bright pink, a little paler at base, fluted. Neck grows in water.

ROSY PINK 2W
Perianth white. Segments broad and overlapping with a ridged texture. Corona medium sized frilled, flared funnel, white with pink rim. Neck grows badly in water.

ROSY SUNRISE 2W-P
Perianth creamy white, creased. Segments twisted, pointed, broad, overlapping. Corona salmon buff shading to salmon pink edge, extra frilled.

ROTONDE 3W-Y
Discarded - Short, severe virus in stock, susceptible to basal rot.

ROUGE 2Y-O
Perianth pinkish-primrose yellow, edges curled in a little, creased, star-shaped, a bit floppy. Segments flat, pointed, broad, overlapping. Corona bright orange red. Neck grows in water.

ROVER 1Y-Y
Perianth yellow. Segments ovate and overlapping, slightly twisted. Corona frilled flared funnel, same colour as perianth. Neck grows badly in water.

ROWALLANE 1Y-Y
Perianth yellow. Segments broad and overlapping. Corona frilled, flared cylinder, same colour as perianth.

ROWENA LEE TEAGLE 2W-ORR
Perianth creamy white, a little creased, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona bright orange red, a shade paler at edge, deeply fluted. Neck grows in water.

ROXANE 1W-W
Perianth creamy white. Segments flat, pointed, broad, overlapping. Corona opens pale lemon yellow, develops to cream. Flower coarse, heavy and out of proportion.

ROYAL ARMOUR 1Y-Y
Perianth bright lemon, darker on mid underside, double triangle-shape, segments round and overlapping. Corona bright golden yellow with coarse rolled crenate edge. Long neck. Neck grows badly in water.

ROYAL CHARGER 2Y-O
Discarded - Short.

ROYAL CHARM 2Y-ORR
Perianth bright rich yellow, barely creased, slightly crepey. Segments rounded, overlapping at base, curled in at edges. Corona basal third orange, developing to deep orange, then with reddish tinge in outer third; straight, with coarsely scalloped edge. Neck grows in water.

ROYAL CROWN 2W-O
Discarded - Short, floppy, flowers below foliage.

ROYAL DELIGHT 1Y-Y
Perianth yellow. Segments ovate, slightly overlapping and twisted. Corona large, frilled, open mouthed, same colour as perianth.

ROYAL GOLD 1Y-Y
No description available.

ROYAL MAIL 2Y-O
Perianth pale lemon yellow, creased, edges curled in. Segments flat, pointed, broad, overlapping. Corona burnt orange, fluted, bowl shaped.

ROYAL OAK 1Y-Y
Discarded - Small.

ROYAL ORANGE 2W-O
Discarded - Short, flowers below foliage.

ROYAL PORCELAIN 2W-WWP
Perianth white. Outer segments very broadly ovate, inner ovate and overlapping. Corona medium sized frilled yellow. Neck grows in water.

ROYAL RANSOM 2Y-O
Perianth pale primrose yellow, suffused apricot, star shaped, edges curled in. Segments flat, pointed, broad, overlapping. Corona bright orange-red, often with white blotches on the edge, fluted.

ROYAL REGIMENT 2W-O
Perianth creamy white, stained yellow base, rounded outline. Segments rounded, overlapping at base, broad, generally smooth, wrinkled at lower edges, curled in at edges. Corona burnt orange, deeply fluted, irregularly lobed. Neck grows in water.

ROYAL ROMANCE 2
Perianth yellow. Segments broad and overlapping. Corona small, frilled cup orange with paler eye. Neck grows badly in water.

ROYAL SEAL 2W-R
Perianth poeticus white, stained yellow base, top segments curl in, bottom segments reflex. Segments flat, pointed, overlapping. Corona small yellow green eye changing to sealing wax red, deeply fluted, frilled, unevenly lobed edge, nearly flat. Neck grows badly in water.

ROYAL WINDSOR 2Y-YRR
Perianth sulphur yellow, crêpey texture, creased, edges curl back. Segments flat, rounded, overlapping. Corona bright orange, fluted, deeply cut edge. Untidy.

ROYALIST 1Y-Y
Perianth soft sulphur yellow, smooth. Segments flat, pointed, broad, overlapping. Corona light lemon yellow, neat. Well proportioned.

RUBICON 2Y-Y
Discarded - Short.

RUBRA 2W-Y
Perianth creamy white, slightly creased, edges curled in a little. Segments flat, pointed, overlapping. Corona bright apricot orange, a little paler at base, fluted.

RUBY KING 2
Perianth sulphur yellow, developing paler, deeply creased, often buckled, pointed outline. Segments flat, pointed, overlapping. Corona bright orange, paler at base, with white blotches on edge, bowl shape, deeply fluted.

RUBY TAIL 2W-R
Perianth white. Outer segments very broadly ovate, inner ovate, and overlapping. Corona medium sized frilled cup orange in colour. Neck grows in water.

RUMBA 2Y-O
Perianth opens sulphur, develops cream stained yellow base, creased, notched, sometimes caught in corona, circular outline. Segments flat, rounded, broad, overlapping. Corona slight gold base shading to bright orange red, some cream blotches on edge, fluted, frilled, lobed, shallow bowl shape. Stiff. Neck grows in water.

RUSSET 3Y-R
Perianth opens primrose, develops cream stained yellow base, creased, sometimes notched, bottom segments reflex, circular outline. Segments flat, rounded, well overlapping. Corona deep orange red, deeply fluted, frilled, lobed, almost flat. Neck grows in water.

RUSTOM PASHA 2Y-O
Perianth clear lemon yellow, a little creased, crêpey texture, notched and often caught in the corona. Segments flat, pointed, overlapping. Corona opens greenish gold developing to deep orange red with slightly paler edge.

SACAJAWEA 2Y-YYO
 Perianth lemon, slightly creased. Segments broad, overlapping, slightly twisted. Corona deep lemon at base becoming orange at edge, open mouthed, frilled. Neck grows badly.

SAILING LIGHT 2W-YYO
 Perianth creamy white, slightly creased, edges curl in a little. Segments flat, pointed, overlapping. Corona cream, shading to bright apricot orange for 1/2 its length, greyish green eye, fluted, almost flat. Hangs head badly when young. Corona edge dies first in water.

SAINT AGNES 8W-O
 Perianth creamy white, stained slight yellow at base, creased, inclined to reflex. Segments flat, rounded, overlapping. Corona bright red shading to greenish gold base, fluted. Style very long.

SAINT BRENDAN 2W-W
 Perianth creamy white, slightly creased and crêpey. Segments flat, pointed, broad, well overlapping, shovel shape. Corona creamy white, deeply frilled, flanged, broad mouth.

SAINT BRIDE 1W-W
 Discarded - Short, hangs head, very susceptible to basal rot.

SAINT EGWIN 2Y-Y
 Perianth clear pale lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona deeper lemon yellow. Neat.

SAINT ISSEY 2Y-Y
 Perianth deep clear lemon, crêpey texture. Segments slightly twisted, pointed, broad, well overlapping. Corona deeper clear golden yellow, edge dimpled and frilled. Well proportioned.

SAINT IVES 2Y-Y
 Perianth bright clear lemon yellow, one segment often buckled. Segments twisted, pointed, overlapping. Corona deeper lemon yellow, fluted, bowl shape.

SAINT JUST 3W-YYO
 Perianth creamy white, stained yellow at base, some segments creased edges curled in a little. Segments flat, pointed, overlapping. Corona bright gold shading to orange red edge for 1/2 of its length.

SAINT KEVERNE 2Y-Y
 Perianth deep, clear lemon yellow, smooth. Segments flat, pointed, broad, overlapping. Corona deeper gold, almost straight, well proportioned. Neck grows in water.

SAINT KEYNE 8W-O
 Perianth pure white stained slight yellow at base, creased, reflexed. Segments flat, rounded, overlapping. Corona bright orange red, shading to orange base, with greenish eye, almost flat. Style reaches half-way between short and long stamens.

SAINT LOUIS 3Y-YOO
 Perianth cream, deeper at base, creased, edges curled in, star-shaped. Segments flat, pointed, broad, overlapping. Corona gold with bright orange red edge for 1/3 of its length, deeply fluted.

SAINT MARTIN 2
 Perianth opens pale primrose yellow, darker centre line on 3 outer segments, develops to dirty cream, edges curled in, crêpey texture, star shaped. Segments flat, pointed, barely overlapping. Corona pale orange shading to gold base, fluted. Neck grows in water.

SAINT MARY 1W-W
 Discarded - Hangs head.

SAINT MORITZ 2W-W
 Discarded - Badly balanced.

SAINTFIELD 2W-P
 No description available.

SALMON TROUT 2W-P
 Perianth white. Segments broad, overlapping and slightly inflexed. Corona medium sized frilled cup, pink with yellow center. Neck grows in water.

SALOME 2W-PPY
 Perianth creamy white, smooth. Segments broad, overlapping for 3/4 of length, edges curled in at tips. Corona pink fading to orange at edge, slightly flanged, frilled.

SALTASH 2Y-O
 Perianth clear lemon yellow, crêpey texture, star shaped. Segments flat, pointed, overlapping. Corona bright burnt orange sometimes with white blotches on edge. Neck grows in water.

SAMITE 1W-W
 Perianth creamy white, smooth. Segments flat, pointed, broad, overlapping. Corona opens creamy primrose, develops to cream. Neat.

- SANTA CLAUS** 4W-W
Perianth creamy white, three back segments bronze green, rounded outline. Segments broad, rounded, overlapping, multiple rows with symmetrical arrangement. Corona segments rudimentary.
- SANTA LUCIA** 2W-W
Discarded - Short, flowers below foliage.
- SANTIAM** 2Y-YYO
Perianth pale yellow. Segments very broadly ovate in outer whorl, ovate in inner, and overlapping. Corona large frilled cup, yellow with orange rim. Neck grows in water.
- SARABANDE** 9W-GOR
Perianth poeticus white, creased down centre segments and edges folded back. Segments flat, rounded, barely overlapping. Corona greenish gold eye shading to greenish lemon and orange red with deeper red rim, fluted, almost flat. Style ½ way between long and short stamens. Neck grows badly in water.
- SARAH LANCASHIRE** 3W
Perianth creamy white, crêpey, top segments curl in a little, bottom segments reflex. Segments flat, pointed, overlapping. Corona small green eye, changing to pale greenish lemon with wire rim of gold, fluted, lightly frilled, bowl shape. Neck grows in water.
- SARCHEDON** 9W-GYR
Perianth pure white, slightly yellow at base, a little creased. Segments flat, rounded, overlapping. Corona bright lemon yellow changing to deep red for 1/4 of its length with a deeper yellow eye. Style extremely long. Neck grows in water.
- SARGASSO** 2Y-YOO
Perianth light lemon yellow, slightly creased. Segments flat, rounded, overlapping. Corona gold shading to bright orange edge for 1/2 its length, edge deeply lobed. Neck grows in water. Stiff and sturdy.
- SASSENHEIM'S GIANT** 1Y-Y
Perianth lemon yellow, crêpey texture, floppy. Segments twisted, pointed, separated. Corona deep lemon yellow, broad mouth, badly proportioned.
- SATIN QUEEN** 2W-YYO
Discarded - Short, flowers below foliage.
- SCANDAL** 3W-O
Perianth creamy white, stained yellow base, creased, notched, often buckled, reflexed. Segments flat, rounded, overlapping. Corona deep burnt orange red, barely fluted, slightly frilled, bowl shape. Neck grows in water.
- SCAPA** 1W-W
Perianth creamy white, creased, crêpey. Segments flat, pointed, overlapping. Corona cream crenate. Stiff.
- SCARLET ELEGANCE** 2Y-R
Perianth bright lemon yellow, odd segments buckled. Segments flat, pointed, overlapping. Corona brilliant deep red shading to gold base. Neck grows in water and corona edge dies first. Stiff.
- SCARLET GEM** 8Y-O
Perianth pale gold suffused red. Segments flat, rounded, overlapping. Corona bright orange red, almost flat, fluted.
- SCARLET LEADER** 2Y-O
Perianth dirty cream, stained yellow at base, creased. Segments flat, pointed, overlapping. Corona bright orange scarlet, often with white blotches on edge, deeply frilled. Neck grows in water.
- SCARLET PIMPERNEL** 2W-O
Perianth pure poeticus white, stained yellow at base. Segments flat, rounded, overlapping. Corona sealing-wax red, deeply fluted, and lobed edge.
- SCARLET ROYAL** 2Y-R
Perianth yellow. Segments very broadly ovate and overlapping. Corona medium sized frilled cup, deep orange in colour. Neck grows in water.
- SCARLETT O'HARA** 2Y-R
Perianth lemon yellow, crêpey texture, creased, often caught in corona. Segments flat, pointed, broad, overlapping. Corona bright orange, a little paler at base, fluted, bowl shape. Poor poise. Neck grows in water.
- SCILLY WHITE** 8W-Y
Perianth creamy white, edges curl in, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona deep cream, almost straight sided, slightly fluted. Style 1/2 length of perianth tube.
- SCINTILLA** 2W-R
Perianth opens primrose, developing pale tips and centre, deeply creased, buckled. Segments flat, pointed, overlapping. Corona deep burnt orange red, deeply fluted, uneven lobed edge.

- SCORCHER** 2Y-O
Perianth sulphur, creased, star-shaped, only overlapping at base. Segments pointed, narrow. Corona pale orange, short, open mouthed, ribbed, gently lobed rim. Neck grows badly.
- SCORPION** 2Y-R
Perianth opens sulphur yellow, developing to cream stained yellow at base, creased, edges curl in. Segments flat, pointed, overlapping. Corona gold base, shading to orange red for half its length, fluted, bowl shape.
- SCOTCH GOLD** 1Y-Y
Perianth yellow. Segments ovate and overlapping. Corona frilled flared cylinder, same colour as perianth. Neck grows in water.
- SCRABO** 2Y-O
Perianth sulphur yellow, paler at tips, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona bright orange red, fluted, almost flat. Neck grows in water.
- SEA GREEN** 9W-GYR
Discarded - Small.
- SEA URCHIN** 2W-Y
Perianth creamy white, creased, pointed outline. Segments flat, pointed, overlapping. Corona opens pale lemon, developing buff primrose with thick heavily frilled edge, with white blotches. Heavy head, ugly.
- SEAFORD** 2W-W
Perianth white. Segments broad and overlapping. Corona large frilled cup, creamy white in colour.
- SEAGULL** 3W-YYO
Perianth creamy white. Segments flat, pointed, barely overlapping. Corona bright greenish lemon with orange edge for 1/4 of its length.
- SEALING WAX** 2Y-O
Perianth egg yolk yellow, smooth. Segments broadly ovate, overlapping extensively. Corona deep orangey red, short, straight sided, fluted with folded edge. Neck grows stiffly in water.
- SEBASTOPOL** 2W-Y
Perianth creamy white, stained slight yellow at base, a little creased, crêpey, edges curl in. Segments flat, pointed, overlapping. Corona bright clear lemon, frilled, slightly flanged. Stiff.
- SELM LAGERLOF** 2W-YOO
Perianth creamy white, creased, floppy. Segments flat, pointed, broad, well overlapping. Corona brilliant orange shading to apricot orange base, almost flat, deeply fluted. Rough.
- SELTAN** 2W-P
Discarded - Short, flowers below foliage.
- SEMIRAMIS** 2W-Y
Perianth white. Segments very broadly ovate, overlapping, slightly inflexed and with a ridged texture. Corona very large frilled bowl, laid back onto perianth, yellow in colour. Neck grows badly in water.
- SEMPRE AVANTI** 2W-O
Perianth opens primrose, develops dirty creamy yellow, stained yellow at base, crêpey texture. Segments flat, pointed, broad, overlapping. Corona bright solid orange, fluted.
- SERAGLIO** 3Y-YYR
Perianth primrose yellow developing to cream, stained yellow at base, creased. Segments flat, rounded, overlapping. Corona greenish lemon with band of bright orange red for 1/4 of its length, almost flat.
- SHAGREEN** 3W-GWW
Perianth creamy white, slightly creased, circular outline. Segments flat, rounded, overlapping. Corona bright green base 1/3 changing to creamy white with narrow rim of buff, fluted, lightly frilled, almost flat.
- SHAH** 7Y-Y
Perianth rich deep yellow, smooth. Segments flat, pointed, broad, overlapping. Corona deeper rich yellow, smooth, of trumpet proportions.
- SHALDON** 2W-Y
Perianth white. Segments broad and overlapping, with a ridged texture. Corona large frilled cup, yellow in colour.
- SHANACH** 9W-YYR
Discarded - Small.
- SHANGHAI** 2Y-Y
Perianth sulphur, developing pale tips, crêpey, a little creased, slightly hooded. Segments flat, pointed, broad, well overlapping. Corona clear lemon, deeply frilled, broad bowl shape. Neck grows badly.

- SHANNON** 2W-W
Perianth creamy white. Segments flat, pointed, broad, overlapping, shovel shape. Corona opens primrose, develops cream with paler edge, of trumpet proportion. Hangs head.
- SHANTALLOW** 3W-GGY
Perianth white. Segments round, overlapping and slightly reflexed. Corona small frilled cup, white, with green eye and yellow rim. Neck grows badly in water.
- SHEEROE** 2Y-R
Discarded - Small, flowers below foliage.
- SHEIK** 2W-Y0Y
Perianth creamy white, stained yellow base and lower edges, pointed outline. Segments pointed, overlapping at base, fairly smooth, curled in at edges. Corona deep orange, bowl-shaped, fluted, irregularly lobed.
- SHEPHARD** 3W-OR
Perianth white. Segments round and overlapping. Corona small, frilled cup, orange with an orange/red rim. Neck grows badly in water.
- SHEPPY** 1Y-Y
Discarded - Small, flowers below foliage.
- SHINING WATERS** 2W-W
Perianth creamy white, creased, star shape. Segments flat, pointed, overlapping. Corona creamy white with rolled back edge, badly proportioned.
- SHIRLEY WYNESS** 2W-YYP
Perianth creamy white, creased, top 3 segments incurved, notched, often caught in corona. Corona opens pale lemon, develops cream with pale creamy peach edge, some white blotches on deeply frilled edge.
- SHOT SILK** 5W-W
Perianth creamy white. Segments flat, pointed, barely overlapping. Corona creamy white, corrugated horizontally. Long style.
- SHOWMAN** 2W-Y
Perianth creamy white, creased, edges curl in. Segments flat, pointed, overlapping. Corona buff primrose base for 2/3 its length, shading to creamy apricot with cream edge, fluted, deeply cut lobed edge.
- SIAM** 2W-P
Perianth creamy white, creased, notched, often caught in corona. Segments flat, pointed, overlapping. Corona pale creamy salmon shading to deeper edge, ageing paler, very broad mouth, often split, ugly. Neck grows in water.
- SIBERIA** 2W-W
Discarded - Short.
- SIDELIGHT** 9W-GOR
Perianth poeticus white, edges curl in, circular outline. Segments flat, rounded, overlapping. Corona moss green eye, shading through greenish lemon for 1/2 its length and a narrow band of deep red, fluted, almost flat. Style as long as longest stamens. Neck grows stiffly in water.
- SIDNEY TORCH** 2Y-YOO
Perianth sulphur yellow, deeply creased, notched, circular outline. Segments flat, rounded, well overlapping. Corona greenish lemon shading to bright orange red edge for half its length, deeply fluted, flat.
- SIGNAL LIGHT** 2W-O
Perianth creamy white, stained yellow at base, a little creased, slightly reflexed. Segments flat, pointed, overlapping. Corona deep burnt orange red, fluted, lightly frilled, shallow bowl shape. Neck grows stiffly in water.
- SIGRID UNSET** 3W-YOO
Perianth white stained yellow at base, curled in at edges. Segments flat, rounded, well overlapping. Corona gold shading to bright deep orange for 1/2 of its length, fluted. Stiff and sturdy.
- SILENT BEAUTY** 3W-YYR
Perianth white. Segments round and overlapping. Corona small frilled cup yellow with an orange/red rim. Neck grows badly in water.
- SILENT GLOW** 3W-R
Perianth white. Segments very broadly ovate and overlapping. Corona small frilled cup, red in colour. Neck grows badly in water.
- SILENT MORN** 3W-YYO
No description available.
- SILVER BUGLE** 2W-W
Perianth creamy white, a little creased. Segments flat, pointed, broad, overlapping. Corona creamy primrose.

- SILVER CHIMES** 8W-W
Perianth creamy white, smooth. Segments flat, pointed, broad, overlapping. Corona deep primrose with paler edge, almost straight sided, straight edge, smooth.
- SILVER LINING** 2W-W
Perianth creamy white, a little creased, edges curled in. Segments flat, pointed, broad, overlapping. Corona opens buff primrose, develops to pale cream with paler edge. Neck grows in water and corona edge dies first.
- SILVER MOON** 2W-O
Perianth white, smooth, 3 segments reflexed. Segments incurling edges, broad, rounded, overlapping. Corona opens sulphur, orange, sulphur, outer rim fades to cream, open mouthed to flat, fluted, deeply lobed rim. Neck grows badly in water.
- SILVER PENNY** 3W-O
Perianth creamy white, creased, edges curled back. Segments flat, rounded, barely overlapping. Corona apricot shading to light orange edge with greenish gold eye, fluted, almost flat. Long style.
- SILVER PRINCESS** 3W-W
Perianth white. Segments ovate and overlapping. Corona small frilled cup, white with a green eye. Neck grows badly in water.
- SILVER SALVER** 3W-GWW
Perianth white, edges curled in a little. Corona pure white (whiter than perianth) with thin buff line at edge and bright green eye, deeply fluted.
- SILVER SAND** (NL)
Perianth white, smooth, slightly hooded. Segments broad, ovate, edges incurling. Corona green at base shading to cream then deep cream at rim, short, open mouthed, fluted, slightly frilled edge. Neck grows badly in water.
- SILVER STANDARD** 2W-W
Perianth cream with primrose edges, slightly creased. Segments broad and overlapping, slightly hooded. Corona pale primrose, flat to open mouthed, deeply frilled. Neck grows badly in water.
- SILVER WEDDING** 1W-W
Discarded - Hangs head badly.
- SILVERTON** 1W-W
Perianth white. Segments broad and overlapping. Corona large frilled open mouthed, and yellow in colour.
- SILVESTER** 11W-Y
Perianth white. Segments very broadly ovate and overlapping. Corona split frilled intermixed orange/white in colour. Neck grows in water.
- SILVRETTA** 1Y-Y
Discarded - Short, flowers below foliage, rough flower.
- SINCERITY** 1W-Y
Perianth creamy white, slightly creased. Segments flat, pointed, broad, well overlapping. Corona pale creamy lemon, neat, hangs head badly.
- SIR PERCY** 2Y-O
Perianth yellow. Segments very broadly ovate and overlapping. Corona medium sized frilled cup, orange. Neck grows in water.
- SIR SAMUEL** 2W-P
Perianth poeticus white, smooth. Segments would fit into a double triangle, pointed. Corona pale pink, edge slightly paler, fades to cream, straight sided, long and narrow, ribbed, barely wavy edge. Neck grows in water. Weak stems.
- SIR WATKIN** 2Y-Y
Perianth primrose yellow, a little paler at tips, ageing to paler colour, edges curled back, making thin appearance. Segments twisted, pointed, barely overlapping. Corona deep lemon yellow.
- SIR WINSTON CHURCHILL** 4W-O
Perianth white. Double segments, rounded and overlapping. Corona orange, and interspersed with inner perianth whorl. Multiheaded. Neck grows badly in water.
- SISTER REINALDE** 2W-YYO
Perianth creamy white, creased, floppy. Segments flat, rounded, well overlapping. Corona greenish lemon with thin pale orange line at edge, deeper at base.
- SLEMISH** 2W-W
Discarded - Short, low flower production, very susceptible to basal rot.
- SLEVEEN** 2W-W
Discarded - Hangs head.

SLIEVEBOY 1Y-Y
Perianth yellow. Segments very broadly ovate and overlapping. Corona frilled flared cylinder, same yellow as perianth. Neck grows in water.

SLIGO 2Y-Y
Perianth lemon yellow, slightly creased, crêpey texture. Segments flat, pointed, broad, overlapping. Corona lemon yellow, of trumpet proportions.

SMARAGD 2W
Perianth creamy white, creased, edges curl back, floppy. Segments flat, pointed, barely overlapping. Corona pale greenish lemon, tinted with pale apricot at edge, deeply fluted. Rough. Neck grows in water.

SMILING MAESTRO 2Y-R
Perianth yellow. Segments broad, overlapping and slightly inflexed. Corona medium sized frilled orange cup laid back onto perianth segments. Neck grows in water.

SMILING MARY 3W
Perianth white. Segments very broadly ovate and overlapping. Corona small frilled cup, orange/red with yellow eye. Neck grows in water.

SMILING QUEEN 2W-YOO
Perianth pure white, crêpey substance, floppy. Segments flat, rounded, overlapping. Corona bright orange with yellow base, very densely frilled at edge.

SMYRNA 9W-GOO
Perianth white, slightly creased, circular outline. Segments flat, rounded, well overlapping. Corona bright orange red, shading a little paler to small moss green eye, flat, fluted. Style extremely long.

SNEEUWPRINSES 3W-YYO
Perianth creamy white, stained yellow at base, edges curl in, creased. Segments flat, pointed, overlapping. Corona gold shading to bright orange red for 1/3 its length, fluted, shallow bowl shape. Neck grows in water.

SNOW BUNTING 7W-Y
Discarded - Small, flowers below foliage, susceptible to basal rot, jonquilla hybrid.

SNOW DREAM 2W-W
Discarded - Short, very susceptible to basal rot.

SNOW GEM 3W-O
Perianth poeticus white, slightly twisted. Segments rounded, nearly a circular outline. Corona yellow at base becoming red at halfway up, crenate, slightly lobed at rim. Neck grows badly in water.

SNOW KING 9W-YYR
Perianth poeticus white, stained yellow at base, notched, creased, circular outline. Segments flat, rounded overlapping. Corona shading from gold to orange with a narrow rim of deep red, fluted, flat. Style longer than stamens. Neck grows badly in water.

SNOWBALL 4W-W
Discarded - Very low flower and bulb production.

SNOWBIRD 5W-W
Perianth creamy white. Segments flat, pointed, barely overlapping. Corona creamy white. Long style.

SNOWLINE 2W-W
Discarded - Very susceptible to basal rot.

SNOWSHILL 2W-W
Perianth white. Segments ovate and overlapping. Corona medium sized frilled cup, same white as perianth. Neck grows badly in water.

SOIRÉE 2W-Y
Discarded - Short, rough.

SOLARIO 2W-Y
Perianth cream, stained deep yellow at base, crêpey texture. Segments flat, pointed, overlapping. Corona bright lemon yellow, of almost trumpet proportions.

SOLE MIO 1Y-Y
Perianth deep clear lemon yellow, green stripe on back, crêpey texture, edges curled back. Segments twisted, pointed, overlapping. Corona deep golden yellow.

SOLE PONENTE 1Y-Y
Perianth yellow. Segments ovate and overlapping. Corona large frilled flared cylinder, slightly deeper yellow than perianth. Neck grows badly in water.

SOLINUS 2W-P
Perianth white. Segments broad and overlapping, slightly inflexed. Corona medium sized frilled flared cylinder, pink in colour. Neck grows in water.

SOLOGNE 2Y-O
Discarded - Rough flower.

SOLVEIG 1W-Y
Perianth creamy white, stained light yellow at base, creased. Segments flat, pointed, overlapping. Corona clear lemon yellow, open mouthed.

SOMALI 2Y-YOO
Discarded - Low flower yield, susceptible to basal rot.

SONATA 9W-GYR
Perianth white, edges curl in a little. Segments flat, rounded, overlapping. Corona greenish lemon with thin pencil line of deep red at edge, flat, fluted. Style extremely long. Neck grows in water.

SONIA SLOAN 2W-OOY
No description available.

SORBET 11W-YYO
Perianth white. Segments rounded, overlapping, creased, buckled and curled. Corona orange, striped/streaked segmented with frilled edge. Neck grows badly in water. Weak stems.

SOUTH POLE 1W-W
Perianth white. Segments very broadly ovate, overlapping and spreading. Corona large frilled rolled trumpet, same white as perianth. Neck grows in water.

SPACE AGE 2Y-Y
Perianth yellow. Segments ovate, overlapping, slightly ridged and slightly inflexed. Corona medium sized frilled cup, deeper yellow than perianth. Neck grows in water.

SPANISH GOLD 1Y-Y
Discarded - Small, flowers below foliage.

SPARKLING EYE 8W-GOO
Perianth creamy white, creased, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona green eye, changing to deep orange, fluted, shallow bowl shape.

SPECTATOR 2W-YYO
Discarded - Hangs head badly, susceptible to basal rot.

SPELLBINDER 1Y-WWG
Perianth pale primrose, slightly creased and twisted. Segments pointed, overlapping. Corona pale primrose with lemon edge, flanged, deeply cut edge.

SPELTER 2Y-O
Perianth deep lemon yellow, overlaid orange, paler at tips, edges curl in, slightly creased. Segments flat, rounded, broad, well overlapping. Corona deep bright orange throughout, slightly fluted. Poor poise. Neck grows stiffly in water.

SPENCER TRACY 2W-O
Perianth white, creased, often caught in corona. Segments flat, rounded, overlapping. Corona bright orange, deeply fluted and frilled into centre of corona. Neck grows in water.

SPICK AND SPAN 2Y-YYO
Perianth lemon yellow. Segments flat, pointed, broad, well overlapping. Corona bright orange red, paler at base, fluted, bowl shape. Very stiff.

SPITZBERGEN 1W-Y
Perianth white, smooth. Segments flat, pointed, broad, shovel shaped, well overlapping. Corona pale clear lemon yellow paling a little with age. Stiff, well balanced flower.

SPLIT 11W-W
Perianth creamy white, creased, crêpey texture, pointed outline. Segments flat, pointed, barely overlapping. Corona cream, split into 6 uneven segments, deeply frilled edge, ugly. Neck grows badly in water.

SPRING BEAUTY 3Y-O
Perianth pale creamy primrose, creased, crêpey texture. Segments flat, pointed, overlapping. Corona bright burnt orange.

SPRING FEVER 2Y-O
Discarded - Short, rough, flowers below foliage.

SPRING GLORY 1W-Y
Perianth creamy white stained yellow at base. Segments twisted, pointed, separated, giving spidery appearance. Corona deep lemon yellow.

SPRING MORNING 1Y-Y
Perianth pale sulphur, a little green on back, pointed outline. Segments slightly twisted, pointed, overlapping. Corona mid lemon, deeply cut crenate edge, flanged. Neck grows in water.

- SPRY** 2Y-R
Perianth clear lemon, paler at tips, suffused orange at base, edges curl in a little. Segments flat, rounded, broad, overlapping. Corona deep burnt orange red, slightly fluted, lightly frilled, straight sided, small in relation to perianth. Very stiff. Corona edge dies 3 days before perianth. Neck grows in water.
- SQUIRE** 1Y-Y
Perianth bright lemon, smooth. Segments rounded, overlapping at base. Corona slender, waisted, crenate edge. Very good flower substance.
- STADIUM** 2W-Y
Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona pale lemon yellow, with white blotches on edge, slightly fluted, very broad mouth, badly proportioned.
- STAINLESS** 2W-W
Perianth creamy white, very slightly creased, raised centre line. Segments rounded, overlapping at base, curled in at edges. Corona whiter, bowl-shaped, fluted. Very good flower substance. Neck grows badly in water.
- STANDARD BEARER** 1Y-Y
Discarded - Hangs head.
- STANDARD VALUE** 1Y-Y
Perianth sulphur yellow, top 3 segments incurved, creased. Segments flat, pointed, overlapping. Corona bright lemon yellow, broad mouth. Badly proportioned. Neck grows in water.
- STATUE** 2W-Y
Perianth creamy white, slightly creased, very flat, pointed outline. Segments flat, pointed, overlapping. Corona pale clear lemon yellow shade paler at base, fluted, almost straight sided. Large but well proportioned. Neck grows stiffly in water.
- STELLA** 2W-Y
Perianth creamy white, edges curl in at tips and curl back at base, thin star shape. Segments twisted, pointed, separated. Corona opens primrose, develops creamy white, lightly fluted, deep bowl shape. Neck grows badly in water.
- STERLING** 1Y-Y
Perianth bright lemon, slightly creased. Segments flat, pointed, overlapping. Corona deep lemon gold, crenate, rolled back edge, slender. Remains slightly hooded in water.
- STILTON** 9W-YYR
Perianth white. Outer segments very broadly ovate, inner ovate, and overlapping. Corona medium sized frilled cup, yellow with a green eye and a red rim. Neck grows badly in water.
- STOREEN** 9W-YYO
Perianth creamy white, edges curled in a little. Segments flat, rounded, well overlapping. Corona gold changing to edge of bright orange for 1/4 of its length, with deeper gold eye.
- STORMONT** 1W-W
Discarded - Short, hangs head, flowers below foliage.
- STOUT LAD** 2Y-Y
Discarded - Poor colour.
- STRAIGHT** 1W-Y
Perianth creamy white, creased, crêpey. Segments flat, pointed, overlapping. Corona creamy lemon shading to paler base, slightly frilled, flanged, neat edge.
- STRAIGHT FLUSH** 2Y-O
Perianth yellow. Segments very broadly ovate and overlapping. Corona small/medium sized frilled cup, orange in colour. Neck grows in water.
- STRAY PINK** 2W-P
Perianth cream, creased, crêpey texture, edges curl in, star shape. Segments flat, pointed, overlapping. Corona opens deep peach, develops bright solid rose pink, a shade paler at base, trumpet shape. Hangs head badly. Neck grows in water.
- STRESA** 1W-W
Perianth creamy white, green stripes on back, stained yellow at base. Segments twisted, pointed, narrow, barely overlapping. Corona opens clear lemon yellow, develops to pale cream, deeply frilled.
- STROMBOLI** 2W-O
Perianth poeticus white, edges incurving, head hangs down. Segments rounded, some notched, overlapping. Corona bright orange, flat, creased, ragged edge. Neck grows badly in water.
- STRONGHOLD** 1Y-Y
Perianth sulphur yellow, creased, crêpey texture. Segments flat, pointed, overlapping. Corona deeper lemon yellow, very broad mouth, coarse, badly proportioned.

- SUCCESSOR** 1Y-Y
Perianth lemon yellow, creased, crêpey texture. Segments flat, pointed, overlapping. Corona deep lemon yellow, straight sided, broad mouthed. Squat and badly proportioned, stamens well separated away from stigma.
- SUDA** 2W-O
Perianth creamy white. Segments flat, pointed, overlapping. Corona opens primrose develops apricot, developing paler colour with age, of trumpet proportions.
- SUDAN** 2Y-R
Perianth clear lemon, white tips, creased, crêpey substance. Segments flat, pointed, overlapping. Corona deep orange-red, fluted, bowl shape. Neck grows in water.
- SUILVEN** 3W-W
Perianth white. Segments broad and overlapping. Corona small, frilled cup, white with yellow rim. Neck grows badly in water.
- SULPHUR** 1Y-Y
Perianth soft sulphur yellow. Segments slightly twisted, pointed, overlapping. Corona clear lemon yellow, almost straight with rolled back edge.
- SULPHUR PRINCE** 1Y-Y
Perianth soft creamy sulphur yellow. Segments flat, pointed, overlapping. Corona pale clear lemon yellow.
- SUMATRA** 2Y-O
Perianth clear pale lemon yellow, one segment often buckled. Segments flat, pointed, broad, overlapping. Corona orange shading to gold base, fluted. Stiff.
- SUN CHARIOT** 2Y-O
Perianth bright lemon, paler at tips, overlaid slight orange at base, creased, crêpey, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona bright deep orange red, fluted, frilled, bowl shape. Neck grows in water.
- SUN DANCE** 1Y-Y
Perianth bright clear yellow, pointed outline. Segments almost flat, pointed, overlapping. Corona deep rich gold. Very broad mouthed. Barely trumpet length.
- SUN FIRE** 3Y-R
Perianth soft lemon overlaid unevenly with bright gold. Segments broad, rounded, overlapping at base, smooth, curled in at edges. Corona deep orange in lower half, developing to burnt orange upper half, reddish tinged rim; expanded, frilled edge. Very good flower substance. Neck grows badly in water.
- SUN SALUTE** 2Y-O
Discarded - Small, short, flowers below foliage.
- SUNBURST** 4Y-Y
No description available.
- SUNCREST** 2
Discarded - Flowers below foliage.
- SUNLIT HOURS** 2Y-Y
Perianth yellow. Segments broad, overlapping, with ridged texture. Corona medium sized frilled cup, same yellow as perianth. Neck grows in water.
- SUNNY BROOK** 2W-O
Perianth creamy white, a little floppy. Segments flat, pointed, broad, overlapping. Corona bright apricot orange, a little paler at base, fluted, extra frilled.
- SUNRIDGE** 2Y-OOY
Perianth yellow. Segments ovate and overlapping. Corona medium sized frilled cup, pale orange in colour. Neck grows badly in water.
- SUNRISE** 3W-YYO
Perianth opens primrose, develops creamy white, stained yellow at base with two yellow stripes on each segment, star shape. Segments flat, pointed, overlapping. Corona pale orange, shading to deeper edge, fluted, shallow bowl shape.
- SUNSILK** 2Y-Y
Perianth yellow. Segments very broadly ovate and overlapping. Corona medium sized frilled cup, slightly deeper yellow than perianth. Neck grows in water.
- SUNSTAR** 3W-R
Perianth creamy white stained yellow at base. Segments flat, rounded, well overlapping. Corona deep orange red shading through deep orange to greenish eye, fluted. Stiff.
- SVENGALI** 1W-Y
Perianth sulphury cream, stained yellow at base and green on back. Segments twisted, pointed, overlapping. Corona bright clear yellow, deeply lobed, wide mouthed.

SWANSDOWN 4W-W
 Perianth segments creamy white, flat, rounded, overlapping, one row complete. Corona cream, complete, filled with short twisted corona and stamen segments. The amount of doubling can vary to almost single flowers.

SWEET ADARE 1W-Y
 Perianth creamy white, stained yellow at base, crêpey substance. Segments twisted, pointed, barely overlapping. Corona primrose yellow, wide bell mouth.

SWEET CHARITY 2W-YYO
 Perianth creamy white. Segments very broad, rounded, overlapping at base, curled in at edges, fairly smooth, full rounded outline. Corona peach in lower half becoming deep cream in upper, fluted, coarsely frilled, deeply lobed edge. Neck grows in water.

SWEET MEMORY 2W-P
 Perianth creamy white, stained pale yellow base. Segments creased, buckled, crêpey, rounded and overlapping. Corona apricot 3/4 cream apricot edge 1/4. Fluted, sharply expanding flat type, deeply lobed. Slender stems, delicate. Neck grows badly in water.

SWEETHEART 2W-YYO
 Perianth creamy white, creased, a little floppy. Segments flat, rounded, overlapping. Corona primrose-apricot varying to creamy apricot, deeply frilled.

SWEETNESS 7Y-Y
 Perianth deep canary yellow, smooth. Segments flat, pointed, overlapping. Corona deeper canary yellow, straight sided.

SWISS MISS 2W-P
 Discarded - Flowers below foliage.

SYLVESTER 1W-W
 Perianth white. Segments broad and overlapping. Corona large frilled open mouthed and white in colour.

SYLVIA O'NEILL 3W-WWY
 Discarded - Low flower production.

SYRACUSE 3W-WWY
 Perianth white. Segments ovate and overlapping. Corona small frilled cup, white with yellow rim. Neck grows badly in water.

TABASCO 2
 Discarded - Severe virus in stock.

TAIN 1W-W
 Discarded - Short, flowers below foliage.

TAKORADI 4W-W
 Perianth white. Segments broad, round, well overlapping with slightly curled edges. Segments intermixed with cream coloured frilled corona segments. Neck grows in water.

TALLAND 2Y
 Perianth gold, slightly creased. Segments broad and overlapping. Corona orange, open mouthed, crenate. Neck grows badly in water.

TAMAR FIRE 4Y-R
 Perianth clear mid lemon, slight green back base. Segments rounded, overlapping, a little creased, some buckled, one complete row, six segments intermixed with deep orange corona segments. Neck grows badly in water.

TAMARA 2Y-Y
 Perianth lemon, outer segments broad, inner segments more ovate, segments twisted and overlapping. Corona lemon, well flared, edge frilled. Very long stems. Neck grows in water.

TAMINO 2Y-O
 Perianth soft pale gold, crêpey substance. Segments flat, rounded, well overlapping. Corona bright burnt orange red throughout, fluted. Neat.

TAMSYN 1Y-Y
 Perianth yellow. Segments rounded, overlapping and reflexed. Corona yellow with a crenate frilled edge, small flower, sturdy stem.

TANNAGHMORE 1W-Y
 Perianth creamy white, stained green-yellow back base. Segments pointed, overlapping, slightly creased, curled in at edges. Corona sulphur, paler at edges, slender with rounded flared mouth, lobed edge.

TANNHAUSER 2Y-YYO
 Perianth soft pale gold, paler at tips, creased, circular outline. Segments flat, rounded, broad, overlapping. Corona greenish lemon base, changing to orange red edge for 1/3 of its length, deeply fluted, frilled, lobed, almost flat.

- TARZAN** 2W-O
Perianth creamy white, stained yellow at base, creased, top 3 segments buckled, often caught in corona. Segments slightly twisted, pointed, overlapping. Corona lemon, suffused orange with narrow edge of pale lemon, some white blotches on edge, fluted. Very rough, large, ugly. Neck grows badly in water.
- TAWNY GOLD** 1Y-Y
Perianth yellow. Segments ovate, twisted and overlapping. Corona large frilled flared cylinder, same yellow as perianth. Neck grows in water.
- TAWNY LAD** 2Y-O
Perianth lemon with darker rim to segments, tinged orange, fading at edge and tips. Segments rounded, overlapping, smooth, curled in at edges. Corona burnt orange, reddish tinge, cup-shaped, wavy lobed edge.
- TEAROSE PHOENIX** 4Y-Y
Perianth segments apricot primrose, twisted, pointed, separated. A few corona segments, deeper apricot, intermixed with perianth segments which are tightly packed and twisted.
- TEBESSA** 3W-R
Perianth creamy white, smooth. Segments flat, pointed, broad, well overlapping. Corona deep solid orange red, fluted, flat, uneven edge.
- TEBOURBA** 3W-R
Perianth creamy white, edges curled in a little. Segments flat, pointed, broad, well overlapping. Corona deep bright orange red with greenish red eye, deeply fluted.
- TECOMA** 2Y-O
Perianth pale sulphur, slightly creased, slightly twisted. Segments acute, overlapping at base, fades at tips. Corona egg yolk yellow, open mouthed, crenate, rim lobed. Neck grows badly in water.
- TEHERAN** 2Y-O
Perianth bright lemon yellow, slightly creased, edges curl in a little. Segments flat, pointed, broad, well overlapping. Corona bright orange red, fluted, shallow bowl shape. Neck grows in water.
- TELAMONIUS PLENUS** 4Y-Y
Perianth segments clear yellow, green on back when first opening, twisted, pointed, separated, curled back at edges. Corona segments deep bright gold, narrow rounded segments tightly packed and intermixed evenly throughout with perianth segments.
- TELSTAR** 1W-W
Perianth white. Segments broad and overlapping. Corona large frilled funnel, same white as perianth. Neck grows in water.
- TEMPLEMORE** 2W-W
Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona opens creamy buff, develops cream with yellow green base, slightly fluted, frilled, neat. Neck grows in water.
- TEMPLEPATRICK** 2W-W
Discarded - Short, flowers below foliage.
- TEMPURA** 2Y-Y
No description available.
- TENEDOS** 2W-Y
Perianth creamy white, edges rolled in, star shaped. Segments flat, pointed, overlapping. Corona opens lemon yellow and develops to cream throughout, of almost trumpet proportion.
- TEXAS** 4Y-O
Perianth segments pale lemon yellow, flat, pointed, overlapping, 2 complete rows and some twisted segments in centre. Corona bright orange, many large segments intermixed tightly with perianth segments.
- THALIA** 5W-W
Perianth creamy white, segments flat, pointed, separated. Corona a little whiter than perianth, yellow inside base, style reached 1/2 length of stamens. Neck grows in water.
- THE FIRST** 1Y-Y
Perianth soft primrose yellow, green on back. Segments flat, pointed, overlapping. Corona deep lemon yellow. Weak neck, floppy flower. Neck grows in water.
- THE GIFT** 1Y-Y
Perianth yellow. Segments broad, overlapping and inflexed. Corona large frilled flared cylinder, slightly deeper yellow than perianth. Neck grows in water.
- THE KNAVE** 6W-Y
Perianth creamy white, Segments broad, rounded, overlapping at base, creased, strongly reflexed. Corona creamy yellow, darker towards top, straight, crenate edge.

THE SUN 3W-O
 Perianth pure white stained yellow at base. Segments flat, pointed, overlapping. Corona deep burnt orange red with deeper red eye, fluted. Style a little longer than longest stamens.

THERAPIA 3W-YYO
 Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona bright gold shading to bright orange edge for 1/3 of its length, fluted.

THERM 3Y-O
 Perianth sulphur yellow, paler at tips, slightly creased. Segments flat, rounded, overlapping. Corona bright orange red, a little paler at base, fluted, bowl shape. Neck grows in water.

THETIS 9W-R
 Discarded - Small, flowers below foliage.

THOMAS HARDY 9W-R
 Discarded - Small.

THOMAS MARK 2Y-YOO
 Perianth clear lemon yellow, 3 segments reflexed. Segments flat, pointed, overlapping. Corona deep yellow edged bright orange for 1/2 of its length, deeply frilled edge, almost flat. Badly proportioned with corona too wide for diameter of perianth.

THRILLER 2Y-R
 Perianth yellow. Segments very broadly ovate, overlapping and slightly inflexed. Corona large frilled cup, yellow with an orange rim. Neck grows in water.

THURSO 1W-W
 Discarded - Flowers below foliage, coarse, ugly flower, susceptible to basal rot.

TIBET 2W-W
 Perianth white, slightly creamy, slightly creased. Segments broad, ovate, overlapping extensively, edges slightly incurling. Corona white, slightly creamy, short, open mouthed with slightly curled back rim, deeply frilled edge. Neck grows in water.

TILLA DURIEUX 3W-GYO
 Perianth creamy white stained yellow at base, creased. Segments flat, rounded, overlapping. Corona orange gold shading to orange red edge for 1/2 its length, and a greenish gold eye, fluted.

TINKER 2Y-O
 Perianth deep lemon, twisted. Segments edges incurling, small, ovate, slightly overlapping. Corona deep orange, straight sided, slightly ribbed, edge slightly frilled. Neck grows badly in water.

TINO ROSSI 2Y-O
 Perianth cream, stained slight yellow at base, one or two segments usually creased, rounded outline. Segments flat, rounded, overlapping. Corona bright deep orange, fluted, widespread.

TINSEL 3W-WWY
 Perianth creamy white, creased, edges curl in. Segments flat, pointed, overlapping. Corona greyish green eye, changing to pale greenish primrose with pencil edge of apricot orange, fluted. Neck grows in water.

TINTORETTO 1Y-Y
 Perianth pale primrose yellow, crêpey texture. Segments flat, pointed, broad, overlapping. Corona deep lemon developing to pinkish buff with yellowish edge. Often has small white blotches on edge of corona.

TITANIA 6W-W
 Discarded - Small, short, cyclamineus hybrids.

TITCH 1Y-Y
 Perianth yellow. Segments acute, twisted with no overlap. Corona large frilled flared, open mouthed, same colour as perianth.

TITOKI 1
 Perianth yellow. Segments ovate, twisted with slight overlap. Corona large frilled flared cylinder, slightly deeper yellow than perianth.

TITTLE-TATTLE 7Y-GYY
 Discarded - Small, jonquilla hybrid.

TIVOLI 2Y-R
 Perianth bright lemon with paler tips, slightly creased, crêpey, pointed outline. Segments flat, pointed, overlapping. Corona deep orange, paler at base, fluted, frilled, lobed, rather uneven edge, bowl shape.

TOLEDO 2
 Discarded - Small, rough flower.

TOORAK GOLD 2Y-Y
Perianth lemon, slightly hooded, slightly upward facing. Segments notched, overlapping. Corona short relative to flower size, open mouth with edge rolled back, gently lobed. Neck grows stiffly in water.

TOPAZ 7W-Y
Discarded - Flowers below foliage, jonquilla hybrid.

TOPIC 2W-P
Perianth dirty white, creased and floppy. Segments flat, pointed, broad, well overlapping. Corona pale buff primrose with thin salmon pink edge, ageing to pale salmon buff. Neck grows in water.

TOP-NOTCHER 1Y-Y
Perianth yellow. Segments ovate and overlapping. Corona medium sized frilled flared cylinder, same yellow as perianth. Neck grows badly in water.

TOPSCORE 2
Perianth bright lemon, paler at tips, creased, edges curl in. Segments flat, pointed, overlapping. Corona bright orange, fluted, frilled, deep bowl shape. Stiff. Neck grows in water.

TORCHDANCE 2Y-Y
Perianth bright lemon, slightly creased, edges curl in, pointed outline. Segments flat, pointed, overlapping. Corona deeper lemon, frilled, almost straight sided, neat. Stiff.

TOREADOR 3W-R
Perianth white, broad, overlapping. Segments edges incurling. Corona orange, flat, ragged frilled edge. Neck grows stiffly in water.

TORERO 2Y-O
Discarded - Severe virus in stock.

TORNAMONA 2W-W
Perianth creamy white, slightly creased and crêpey, pointed outline. Segments flat, pointed, overlapping. Corona whiter than perianth, lightly frilled, neat, trumpet proportion. Neck grows in water.

TORQUAY 2Y-O
Perianth pale yellow. Segments very broadly ovate and well overlapping. Corona small frilled cup, orange in colour. Neck grows in water.

TOSCANINI 2W-WWP
Perianth creamy white, creased, catches in corona, edges notched. Segments twisted, pointed, overlapping. Corona salmon pink shading to paler colour at base. Rough and ugly.

TOUJOURS 1Y-Y
Perianth soft mid-lemon. Segments somewhat creased, rounded, overlapping. Corona deeper, bright lemon, top half fluted, becoming corrugated in places.

TRAMORE 2W-Y
Perianth creamy white stained yellow at base, slightly creased, pointed outline. Segments flat, pointed, overlapping. Corona bright clear lemon, frilled, crenate, trumpet proportion. Very stiff.

TREDORE 3Y-O
Perianth opens creamy primrose yellow, develops to cream, stained yellow at base. Segments flat, pointed, broad, overlapping. Corona deep orange red, fluted.

TREEN 2Y-OOR
Discarded - Poor colour, long neck.

TREFFRY 3W-O
Perianth white stained yellow at base, a little creased. Segments flat, rounded, overlapping. Corona deep greenish lemon, shading to bright orange red edge for 1/2 its length, fluted.

TRENANCE 2Y-Y
Perianth soft pale lemon yellow, white tips. Segments flat, pointed, broad, well overlapping. Corona clear deep lemon yellow. Of almost trumpet proportions.

TRENITHON 1Y-Y
Perianth yellow. Segments very broadly ovate, overlapping and slightly inflexed. Corona large frilled flared cylinder, same yellow as perianth. Neck grows in water.

TRENOON 2Y-Y
Discarded - Small.

TRESAMBLE 5W-W
Perianth creamy white. Segments pointed, broad, overlapping. Corona cream with paler edge, yellow green base. Very long style.

TRESITHICK 2Y-Y
Discarded - Poor flower.

- TRESKERBY** 3Y-R
Perianth cream stained yellow at base, slightly hooded. Segments flat, pointed, barely overlapping. Corona burnt orange shading to gold base.
- TRESKEWES** 5Y-Y
Perianth pale greenish lemon yellow, a little creased, edges curl back. Segments flat, pointed, separated. Corona pale greenish lemon yellow, slightly fluted, open mouthed. Neck grows badly in water.
- TRESSERVE** 1Y-Y
Perianth sulphur yellow, smooth, star-shaped. Segments twisted, pointed, barely overlapping. Corona deep lemon yellow, with deep crenate edge.
- TREVISKY** 2Y-O
Perianth clear lemon yellow, white tips, crêpey substance. Segments flat, pointed, overlapping. Corona clear orange red, fluted, almost straight sided.
- TREVITHIAN** 7Y-Y
Perianth lemon yellow, crêpey texture. Segments flat, rounded, overlapping. Corona bright lemon yellow, well balanced.
- TREWINCE** 2Y-Y
Perianth pale lemon yellow, slightly creased. Segments twisted, pointed, overlapping. Corona bright deep lemon yellow.
- TREWIRGIE** 6Y-Y
Perianth bright deep lemon yellow, smooth, reflexed. Segments flat, pointed, broad, overlapping. Corona deeper bright lemon yellow. Well proportioned.
- TREWITHEN** 1Y-Y
Perianth sulphur, slightly creased. Segments would fit into a double triangle. Corona deep sulphur, straight sided, opening slightly at rim, crenate, lightly frilled. Well proportioned flower. Weak, hollow stems. Neck grows in water.
- TRIANON** 3W-Y
Perianth white. Segments broad and overlapping. Corona large frilled yellow bowl.
- TRIFINE** 2Y-O
Perianth lemon, creased, slightly twisted. Segments triangular, only overlapping at bottom third of segments, colour fades at tips. Corona gold to orange, open mouthed, crenate and slightly frilled. Neck grows in water.
- TROCADERO** 1W-Y
Discarded - Untidy corona edge, perianth catches in corona.
- TROSTAN** 1W-Y
Perianth creamy white, creased. Segments slightly twisted, pointed, overlapping. Corona cream, crenate, rolled back edge.
- TROUBADOUR** 2W-P
No description available.
- TROUSSEAU** 1W-Y
Perianth creamy white. Segments flat, pointed, broad, overlapping. Corona opens clear lemon yellow, develops to rich deep buff, neat shape and edge, well proportioned.
- TRUELOVE** 3W-R
Perianth poeticus white, stained yellow at base, edges curl in, a little creased. Segments flat, pointed, broad, overlapping. Corona deep sealing wax red, shading paler at base, fluted. Stiff. Neck grows stiffly in water.
- TRUMPETER** 1Y-Y
Perianth gold, smooth, slightly twisted, slightly hooded. Segments small and neat, overlapping to half length. Corona gold, flanged with neat frill. Overall crisp and neat appearance. Neck grows in water.
- TRUTH** 2W-W
Perianth pure white, perfect double triangle, smooth. Segments flat, pointed, broad, well overlapping, shovel shaped. Corona white, straight with rolled back edge, almost trumpet proportion.
- TRYST** 2W-W
Perianth creamy white, slightly creased, circular outline. Segments flat, rounded, broad, well overlapping. Corona opens primrose, develops cream, almost pinched in, fluted. Neck grows stiffly in water.
- TUDOR KING** 1W-Y
Discarded - Short, susceptible to basal rot.
- TUDOR MINSTREL** 2W-Y
Perianth creamy white, stained yellow at base, slightly creased. Segments flat, pointed, broad, overlapping. Corona chrome yellow, slightly fluted, lightly frilled, bowl shape. Hangs head when young. Neck grows badly in water.

- TULLYGARLEY** 2Y-R
Perianth deep lemon yellow, crêpey texture, slightly creased. Segments flat, rounded, broad, well overlapping. Corona deep burnt orange red, fluted. Hangs head badly. Neck grows in water.
- TULLYGLASS** 2W-WWY
Perianth creamy white. Segments broad and overlapping. Corona medium sized frilled cup, creamy yellow with yellow rim. Neck grows in water.
- TULLYROE** 2W-R
Perianth poeticus white, smooth, edges incurled. Segments would fit into a double triangle if flat, triangular shape. Corona dark orange, short and slightly flat, slightly lobed rim. Neck grows badly in water.
- TULLYROSE** 2W-P
No description available.
- TULLYROYAL** 2W-P
Perianth creamy white, circular outline. Segments rounded, overlapping, creased, some buckled, crepey particularly at edges. Corona moss green throat, cream base developing to pale salmon upwards, sometimes white blotches; expanded, frilled edge.
- TULYAR** 3W-R
Perianth white. Outer segments rounded, inner very broadly ovate and well overlapping. Corona small frilled orange cup. Neck grows badly in water.
- TUNIS** 2W-WWY
Perianth muddy cream, creased. Segments twisted, pointed, overlapping. Corona opens lemon yellow, develops to buff, deeper at edge.
- TURIN** 3W-GOO
Perianth white, stained yellow at base, creased. Segments flat, pointed, broad, overlapping. Corona orange gold shading to orange red for 1/2 its length, almost flat, fluted.
- TUSKAR LIGHT** 2W-YYR
Discarded - Long neck, poor poise, very susceptible to basal rot.
- TUTANKHAMUN** 2W-GWW
Perianth white, stained green at base. Segments rounded, overlapping at base, creased, curled in at edges, segments broad with star-shaped outline. Corona moss green throat, then white, expanded, frilled and flanged edge. Very good flower substance.
- TWINK** 4W-O
Perianth segments cream, stained yellow at base, twisted, pointed, separated, 2 complete rows, outer segments curled back at edges. Corona segments orange intermixed throughout, with 2 or 3 twisted perianth segments in centre.
- ULSTER BEAUTY** 2W-WWY
Perianth creamy white, slightly creased, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona opens pale lemon, develops deep cream shading to creamy lemon edge, lightly frilled, almost straight sided. Neck grows in water.
- ULSTER MAIDEN** 2W-R
Perianth white. Segments broad and overlapping. Corona small frilled orange cup. Neck grows badly in water.
- ULSTER PRINCE** 1Y-Y
Perianth pale lemon, crêpey. Segments flat, pointed, overlapping. Corona deeper lemon, frilled, flanged, neat. Stiff.
- ULSTER QUEEN** 1W-W
Perianth creamy white, stained green/yellow back base. Segments rounded, overlapping at base; bottom segments very broad, slightly creased, narrower top segments twisted, notched. Corona slightly darker, moss green at throat, ivory white, wide mouth, lobed edge. Very good flower substance.
- ULTIMUS** 2Y-O
Perianth pale lemon, edges incurling. Segments acute, well overlapping. Corona orange, short, open mouthed, rim frilled. Neck grows badly in water.
- UNIQUE** 4W-Y
Perianth creamy white. Segments broad, well rounded, overlapping, crepey; one complete row with nine buckled segments, pointed forwards, intermixed with bright pale lemon corona segments, edged deep cream, frilled edge; compact centre. Weak stems.
- UNSURPASSABLE** 1Y-Y
Perianth clear bright lemon yellow, crêpey texture. Segments twisted, pointed, overlapping, curled back at edges giving thin appearance. Corona deep golden yellow.
- URANIA** 3
Perianth yellow. Segments broad and overlapping. Corona medium sized frilled orange cup. Neck grows in water.

- URSA 2W-WPP
Perianth white. Segments broad and overlapping. Corona medium sized, frilled cup, pink with white centre. Neck grows in water.
- VAHSEL BAY 2W-W
Perianth creamy white, creased, crêpey, pointed outline. Segments flat, pointed, overlapping. Corona cream, deeply frilled, crenate, flanged, broad mouth, dumpy appearance.
- VALENCIA 4Y-O
Perianth segments pale lemon, flat, pointed, separated, 3 complete outer rows with a few twisted segments in centre, evenly intermixed with pale orange corona segments. Amount of doubling varies.
- VALIANT SPARK 2Y-O
Perianth deep lemon yellow, crêpey texture, creased, top 3 segments buckled, notched. Segments slightly twisted, pointed, overlapping. Corona deep solid burnt orange, fluted. Poor poise. Neck grows badly in water. Corona edge dies first.
- VAN WAVEREN'S GIANT 1Y-Y
Perianth creamy sulphur yellow, deeper yellow on back, stained deep yellow at base. Segments twisted, pointed, barely overlapping. Corona bright lemon yellow.
- VAN WERELD'S FAVOURITE 1W-Y
Perianth dirty creamy white, stained yellow at base. Segments flat, pointed, broad, overlapping. Corona clear lemon yellow.
- VARIANT 2Y-R
Perianth yellow. Outer segments broad, inner ovate and overlapping, with a ridged texture. Corona large frilled red cup. Neck grows badly in water.
- VARNA 3Y-YOO
Perianth opens pale lemon yellow, develops to cream stained yellow at base, smooth. Segments flat, pointed, overlapping. Corona bright orange red, shading to orange gold at base, occasional white blotches on edge, goblet shape, stiff and sturdy.
- VELVETEEN 2Y-Y
Perianth light lemon yellow, creased. Segments flat, pointed, overlapping. Corona gold, overlaid orange for half its length, colour a little variable, straight sided.
- VENUS 2W-YYO
Discarded - Rough, floppy flower.
- VERDI 1W
Perianth pale lemon yellow. Segments broad and overlapping. Corona large frilled flared and deeper yellow in colour.
- VERGER 3W-R
Perianth pure white stained slightly yellow at base, crêpey texture, a little creased. Segments flat, rounded, overlapping. Corona bright orange red, flat, fluted. Neck grows in water.
- VERONA 3W-W
Perianth white. Segments very broadly ovate and overlapping. Corona medium sized frilled bowl, pale yellow. Neck grows in water.
- VICLAND 2W
Perianth opens primrose, develops cream stained deep yellow at base, creased, sometimes buckled, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona bright lemon base 1/2 shading to bright orange red edge, fluted, frilled, lobed, bowl shape. Neck grows in water.
- VICTOR BORGE 1W-Y
Discarded - Ugly, poor perianth, flowers below foliage.
- VICTORIA 1W-Y
Perianth cream stained yellow at base. Segments flat, pointed, overlapping. Corona bright lemon yellow.
- VICTORIOUS 2W-Y
Perianth creamy white, slightly creased. Segments flat, pointed, broad, overlapping. Corona opens pale lemon yellow, developing to pale creamy sulphur with thin lemon rim.
- VIGIL 1W-W
Perianth white. Outer segments broad, inner ovate and overlapping. Corona large frilled flared cylinder, same white as perianth.
- VIKING 1Y-Y
Perianth egg yolk yellow, smooth, rounded, flower semi-upward pointing. Segments overlapping to 3/4 of length. Corona long, fairly narrow, almost straight sided, egg yolk yellow, edge deeply frilled. Stems can be hollow. Neck grows stiffly in water.

VILLAGE BEAUTY 3W-YY
 Perianth creamy white, stained yellow at base, creased, reflexed. Segments flat, pointed, barely overlapping. Corona greenish gold with narrow rim of orange red, fluted, uneven lobed edge. Neck grows in water.

VILNA 2W-W
 Discarded - Poor looking flower.

VIRTUE 2Y-Y
 Perianth clear lemon yellow, crêpey substance. Segments flat, pointed, overlapping. Corona deeper yellow, of trumpet proportions. Stiff and sturdy.

VIVEX 2Y-YYO
 Perianth yellow. Segments rounded and overlapping. Corona medium/large frilled bowl, yellow with orange rim.

VIXI 2Y-Y
 No description available.

VOLGA 2W-W
 Perianth white, sometimes notched. Segments flat, pointed, broad overlapping. Corona opens cream, develops creamy white. Very stiff, hangs head.

VOLTERRA 2Y-O
 No description available.

VULCAN 2Y-O
 Perianth deep lemon, slightly reflexed. Segments ovate to rounded, overlapping. Corona orange, straight sided, slightly frilled rim. Neck grows badly in water.

W.P.MILNER 1W-W
 Perianth creamy sulphur, green stripe on back, stained yellow at base. Segments twisted, pointed, separated. Corona opens pale lemon yellow, develops to pale primrose.

WAHKEENA 2W-Y
 Perianth white. Segments broad and overlapping. Corona medium sized frilled cup, yellow in colour. Neck grows in water. Weak stems.

WALT DISNEY 2Y-O
 Perianth deep clear lemon yellow, crêpey substance, star shaped. Segments flat, pointed, overlapping. Corona bright solid orange, often white blotches on edge. Neck grows in water, corona edge dies first.

WANDIN GLORY 1Y-Y
 Perianth yellow. Segments ovate and overlapping. Corona medium sized frilled open mouthed, same yellow as perianth. Neck grows badly in water.

WARIALDA 1Y-Y
 Perianth yellow. Segments ovate, twisted with slight overlap. Corona large flared, frilled, and same yellow as perianth.

WARLOCK 2W-YYO
 Perianth creamy white, stained yellow at base, creased. Segments flat, pointed, overlapping. Corona bright gold, shading to orange edge for 1/3 of its length.

WARM SPRINGS 2Y-YYO
 Perianth soft pale sulphur yellow, creased, a little floppy. Segments flat, pointed, well overlapping. Corona deep gold shading to orange edge for 1/4 its length, almost flat, fluted.

WARRIOR 2Y-O
 Perianth bright lemon yellow, smooth. Segments flat, pointed, overlapping. Corona bright orange red, shade paler at edge. Neck grows in water.

WARWICK 1Y-Y
 Perianth yellow. Segments broad and overlapping and slightly inflexed. Corona large frilled flared, same yellow as perianth. Neck grows in water.

WATERPERRY 7W-PPP
 Perianth muddy cream, creased, crêpey, edges curl in. Segments flat, pointed and overlapping. Corona opens primrose, develops buff yellow with peach edge of 1/4 its length. Neck grows badly in water.

WATERVILLE 2W-W
 Perianth creamy white, slightly creased, edges curl in a little, circular outline. Segments flat, rounded, broad, well overlapping. Corona creamy primrose deeper at edge, greenish gold eye, fluted, shallow bowl shape.

WEARDALE PERFECTION 1W-Y
 Perianth opens pale primrose, develops dirty white, stained yellow at base, crêpey texture. Segments twisted, pointed, barely overlapping. Corona deep bright lemon yellow. Small but well proportioned.

WEDDING BELL 2W-W
 Perianth creamy white, smooth, pointed outline. Segments flat, pointed, overlapping. Corona creamy white, almost straight edge, flanged, neat. Neck grows in water.

WEINER BLUT 2W-O
 No description available.

WEISSHORN 1W-W
 Discarded - Short.

WELCOME (NL)
 Perianth rich golden. Segments rounded, overlapping. Corona rich golden, expanded, flanged, deeply crenate. Neck grows in water.

WELL-BORN 2W-W
 Perianth creamy white, slightly creased. Segments flat, pointed, broad, well overlapping. Corona cream, of trumpet proportion.

WEXFORD 2Y-Y
 Perianth lemon yellow, a little crêpey. Segments flat, pointed, broad, well overlapping. Corona deeper lemon yellow, barely trumpet length. Stiff and neat.

WHANG-HI 6Y-O
 Discarded - Small, short, cyclamineus hybrid.

WHITE CHIEF 1W-W
 Perianth poeticus white, creased and twisted. Segments broad and straight sided. Corona poeticus white, flanged, lightly frilled rim. Hollow stems. Neck grows a little in water.

WHITE CITY 3W-W
 Perianth creamy white, creased. Segments flat, pointed, broad, overlapping. Corona pale creamy yellow with greyish eye, fluted.

WHITE EMPEROR 1W-W
 Perianth white. Segments flat, pointed, overlapping. Corona opens deep buff primrose, develops to cream. Neat.

WHITE GOLD 2W-Y
 Perianth creamy white, stained slightly yellow at base, darker back base. Segments broad, rounded, overlapping at base, slightly creased, curled in at edges. Corona bright lemon yellow, fluted upper half, wide belled mouth, crenate lobed edge. Neck grows in water.

WHITE LADY 3W-Y
 Perianth creamy white, star-shaped. Segments flat, pointed, barely overlapping. Corona opens pale lemon develops to almost white with lemon base. Neck grows in water.

WHITE LION 4W-WYY
 Perianth segments creamy white, flat, pointed, broad, overlapping, complete. Corona segments open primrose yellow, develop to creamy buff, broad, intermixed evenly with the broad perianth segments. A solid, well shaped flower.

WHITE MAGIC 2W-GWW
 Perianth white. Segments broad, overlapping with a ridged texture. Corona medium sized frilled funnel, same white as perianth. Neck grows in water.

WHITE MAIDEN 2W-W
 Discarded - Severe virus in stock.

WHITE MARVEL 4W-W
 Perianth white. Segments ovate and double. Corona creamy white, double, and intermixed with inner perianth whorl. Neck grows badly in water.

WHITE NILE 2W-W
 Perianth creamy white, creased. Segments flat, pointed, overlapping. Corona opens primrose yellow, develops to pale cream. Very long style.

WHITE PLUME 2W-W
 Perianth white, slightly creased and crêpey, bottom segments reflex, circular outline. Segments flat, rounded, overlapping. Corona creamy white with moss green eye ageing to greyish green, deeply fluted, frilled, deeply lobed, straight sided, hangs head when young. Neck grows in water.

WHITE PRINCE 1W-W
 Perianth white. Segments very broadly ovate, overlapping and slightly inflexed. Corona medium sized, frilled, rolled and open mouthed, same white as perianth. Neck grows badly in water.

WHITE PROSPECT 1W-W
 Perianth white. Outer segments very broadly ovate, inner ovate and overlapping. Corona large frilled, flared funnel, same colour as perianth. Neck grows in water.

- WHITE QUEEN** 2W-W
Perianth poeticus white, very creased. Segments catch in corona. Segments twisted, rounded, overlapping. Corona buff primrose with paler edge, very deeply frilled.
- WHITE SAIL** 4W-W
Perianth opens deep cream develops creamy white, with cream centre stripe, flat, rounded, overlapping, 1 row complete with 6-8 narrow twisted segments in centre intermixed with very short cream corona segments. Crêpey, creased, circular outline. Amount of doubling variable.
- WHITE SPIRE** 2W-W
Perianth white. Outer segments very broadly ovate, inner ovate, and overlapping. Corona medium sized frilled cup, white, with green eye. Neck grows badly in water.
- WHITE SPRITE** 3W-W
Perianth white. Outer segments rounded, inner ovate and overlapping. Corona small frilled cup, white with green eye. Neck grows badly in water.
- WHITE TARTAR** 1W-W
Discarded - Coarse flower, poor poise, weak neck.
- WHITE TRIUMPHATOR** 1W-W
Perianth creamy white, slightly creased. Segments flat, pointed, broad, overlapping. Corona opens buff primrose develops to pale creamy white. Rather large and coarse.
- WHITE WAX** 1W-W
Perianth muddy cream, stained yellow on back, pointed outline. Segments flat, pointed, overlapping. Corona primrose yellow, developing cream, slender. Rough.
- WHITEHEAD** 2W-W
Perianth creamy white, stained green back base. Segments rounded, overlapping, bottom segments very broad, slightly creased, curled in at edges. Corona moss green throat, pale cream, expanded, frilled edge. Very good flower substance. Neck grows in water.
- WHITELEY GEM** 2Y-O
Perianth clear soft yellow, creased, perianth segments often caught in corona. Segments flat, pointed, barely overlapping. Corona bright orange red, shading to gold base. Neck grows in water.
- WHITE'S HYBRID** 8W-YYR
Discarded - Short, flowers below foliage.
- WHITEWELL** 2W-Y
Perianth creamy white, stained yellow at base, crêpey texture. Top segments long and pointed, lower segments shorter and rounded. Segments flat, pointed, overlapping. Corona orange gold, fluted.
- WIELAND WAGNER** 2W-Y
Discarded - Ugly flower, short.
- WILD ROSE** 2W-P
Perianth creamy white, slightly creased. Segments flat, pointed, broad, overlapping. Corona clear rose pink with slightly deeper edge, and greenish eye. Well balanced.
- WILL SCARLETT** 2W-O
Perianth opens primrose, develops creamy white stained yellow at base, edges curl back. Segments twisted, pointed, separated. Corona light orange, shading to orange edge for ½ its length, fluted, deep lobed uneven edge. Hangs head badly. Neck grows in water.
- WILLEM MENGELBERG** 1
Perianth bright lemon yellow, crêpey texture, one segment often buckled. Segments flat, pointed, overlapping. Corona deeper lemon yellow, very broad mouth, badly proportioned.
- WILLEMSTAD** 2W-O
Perianth creamy white, deeply creased, buckled, notched, edges curl in. Segments flat, pointed, overlapping. Corona bright orange, deeply fluted, unevenly lobed edge. Neck grows in water.
- WILLIAM FARMER** 2Y-Y
Perianth clear lemon, smooth, a little crepey. Segments rounded, broad, overlapping at base, curled in at edges, star shaped outline. Corona mid lemon, paler at base, wide, rounded belled mouth, even crenate edge.
- WILLIAM THE SILENT** 1Y-Y
Perianth lemon yellow, creased, crêpey texture. Segments twisted, pointed, overlapping. Corona deep lemon yellow, very broad mouth, badly proportioned.
- WILLOWFIELD** 3W-R
Perianth creamy white, stained yellow at base, creased, buckled, edges curl in. Segments slightly twisted, pointed, overlapping. Corona deep gold base shading to deep orange red, shallow bowl shape, fluted. Corona edge dies first. Neck grows badly in water.

WIN ALL 3Y-YYR
Discarded - Low flower yield.

WINCHESTER 2Y-YOO
Perianth bright lemon, creased, crêpey, edges curl in a little. Segments flat, pointed, broad, overlapping. Corona orange shading to orange red edge for ½ its length, fluted, frilled, goblet shape. Stiff. Neck grows in water.

WINDBLOWN 4W-Y
No description available.

WINGADEE 2W-OOY
Perianth cream, slightly reflexed but edges of segments incurling. Segments broad, rounded but with prominent mucronate tip, overlapping on ¾ of length. Corona egg yolk yellow at base, deep lemon at edge, short, open mouthed, crenate, with lobed edge. Hollow stems. Neck grows stiffly in water.

WINGED VICTORY 6W-Y
Discarded - Short, cyclamineus hybrid.

WINIFRED VAN GRAVEN 3W-YYR
Perianth poeticus white, stained slight yellow at base, edges curl in a little. Segments flat, pointed, broad, well overlapping. Corona greenish lemon, changing to bright orange red edge for 1/3 its length, fluted. Very long style. Stiff and sturdy.

WINKFIELD'S DOWER 1Y-Y
Perianth pale sulphur yellow, slightly creased, stiff. Segments slightly twisted, pointed, overlapping. Corona bright yellow.

WINNIE WEEDON 9W-YYR
Perianth creamy white, edges curled in, star shaped. Segments flat, pointed, overlapping. Corona lemon yellow, changing to bright red edge for 1/4 of its length, with small deeper lemon eye.

WINNIPEG 2Y-YYO
Perianth sulphur, deeply creased, sometimes buckled. Segments flat, pointed, overlapping. Corona light greenish gold shading to bright orange edge for 1/2 its length, fluted, frilled, shallow bowl shape. Neck grows in water.

WINTER BELLS 8W-Y
Perianth white. Segments broad and overlapping. Corona small lobed yellow cup. Multiheaded.

WINTER GOLD 1Y-Y
Perianth deep golden yellow, some green on back. Segments twisted, pointed, barely overlapping. Corona deep rich gold, almost straight.

WINTERMOON 1W-Y
Perianth white. Segments ovate, spreading and overlapping. Corona large frilled open mouthed, yellow in colour.

WINWICK 1W-W
Perianth white. Segments broad, overlapping and inflexed. Corona large frilled, rolled funnel, same colour as perianth. Neck grows badly in water.

WISDOM 3W-W
Perianth creamy white, slightly creased, edges curled in a little. Segments flat, rounded, well overlapping. Corona cream with slight buff edge and grey-green eye, whitens with age. Stiff and sturdy.

WODAN 2W-YYO
Discarded - Poor colour, untidy corona edge, flowers below foliage.

WOODCOCK 6Y-Y
Perianth bright clear yellow, reflexed, crêpey texture. Segments flat, pointed, broad, overlapping. Corona deeper yellow. Well proportioned.

WOODGREEN 2W-Y
Perianth white with yellow tinge to edges, slightly reflexed and twisted. Segments overlapping at base only. Corona primrose, paler at base becoming darker at edge, open mouthed, only slightly frilled. Neck grows in water.

WOODLAND PRINCE 3W-Y
Perianth creamy white. Segments rounded, overlapping at base, bottom segments broad, top segments narrower, curled in deeply at edges, slightly creased. Corona cream, top one-fifth developing to pale lemon, fluted. Neck grows in water.

WOODLAND STAR 3W-R
Perianth creamy white, stained yellow base, smooth, raised centre line. Segments rounded, overlapping at base, curled in at edges. Corona moss green throat, deep warm gold in lower half changing through orange to reddish rim; shallow bowl-shaped, fluted edge. Neck grows badly in water.

- WOODPECKER** 2W
Perianth opens primrose, develops cream stained yellow at base, slightly creased, edges curl in at tips, slightly reflexed. Segments flat, pointed, overlapping. Corona gold 1/4 shading to bright orange edge, fluted, lightly frilled, bowl shape. Neck grows in water.
- WOODVALE** 2W-WWY
Perianth creamy white, smooth, pointed outline. Segments flat, pointed, broad, overlapping. Corona opens primrose, develops cream, barely frilled, slender trumpet proportion. Neat. Neck grows in water.
- WOONAN** 2Y-R
Perianth yellow. Segments broad and well overlapping. Corona small frilled orange/red cup. Neck grows badly in water.
- WRESTLER** 1Y-Y
Perianth muddy greenish sulphur yellow, green on back. Segments twisted, pointed, overlapping. Corona bright deep lemon yellow.
- YALTA** 3W-YYR
Perianth creamy white, creased, edges curled in. Segments flat, pointed, broad, well overlapping. Corona orange gold changing to deep orange red edge for 1/4 of its length, fluted. Stiff and sturdy.
- YANKEE CLIPPER** 2Y-YYO
Perianth opens primrose with pale tips, develops cream, creased, often notched, sometimes buckled. Segments flat, pointed, overlapping. Corona gold base shading to deep orange, fluted, frilled, shallow bowl shape. Neck grows badly in water.
- YELLOW BEAUTY** 1Y-Y
Perianth clear lemon yellow, smooth, very pointed, star shaped. Segments twisted, pointed, barely overlapping. Corona deeper clear lemon yellow.
- YELLOW CHEERFULNESS** 4Y-Y
Perianth segments sulphur yellow, flat, rounded, overlapping, complete row. Corona complete, deeper yellow with pencil line of orange at edge, filled with corona and stamen segments, neat.
- YELLOW CLOUD** 1
Perianth sulphur yellow, creased. Segments flat, pointed, overlapping. Corona lemon yellow, open mouthed, badly proportioned.
- YELLOW DAZZLER** 1Y-Y
Perianth lemon, slightly reflexed. Segments acute, overlapping, some slightly notched. Corona lemon, flanged, gentle frill. Neck grows stiffly in water.
- YELLOW IDOL** 1Y-Y
Perianth light lemon yellow, creased, sometimes notched. Segments flat, pointed, well overlapping. Corona deeper lemon yellow, almost straight sided.
- YELLOW MOON** 2Y-Y
Perianth light lemon yellow, developing paler tips, smooth. Segments flat, pointed, broad, overlapping. Corona light lemon yellow, fluted, bowl shape. Neck grows stiffly in water.
- YELLOW POPPY** 2Y-YYO
Perianth clear primrose yellow, deeper at base. Segments flat, pointed, broad, overlapping. Corona greenish lemon yellow with thin orange line at edge.
- YELLOW SUN** 2Y-Y
Perianth lemon, slightly hooded. Segments narrow, acute, overlapping at base only. Corona lemon, straight sided, opening at mouth, deeply frilled. Neck grows badly in water.
- YELLOW SUNSET** 1Y-Y
Perianth yellow. Segments ovate, slightly twisted, with a slight overlap. Corona large frilled flared, and yellow as perianth.
- YELLOW TRIUMPHATOR** 1
Perianth yellow. Segments ovate, slightly twisted, with a slight overlap. Corona large frilled, flared and slightly deeper yellow than perianth. Neck grows in water.
- YEO** 2W
Perianth yellow. Segments ovate, overlapping and with a ridged texture. Corona medium sized frilled cup, pale orange with a deeper orange around rim. Neck grows badly in water.
- YORNUP** 2Y-Y
Perianth pale sulphur, slightly twisted, star shaped. Segments some notched, overlapping only at base. Corona gold, open mouthed, slightly frilled at edge. Neck grows badly in water.
- ZANITA** 7Y-GYY
Perianth greenish lemon yellow, edges curl in, crêpey texture. Segments flat, pointed, overlapping. Corona lemon yellow with slight green eye, deeply fluted, bowl shape. Neck grows badly in water.

ZARAH LEANDER 2Y-R
Perianth clear pale lemon yellow, creased badly. Segments flat, pointed, well overlapping. Corona orange red, white blotches on edge, double frilled.

ZEPHYR 2
Discarded - Short, flowers below foliage.

ZERO 2W-W
Perianth creamy white, slight green base, creased, pointed outline. Segments flat, pointed, overlapping. Corona creamy white, of trumpet proportions.

ZEST 1W-Y
Discarded - Untidy flower, flowers below foliage.

ZIRCON 2W-Y
Perianth lemony white. Segments broad and overlapping. Corona large frilled cup, yellow in colour. Neck grows badly in water.

ACKNOWLEDGEMENTS

In compiling this report describing 34 years of the narcissus variety trial at Rosewarne EHS, I should like to pay tribute to the directors of the Station and their staff. The directors were Mr F W Shepherd (1952-1962), Mr H J Eaton (1962-1977) and Mr M R Pollock (1977-1989). Special thanks are due to Miss B M Fry, Mr L W Wallis and Mr A A Tompsett, who had particular responsibilities for this work, and whose efforts have been brought together in this report.

At HRI Kirton, I would like to thank the following staff for their help in compiling this and earlier reports: Mr G P Clark, Mrs L J Withers, Mr R E Smith, Dr C D Paterson, Mrs C Kitchen and Mrs A Dawson.

Funding of the work by the Ministry of Agriculture, Fisheries and Food, until 1990, is hereby acknowledged.

BIBLIOGRAPHY

Earlier reports

ADAS, 1971. Narcissus variety trials. 3rd report 1968-71. Rosewarne Experimental Horticulture Station.

Fry, B M, 1965. Narcissus variety collection. Report Rosewarne EHS and Ellbridge Sub-station for 1964: 32-36.

Fry, B M and Shepherd, F W, 1961. Daffodil variety collection. Report Rosewarne EHS and Ellbridge Sub-station for 1960: 28-48.

Hanks, G R, 1992. Narcissus variety assessment (Fifth report). Report to Horticultural Development Council.

NAAS, 1964. Narcissus variety trials. 1st report 1955-63. Station Leaflet no. 3, Rosewarne Experimental Horticulture Station.

NAAS, 1967. Narcissus variety trials. 2nd report 1964-67. Rosewarne Experimental Horticulture Station.

Paterson, C D, 1991. Narcissus variety assessment (Fourth report). Report to Horticultural Development Council.

Checklists, etc

ADS, 1989. Daffodils to show and grow. Abridged classified list of daffodil names. 4th edition. American Daffodil Society : Hernando.

Kington, S, 1989a. The international daffodil checklist. Royal Horticultural Society: London.

Kington, S, 1989b. The international daffodil register (1969), 15th supplement. (1st supplement to 1989 checklist). Royal Horticultural Society: London.

Kington, S, 1990. The international daffodil register (1969), 16th supplement. (2nd supplement to 1989 checklist). Royal Horticultural Society: London.

Kington, S, 1991. The international daffodil register (1969), 17th supplement. (3rd supplement to 1989 checklist). Royal Horticultural Society: London.

Kington, S, 1992. The international daffodil register (1969), 18th supplement. (4th supplement to 1989 checklist). Royal Horticultural Society: London.

Kington, S, 1993. The international daffodil register (1969), 19th supplement. (5th supplement to 1989 checklist). Royal Horticultural Society: London.

Husbandry


ADAS, 1970. Narcissus in SW England. Station leaflet No. 4 (second edition). Rosewarne Experimental Horticulture Station.

ADAS, 1985. Narcissus bulb production. Booklet 2150 (Revised). MAFF Publications: Alnwick.

Rees, A R, Wallis, L W and Tompsett, A A, 1973. Effects of planting density, plant arrangement and frequency of lifting on flower and bulb production of narcissus in S W England. Journal of Horticultural Science, 48: 59-73.

GRH3.REP

FIGURE 1. PERIANTH AND SEGMENT SHAPES



COMMERCIAL IN CONFIDENCE

FIGURE 2. CORONA SHAPES AND EDGES AND LEAF POISE

