

Project title	Onions: Improving risk assessment for stem nematode
Project number:	FV 327
Project leader:	Dr Steve Ellis, ADAS
Report:	Final report, August 2009
Previous report	Annual report, August 2008
Key staff:	Dr Steve Ellis, ADAS UK Ltd Dr Bill Parker, ADAS UK Ltd David Norman, Fresh Produce Consultancy Ltd
Location of project:	Cambridgeshire & ADAS, High Mowthorpe, North Yorkshire
Project coordinator:	Mr R Brown, E C Brown, Wypemere Farm, Angel Bridge, Whittlesey, Cambs, PE7 2HQ
Date project commenced:	3 May 2007
Date project completed (or expected completion date):	31 August 2009
Key words:	Onions, stem nematode, <i>Ditylenchus dipsaci</i> , soil sampling, GPS mapping, soil extraction

Whilst reports issued under the auspices of the HDC are prepared from the best available information, neither the authors nor the HDC can accept any responsibility for inaccuracy or liability for loss, damage or injury from the application of any concept or procedure discussed.

The contents of this publication are strictly private to HDC members. No part of this publication may be presented, copied or reproduced in any form or by any means without prior written permission of the Horticultural Development Company.

The results and conclusions in this report are based on an investigation conducted over a one-year period. The conditions under which the experiments were carried out and the results have been reported in detail and with accuracy. However, because of the biological nature of the work it must be borne in mind that different circumstances and conditions could produce different results. Therefore, care must be taken with interpretation of the results, especially if they are used as the basis for commercial product recommendations.

AUTHENTICATION

We declare that this work was done under our supervision according to the procedures described herein and that the report represents a true and accurate record of the results obtained.

Dr S Ellis
Research Entomologist
ADAS

Signature  Date28.9.09.....

Mr D Norman
Director
Fresh Produce Consultancy Ltd

SignatureApproved by email..... Date28.9.09.....

Report authorised by:

Mr J Clarke
Horticulture Sector Manager
ADAS

SignatureApproved by email.....Date28.9.09.....

TABLE OF CONTENTS

GROWER SUMMARY	1
HEADLINES	1
BACKGROUND AND EXPECTED DELIVERABLES	1
SUMMARY OF THE PROJECT AND MAIN CONCLUSIONS	2
<i>Comparing nematode extraction methods</i>	2
<i>Mapping the in-field distribution of stem nematode</i>	2
VALIDATION SAMPLING	3
FINANCIAL BENEFITS	4
ACTION POINTS FOR GROWERS.....	5
SCIENCE SECTION	6
INTRODUCTION	6
MATERIALS AND METHODS.....	7
<i>Comparison of nematode extraction methods</i>	7
<i>Mapping stem nematode distribution within fields</i>	8
<i>Impact of reducing sample size on sampling accuracy</i>	8
<i>Validation sampling</i>	10
RESULTS.....	10
<i>Comparison of nematode extraction methods</i>	10
<i>Mapping stem nematode distribution within field</i>	12
<i>Impact of reducing sample size on sampling accuracy</i>	15
<i>Validation sampling</i>	20
DISCUSSION	25
CONCLUSIONS	27
TECHNOLOGY TRANSFER.....	28
GLOSSARY	28

Grower Summary

Headlines

- There are distinct patches and gaps in nematode distribution within fields. The best chance of detecting the pest is through systematic sampling across a grid, which must comprise 100 sample points for every 4 hectares.
- The Seinhorst two-flask technique is the most effective laboratory method of extracting stem nematodes.

Background and expected deliverables

Stem nematode (*Ditylenchus dipsaci*) is a very destructive pest of bulb onions. Assured Produce protocols strongly recommend the use of representative soil sampling as part of risk assessment for the pest. This ignores the inherent problems of detecting it in soil. The pest is primarily an endoparasite (spending most of its life in the plant) so numbers in soil are usually small. This project aimed to develop optimum methods of soil sampling and analysis for detecting the pest at typical population densities encountered in commercial practice. This involved the use of soil sampling linked to differential GPS (DGPS) sample location. In particular, the project investigated within-field distribution of the pest and developed sampling protocols to give the maximum chance of detection. As not all soil extraction methods are equally effective at recovering stem nematode, different techniques were compared, to determine best practice.

The expected deliverables from this work included:

- Clear guidelines on which extraction method to request from analytical laboratories, to give the best chance of recovering stem nematode from soil.
- Graphical representation of the distribution of stem nematodes within fields at high, medium and low risk from the pest to aid generation of sampling plans.
- Sampling protocols that optimise the chance of detecting stem nematode at typical levels of infestation
- Guidelines to give improved confidence in risk assessment for stem nematode.

Summary of the project and main conclusions

Comparing nematode extraction methods

Soil samples from fields known to be heavily infested with stem nematode were mixed thoroughly and a sub-sample extracted using the Whitehead tray, Seinhorst two-flask, Flegg modified Cobb and centrifugation techniques. A total of five replicate samples were extracted using each technique. There were clear differences between the methods in terms of their efficiency of extraction of stem nematode. Overall the Seinhorst two-flask technique was most effective and on average extracted almost three times as many stem nematodes as the Flegg modified Cobb technique and just over twice as many nematodes as the Whitehead tray and centrifugation techniques (Figure 1). Interestingly, the Seinhorst two-flask technique was also most effective at extracting all other nematode groups recovered from the samples.




Figure 1. Numbers of stem nematodes (\pm standard errors) recovered from soil using different extraction methods

Mapping the in-field distribution of stem nematode

This part of the project took advantage of automated soil sampling equipment linked to differential GPS (DGPS) provided by Fresh Produce Consultancy. Intensive soil sampling was concentrated on three fields known to pose a high, medium or low risk to crops due to stem nematode. GPS-generated sampling grids were prepared for a four hectare block in each field to provide 400 10 x 10m sampling plots. A total of five soil cores were taken from within each plot to provide a bulked sample of about 50 g of soil. Coordinates for the

location of each sample were obtained using DGPS. Individual soil samples were extracted using the Seinhorst two-flask technique and the number of stem nematodes counted.

The nematode data was linked to sampling coordinates and used to map the characteristics of the pest's distribution within each field. As expected, the distribution of nematodes varied depending on whether crops grown in the field were perceived to be at low, medium or high risk from stem nematode. Detailed analysis of the distributions suggests that there are distinct patches and gaps in the nematode distribution across a field rather than it being just random. This will have to be taken into account when developing sampling protocols. Using the distribution maps as a tool, some preliminary work has been done to determine how reducing the number of sampling points affects the mean nematode count. This helped to indicate how few sampling points were needed to produce an accurate assessment of pest numbers and risk of crop damage. For example, by systematically omitting every eighth, fourth or second count it was possible to reduce the number of sampling points to 350, 300 and 200 respectively. Results to date suggest that this has little effect on the apparent mean nematode population in the field. As the pest shows a patchy distribution, a systematic sampling plan is probably the best option as it covers the sample area more uniformly and increases the chances of locating a patch of stem nematode. If you simply sampled randomly, the chances of missing the patch could be higher.

Validation sampling

Following on from mapping the in-field distribution of stem nematode it was possible to suggest a sampling protocol which is likely to give the optimum chance of detecting the pest. Results suggested that a systematic sampling plan is probably the best option as it covers the sampling area uniformly and so increases the chances of locating a patch of stem nematode.

A total of 11 commercial onion fields known to be infested with stem nematode were chosen to test the sampling protocol. In each field a 100-core sample was taken with a pneumatic sampler, single cores being taken from each square of a GPS-generated sample grid containing 100 squares each measuring 20 x 20 m. This was compared with a 25-core sample taken by hand with a cheese corer via a "W" shaped path. This being typical of samples submitted to analytical laboratories for extraction to detect stem nematode. Sufficient soil was taken so that both a Seinhorst two-flask and Whitehead tray extraction could be performed on each sample.

Results showed that in general more stem nematodes were recovered from the pneumatically-sampled soil compared with the hand-sampled soil, irrespective of extraction method. This was most marked for the Seinhorst two flask extractions where about 50% more nematodes were found in the pneumatic compared with the hand samples. The validation samples also confirmed the superiority of the Seinhorst two-flask extraction over the Whitehead tray for extracting stem nematode. For hand samples, 89% more stem nematodes were recovered using the Seinhorst two-flask extraction compared with the Whitehead tray. For pneumatic samples, 86% more nematodes were recovered using the Seinhorst two-flask extraction compared with the Whitehead tray. Seinhorst two-flask extraction of hand- or pneumatic-sampled soil doubled the chances of detecting stem nematode when compared to the Whitehead tray.

Hand sampling via a W-shaped path recovered stem nematode at two more sites than did pneumatic sampling from a systematic sampling grid, although numbers at both these sites were very low. This result is surprising as examination of the in-field distribution of stem nematode suggested that a systematic sampling grid was probably the best option for detecting stem nematode. A number of factors may have contributed to this result. It is possible that nematodes were uniformly distributed in most fields so the W-shaped path was as effective as the sampling grid. Also, it is possible that the random sample happened to hit a patch of nematodes. Finally the volume of soil taken with the hand sampler may have increased the chances of finding nematodes. Despite this result and in view of the potential patchy distribution of stem nematode it is likely that a systematic sampling grid of 100 points covering 4ha will give the best chance of detecting stem nematode.

Financial benefits

Estimates from historical records of soil sampling suggest that about 2.5% of land sampled for stem nematode has a detectable infestation. Therefore, for every 1,000 ha of onions grown on sampled land there may be 25 ha lost due to stem nematode if no treatment was applied. With a total UK crop area of onions of about 8,000 ha the total annual area lost, if untreated, would be about 200 ha with a value of around £1,000,000. Greater confidence in the results of soil sampling to predict stem nematode risk would significantly reduce this loss. Improved risk assessment would also help to minimise unnecessary use of nematicides as insurance treatments and allow growers to move towards nematicide-free production as requested by many supermarkets

Action points for growers

- Request that labs offering analysis of soil samples for detection of stem nematode use the Seinhorst two-flask extraction technique to give the best chance of recovering the pest.
- When sampling fields for stem nematodes, individual soil samples (which can be bulked) should be taken from at least 100 uniformly distributed points per 4 ha to give the best chance of detecting the pest and to ensure that an acceptable measure of the average numbers of nematodes present is obtained.

Science Section

Introduction

Stem nematode (*Ditylenchus dipsaci* (Kuhn) Filipjev) is a destructive pest of many plants. Onions may be attacked at any time after germination and may be killed outright before they reach the seedling stage even by apparently low levels of nematodes in soil. Following the withdrawal from use of Temik 10G (aldicarb) on 31 December 2007 only Vydate (oxamyl) remains available for stem nematode control on onions in the UK, with a specific off label approval (SOLA 2006/1890). As nematode numbers in soil are usually small and show an aggregated distribution, work was needed to define best practice for soil sampling to provide the maximum chance of detecting the pest. This project addressed this need by investigating in-field distribution of stem nematode, optimum sampling patterns for pest detection and comparing soil extraction methods to determine the most effective at recovering nematodes.

The risk of stem nematode damage is currently assessed by considering field history, previous cropping and by representative soil sampling. Soil sampling is recommended by Assured Produce protocols, but in practice current soil sampling strategies ignore the inherent problems of detecting low numbers of stem nematode in soil. With the loss of Temik 10G and pressure from retailers to reduce nematicide use, reliable soil sampling will become increasingly important for identifying fields at risk. Alternative detection strategies, such as sampling weed species for stem nematode, are not likely to be effective as weed distribution across fields is never uniform.

There is little information available on soil sampling for stem nematode in both the UK and Europe. Some workers have commented on the number of samples and depth of sampling that should be used. Caubel (1975) suggests that 50 x 20g samples to a depth of 15 to 20 cm are required where the nematode is concentrated. However, there is no mention of the maximum area to be sampled and the technique may not be effective in fields where there is little information on the existing nematode population, for example fields potentially available for rent. Areas of one third of a hectare were sampled 60 times to a depth of 20 cm by Kleijburg (1960) to provide 1 kg of soil for extraction. Currently in the UK, fields are hand-sampled by walking a W-shaped path and taking 25 cores or, when using mechanised sampling, up to 100 cores may be taken from a 4 ha area (David Norman, *pers comm*).

There is clearly no agreement over the number of cores to be taken from a particular area and this is something that the project addresses.

Muller *et al.* (1993) divided each of 20 replicate 2 kg soil samples into 16 x 125 g sub-samples. This provided 320 samples in which numbers of stem nematodes were assessed. These data were used to calculate the nematode numbers in a range of sample sizes and predict the number of samples required to detect the pest. It was concluded that two 125 g sub-samples needed to be examined to detect stem nematode. It is not clear whether the field was considered at high risk of damage but the principle of intensively sampling onion fields and using these data to predict the accuracy of different sampling patterns to detect the pest was adopted in this study.

Automated soil sampling equipment linked to differential GPS (DGPS) sample location offers the opportunity to take large numbers of soil cores at known locations. This provides data on stem nematode distribution and allows better guidelines for the correct approach to soil sampling for the pest to be developed. This in turn should increase the chances of detecting low levels of stem nematode.

Experience has shown that not all laboratory extraction procedures are equally effective at recovering stem nematode from soil. The most efficient extraction procedures need to be used to ensure that extraction of low numbers of the pest can be reliably achieved, but there is no published literature comparing extraction methods for stem nematode. Therefore this study compared the efficiency of the Whitehead tray, the Flegg modified Cobb technique, centrifugation and the Seinhorst two-flask technique in recovering stem nematodes from soil.

Overall, improved soil sampling, pest extraction and results interpretation will increase growers' confidence in making valid risk assessments, leading to reduced/targeted use of nematicides and allowing cost savings for growers.

Materials and methods

Comparison of nematode extraction methods

A total of four mineral soil samples, each weighing approximately 10 kg, were taken in September 2007 from a field in Cambridgeshire known to be heavily infested with stem nematode. The samples were bulked together and thoroughly mixed. This sample was then sub-sampled to create 20 samples each of approximately 1 kg. Five replicate samples from

each subsample were then extracted to determine the level of stem nematode infestation, using each of the following extraction methods.

1. Whitehead Tray (Whitehead and Hemming, 1965)
2. Flegg modified Cobb technique (Flegg, 1967)
3. Seinhorst two-flask technique (Seinhorst, 1955)
4. Centrifugation (Jenkins, 1964)

Nematode suspensions were cleaned to aid microscopic examination. This was done using a small sieve consisting of a plastic ring (8 cm diameter) cut from vinyl drain-pipe to which a circle of plastic mesh (48 μ mesh) was glued at one end. This sieve was placed inside a Petri dish and the nematode suspension plus debris poured onto it. More water was added carefully down the inside of the Petri dish until the debris in the small sieve was immersed. The Petri dish was then covered and allowed to stand undisturbed for 48 hours. During this period the nematodes wriggled out of the debris, through the mesh and into the Petri dish where they could be collected. The numbers of nematodes recovered were counted and expressed as numbers/litre of soil. These data were subjected to the analysis of variance. The time taken to undertake each stage of the extraction was also noted.

Mapping stem nematode distribution within fields

Intensive soil sampling was done in a 4 ha block in each of three fields in Cambridgeshire, known to be at high (Block A, Denton Lodge Farm, Holme, Peterborough, Cambs. Grid ref TL 183884), medium (Block B, Carolls Farm, Manea, Cambs. Grid ref TL 494904) and low (Block C, Carolls Farm, Manea, Cambs. Grid ref TL 495902) risk respectively of having significant stem nematode populations. Soil samples were taken using automated soil sampling equipment linked to differential GPS (DGPS). GPS-generated sampling grids were prepared for a four hectare block in each field to provide 400 10 x 10 m sampling plots. A total of five soil cores were taken from within each plot to provide a bulked sample of about 50 g of soil. Co-ordinates for the location of each sample were taken using the DGPS. Individual soil samples were extracted using the Seinhorst two-flask technique and the number of stem nematodes counted. Numbers were converted to number/litre soil. The nematode data was linked to sampling co-ordinates and used to map the characteristics of the pest's distribution within each field.

Impact of reducing sample size on sampling accuracy

Analyses were done to assess the impact of reducing sample size on the accuracy of both the overall mean infestation level identified by the sampling, and the apparent spatial

distribution of nematodes in the sample blocks. To generate appropriate reduced data sets, two approaches were taken:

- a) *Systematic reduction*: all sample points from each block were ranked in the order in which they were taken during the sampling process. Data sets reduced by 50, 100, 200, 300, 350 and 375 points from the original 400 in each block were obtained by removing the 8th data point in every series of 8 (50 points removed); the 4th data point in every series of 4 (100 points removed); and every other data point (200 data points removed). Data sets reduced by 300, 350 and 375 points were obtained by retaining only every 6th, 10th and 18th data point respectively.
- b) *Random reduction*: each sample point in each block was assigned a random number (using the random number function in Microsoft Excel). To obtain a sample reduced by 50 points, all sample points were ranked in ascending order by the random number and the first 50 points removed. The process was repeated to remove the number of data points required to generate data sets reduced by 100, 200, 300, 350 and 375 points for each block, using a new randomisation for each selection.

Mean and 95% confidence limits were calculated for each sample size for each block.

To assess the impact on the apparent spatial distribution of the nematodes in each block of reducing sample size, the systematically-reduced data sets (nematode counts and their associated x, y coordinates within the sampling block) were used to calculate an index of aggregation (I_a) using SADIE (Spatial Analysis by DistancE Indices (Perry, 1995)). Index values significantly greater than 1 indicate that counts exhibit a degree of aggregation, and an associated probability value indicates whether the degree of aggregation is significantly different from what would be obtained from random permutations of the counts. In essence therefore, I_a indicates whether or not the counts are randomly distributed.

In addition, SADIE 'red-blue' plots (Perry *et al.*, 1999) were generated for the full data sets and for data sets systematically reduced to 100 points. In simple terms, this analysis calculates an index of clustering for each data point. The analysis assesses whether a high count at a particular location is associated with other high counts at neighbouring data points = a 'patch', and similarly whether low/zero counts are associated with low values at neighbouring points = a 'gap'. These values can then be interpolated to produce a contour map with patches (red) where the cluster index is >1.5 and gaps (blue) where the index is <1.5 .

These analyses could only be done on data from blocks A and B as the data set from block C contained too many zeroes to give reliable results.

Validation sampling

Detection of stem nematode is considered to be more important than an accurate estimate of numbers of the pest as the advice is not to grow onions in soil infested with stem nematode (even populations below 25 nematodes/litre of soil have been known to completely destroy a drilled onion crop). Work on mapping the in-field distribution of stem nematode suggested that when taking a soil sample to detect the pest a systematic sampling plan was probably the best option when compared to a random sampling plan. The systematic plan covers the sample area uniformly and also increases the chances of locating a patch of stem nematode.

Validation samples were taken in spring 2009. A sample was taken using pneumatic, automated sampling apparatus linked to differential GPS. This sample comprised 100 cores taken in a systematic manner from a GPS-generated rectangular sampling grid consisting of 100, 20 m x 20 m sample areas. The total area sampled was 4 ha. The size of the sampling area and number of cores was determined from the results of the analyses to investigate the impact of reducing sample size on sample accuracy. The automated sample was compared with one which was taken by hand-sampling using a 20 cm deep x 2 cm diameter cheese corer. The sample comprised 25 cores taken over a "W" shaped sampling path. This is the type of sample most commonly submitted to analytical labs when attempting to detect stem nematode. The hand sampling produced a sample of 5 kg in comparison with 0.8 kg from the automated sampler. A total of 11 fields were sampled for stem nematode. These were all fields in which a moderate or light infestation was expected, with the exception of one field (Bliss 3) where a heavy infestation of the pest had been recorded. All samples were extracted using the Seinhorst two-flask technique and the Whitehead tray to provide additional data with which to compare the extraction methods.

Results

Comparison of nematode extraction methods

Time to perform nematode extractions

The total time taken to perform each of the nematode extraction methods is shown in Table 1. Centrifugation was the least time-consuming extraction method. This is mainly because the other methods require samples to be left for 24-48 hours as part of the

extraction or to clear the nematode suspension. During this waiting time no staff input is required so other tasks can be undertaken.

Table 1. Time (minutes) taken to extract one soil sample for stem nematodes using the Whitehead tray, Flegg modified Cobb technique, Seinhorst two flask technique and centrifugation

Extraction method	Time (minutes) for different stages of extraction			Total
	Soil preparation	Extraction	Clearing suspension	
Whitehead tray	5	1440	3	1448
Flegg modified Cobb technique	5	6	2880	2891
Seinhorst two-flask technique	5	15	2880	2900
Centrifugation	5	36	0	41

Table 2. Mean numbers of a range of nematodes recovered from replicate soil samples using four extraction methods.

Extraction method	Nematode group (number/l soil)			
	Stem nematode	Stunt/spiral nematodes	Root lesion nematodes	Cyst juveniles
Whitehead tray	59	225	205	80
Flegg modified Cobb technique	46	95	0	45
Seinhorst two-flask technique	128	920	250	110
Centrifugation	54	45	5	10
SED (16 DF)	9.7	37.6	37.4	20.2

Extraction efficiency

A total of four nematode groups were recovered in sufficient numbers using each extraction method to subject to statistical analysis. These were stunt/spiral nematodes (*Tylenchorynchus* spp), cyst juveniles (larvae of *Globodera* spp or *Heterodera* spp), root lesion nematodes (*Pratylenchus* spp) and stem nematode (*Ditylenchus dipsaci*). Numbers of all nematodes recovered differed significantly between extraction methods. ($P < 0.001$ for each nematode group, Table 2.) The Seinhorst two-flask technique was the most effective extraction method for all four nematode groups studied. It was significantly better ($P < 0.05$)

than all other extraction methods for stem nematode and stunt/spiral nematodes and statistically better ($P<0.05$) than centrifugation and the Flegg modified Cobb technique for cyst juveniles and root lesion nematodes. The Whitehead tray was significantly better ($P<0.05$) than centrifugation and the Flegg modified Cobb technique for stunt/spiral nematodes and root lesion nematodes and significantly better than centrifugation for cyst juveniles. Overall centrifugation was probably the least effective extraction method.

Mapping stem nematode distribution within field

Numbers of stem nematode recovered from fields at high medium and low risk from stem nematode were linked to sampling co-ordinates and used to map the characteristics of the pest's distribution within each of the high, medium and low risk field areas (Figures 1 and 2) using punctual kriging as the interpolation routine. These maps tend to suggest the presence of a patchy distribution, particularly for stem nematode in the high risk (Block A) field (Figure 1). Medium infestations (Block B) suggested are more random distribution (Figure 2), or at least a distribution with fewer and less well-defined patches. In the low risk block (Block C), distinct but small patches were evident.




Figure 1. Punctual kriging contour map (courtesy of Fresh Produce Consultancy) to show distribution of stem nematode in a 4ha block known to be at high (Block A) risk from stem nematode



Figure 2. Punctual Kriging contour map (courtesy of Fresh Produce Consultancy) to show distribution of stem nematode in a 4ha block known to be at medium (Block B) and low (Block C) risk from stem nematode

Impact of reducing sample size on sampling accuracy

The effect of reducing data sets for each block by systematic and random reduction are shown in Figures 3, 4 and 5 for blocks A, B and C respectively. The overall mean populations levels for the three blocks for the full data sets (400 points) were block A: 95.8 nematodes/litre of soil; block B: 16.3 nematodes/litre of soil; and block C: 0.14 nematodes/litre of soil (approximately one order of magnitude between each block). Reducing sample sizes from 400 to any value down to 100 had very little effect of the overall mean in any of the blocks, whether sample points were randomly or systematically removed. Sample sizes of <100 tended to have higher errors associated with them. Plots of log mean against log variance for each data set for each block (data not shown) generally showed that outlying points were associated with sample sizes of <100, suggesting that mean/variance relationships were more stable for sample sizes >100. Even where infestation levels were very low (Block C) nematodes were found at all sample sizes except 25 samples (Figure 3).




Figure 3. The effect on the mean nematode infestation level (\pm 95% confidence limits) in Block A (high risk) of reducing the sample point number randomly and systematically




Figure 4. The effect on the mean nematode infestation level (\pm 95% confidence limits) in Block B (medium risk) of reducing the sample point number randomly and systematically




Figure 5. The effect on the mean nematode infestation level (\pm 95% confidence limits) in Block C (low risk) of reducing the sample point number randomly and systematically

Spatial analyses

The values of I_a obtained from the full and systematically reduced data sets from Blocks A and B are given in Table 3.


Table 3. Values of I_a and associated probabilities (P) for full and systematically reduced data sets in Blocks A and B. Values of P in bold are significant

Sample size	Block A		Block B	
	I_a	P	I_a	P
50	1.012	0.396	1.006	0.400
100	1.139	0.208	1.198	0.118
200	1.184	0.148	1.741	0.005
300	1.271	0.094	2.021	0.005
350	1.427	0.031	2.095	0.005
400	1.486	0.039	2.220	0.013

In both blocks, values of I_a declined as sample size decreased, indicating that the observed nematode distribution tended to become more random as the sample size decreased. For block A (high infestation), values of I_a were significant at sample sizes of 400 and 350, but not at smaller sample sizes, suggesting that at sample sizes <350, the nematode distribution was random rather than patchy. For block B, evidence of aggregation in the counts was observed down to a sample size of 200.

The SADIE 'red-blue' plots (Figures 6 and 7) show that significant patches and gaps can be identified in the distribution of nematode in both block A and block B when the full data sets (400 data points are used). When the same analysis is done using at much reduced data sets (100 points), patches and gaps are still present to a lesser extent, but do not correlate particularly well with those indicated on the maps for the full data sets.

a) 400 sample points



b) 100 sample points





Figure 6. SADIE 'red-blue' plots for block A for full (400 data points) and reduced (100 data points) data sets. Crosses indicate the location of the sample points (data are interpolated beyond the physical limits of the sampling block). Red areas = patches, blue areas = gaps

a) 400 sample points



b) 100 sample points




Figure 7. SADIE 'red-blue' plots for block B for full (400 data points) and reduced (100 data points) data sets. Crosses indicate the location of the sample points (data are interpolated beyond the physical limits of the sampling block). Red areas = patches, blue areas = gaps

Validation sampling

In view of the difference in size between the automated pneumatic sample (800 g) and the hand-sample (1.5 kg), the number of nematodes extracted from each sample was divided by the weight of the sample in kilogrammes to give the number per kg in each case. This ensured that any analyses of the datasets from both sampling methods compared like with like. The numbers of nematodes per litre of soil corrected for sample weight are given in Table 4.

Paired “T” tests were used to determine whether there was any difference in numbers of different nematode groups between hand and pneumatic samples both within and between extraction methods. The complete list of comparisons is given below:

- a. Hand sampling and Seinhorst two-flask extraction v’s pneumatic sampling and Seinhorst two-flask extraction.
- b. Hand sampling and Whitehead tray extraction v’s pneumatic sampling and Whitehead tray extraction.
- c. Hand sampling and Seinhorst two-flask extraction v’s hand sampling and Whitehead tray extraction.
- d. Pneumatic sampling and Seinhorst two-flask extraction v’s pneumatic sampling and Whitehead tray extraction.

The paired “T” test used the mean difference between counts for the two sampling and extraction methods to test the null hypothesis that the mean difference in nematode counts for a particular comparison was equal to zero. Paired “T” tests were done on the data for stunt/spiral nematodes, root lesion nematodes and stem nematode. Numbers of all other species recovered were too low for analysis.

The mean numbers of nematodes recovered from the two sampling methods and two extraction methods are given in Table 5.

Table 4. Numbers of nematodes/l soil extracted by the Seinhorst two-flask technique or Whitehead tray from hand or pneumatic samples corrected for sample weight

Field	Stubby root nematodes	Stunt/ Spiral nematodes	Cyst juveniles	Root lesion nematodes	Needle nematodes	Stem nematode
Samples extracted using Seinhorst two-flask technique						
Hand samples						
Heading C9 B	0	0	0	83	0	
Heading C9 C	0	17	17	133	0	
D Norman Bell Big 27	0	300	50	183	0	
D Norman Old Pear 5	0	267	0	233	3	
D Norman TF40A	0	117	0	33	0	
Chans Wheatsheaf 1	17	17	0	200	0	
Jenson Turves 1	0	33	0	117	0	
Kisby 1	0	133	0	117	0	
Barshall E 1	317	267	150	100	0	
Bliss 3 1	0	350	0	150	0	
Wype Butler 1	0	100	0	233	0	
Total	333	1600	217	1583	3	3833
Pneumatic samples						
Heading C9 B	0	0	0	125	0	38
Heading C9 C	0	31	31	188	0	13
D Norman Bell Big 27	0	281	31	250	0	0
D Norman Old Pear 5	31	156	0	63	0	31

Table 4 (cont'd)

D Norman TF40A	0	188	63	125	0	25	
Chans Wheatsheaf 1	0	0	0	63	0	0	
Jenson Turves 1	0	31	0	125	0	0	
Kisby 1	0	125	0	94	0	19	
Barshall E 1	375	125	125	188	0	6	
Bliss 3 1	0	594	0	125	0	13	
Wype Butler 1	0	63	0	188	13	0	
Total	406	1594	250	1531	13	144	3938

Samples extracted using Whitehead tray

Hand samples

Heading C9 B	0	17	0	17	0	3	
Heading C9 C	0	0	17	50	0	0	
D Norman Bell Big 27	0	300	17	117	0	0	
D Norman Old Pear 5	0	83	0	17	0	7	
D Norman TF40A	0	17	0	33	0	10	
Chans Wheatsheaf 1	0	17	0	83	0	0	
Jenson Turves 1	0	17	0	183	0	0	
Kisby 1	0	17	17	17	0	10	
Barshall E 1	67	0	33	17	0	0	
Bliss 3 1	0	900	0	350	0	0	
Wype Butler 1	0	67	0	250	0	3	
Total	67	1433	83	1133	0	33	2750

Table 4 (cont'd)

Pneumatic samples							
Heading C9 B	0	0	31	63	0	0	
Heading C9 C	0	0	94	94	0	0	
D Norman Bell Big 27	0	156	63	188	0	0	
D Norman Old Pear 5	0	125	0	63	0	6	
D Norman TF40A	0	31	63	31	0	19	
Chans Wheatsheaf 1	0	31	0	313	0	0	
Jenson Turves 1	0	31	0	500	0	0	
Kisby 1	0	31	31	31	0	6	
Barshall E 1	63	0	63	63	0	0	
Bliss 3 1	0	531	0	313	0	13	
Wype Butler 1	0	188	0	563	0	6	
Total	63	1125	344	2219	0	50	3800

Table 5. Mean numbers of nematodes (number/l soil) recovered by hand and pneumatic sampling and by Seinhorst two-flask or Whitehead tray extraction

Nematode species	Mean number nematodes/l soil			
	Seinhorst two flask		Whitehead tray	
	Hand sample	Pneumatic sample	Hand sample	Pneumatic sample
Stunt/spiral nematodes	145.5	144.9	130.3	102.3
Root lesion nematodes	143.9	139.2	103.0	201.7
Stem nematode	10.7	16.0	3.7	5.6

There was no significant difference in numbers of stunt/spiral nematodes per litre of soil between hand or pneumatic samples irrespective of whether they were extracted by the Seinhorst two-flask technique or by the Whitehead tray. However, in general more nematodes were found in the hand sample compared with the pneumatic sample. Also numbers of stunt/spiral nematodes extracted by the Seinhorst two-flask technique and the Whitehead tray did not differ significantly for hand samples although there was a trend to find more nematodes in the two-flask extracted soil. Although the number of stunt/spiral nematodes extracted using the two-flask technique or Whitehead tray did not differ statistically there were on average more nematodes where the two-flask technique was used.

Numbers of root lesion nematodes did not differ significantly between hand and pneumatic sampling when extracted using the Seinhorst two-flask technique. However, significantly more root lesion nematodes ($P < 0.05$, $T = -2.60$, 10 df) were recovered from pneumatically sampled soil compared with hand sampled soil using the Whitehead tray extraction. Almost twice as many nematodes were found in the pneumatic than in the hand sample. There was no significant difference in numbers of root lesion nematodes extracted using the Seinhorst two-flask technique compared with the Whitehead tray for both hand and pneumatic samples. However, in general the two-flask technique tended to recover most nematodes from hand sampled soil whereas the Whitehead tray recovered most nematodes from the pneumatically sampled soil.

There was a trend to recover more stem nematodes from the pneumatically-sampled soil compared with hand-sampled soil irrespective of extraction method. This was most marked for the two-flask extraction where about 50% more nematodes were found in the pneumatic samples compared with the hand samples ($T = -1.99$, $P = 0.081$, 8 df). The two-flask

extraction recovered more stem nematodes than the Whitehead tray for both hand and pneumatic soil samples ($T = 2.51$, $P < 0.05$, 8 df for hand samples, $T = 2.29$, $P = 0.051$, 8 df for pneumatic samples). For hand samples 89% more nematodes were recovered using the two-flask compared with the Whitehead tray. For pneumatic samples 86% more nematodes were recovered using the two-flask compared with the Whitehead tray.

The Seinhorst two-flask technique improved the chances of detecting stem nematode approximately two-fold when compared with the Whitehead tray. Two-flask extraction recovered stem nematode from nine sites for hand sampled soil and seven sites for pneumatically sampled soil. The Whitehead tray recovered stem nematode from only four sites for both hand and pneumatically sampled soil. There were no occasions where the Whitehead tray recovered stem nematode but the two-flask did not.

In two fields, Chans Wheatsheaf 1 and Jenson Turves 1 no stem nematode were detected by any extraction or sampling method. The pest had been detected in both of these fields previously but at low levels (5 stem nematodes per litre soil). However, before the validation samples were taken the fields were ploughed so it is possible that nematodes were either buried below sampling depth or distributed over a larger area of the field.

Discussion

Stem nematodes were recovered from all replicate samples used to compare the efficacy of extraction methods, confirming that the crops grown in the sampled field were at high risk from the pest. Numbers recovered using all four extraction methods were sufficiently high to suggest that the field should not be cropped with onions.

The Seinhorst two-flask technique was the most effective extraction method for all nematode groups recovered from the samples. This was most obvious with stem nematode and stunt/spiral nematodes. The Seinhorst two-flask technique recovered at least twice as many stem nematodes and at least four times as many stunt/spiral nematodes as any other extraction methods. The Whitehead tray was generally the second most effective extraction method.

The superiority of the Seinhorst two-flask extraction method over the Whitehead tray was confirmed by the validation sampling, particularly for stunt/spiral nematodes and stem nematode. The difference between the two extraction methods was statistically significant

for stem nematode with 89% and 86% more nematodes recovered using the two-flask compared with the Whitehead tray for hand and pneumatically sampled soil respectively. Two-flask extraction also doubled the chances of detecting stem nematode when compared with the Whitehead tray. The differences were not as marked for stunt/spiral nematodes but in general most were recovered following two-flask extraction.

The results for root lesion nematodes were inconsistent and difficult to explain. There was little difference between the hand and pneumatic samples when extracted using the Seinhorst two-flask technique, whereas almost twice as many nematodes were recovered from the pneumatic sample as from the hand sample for Whitehead tray extraction.

The Flegg modified Cobb technique and centrifugation were least effective of all the extraction methods compared. The Flegg modified Cobb technique is principally intended for extraction of larger nematode species such as needle (*Longidorus* spp) and dagger nematodes (*Xiphinema* spp). Therefore, it is not surprising that it was ineffective for lesion nematodes, cyst juveniles and stem nematode which are all relatively small species.

Although centrifugation was the least time consuming extraction method, the extracted nematodes were less easy to examine than those from any of the other methods tested. This is because the nematodes became distorted following extraction in sucrose solution and so were less easy to identify than those extracted in water. The effect of the sucrose solution was more marked with stunt/spiral nematodes, root lesion nematodes and cyst juveniles than with stem nematodes. A further disadvantage of centrifugation is that some organic debris remained in the sample which made microscopic examination more difficult than with the other extraction methods. Although centrifugation has the advantage over other extraction methods in that it will extract dead nematodes, it was the least user-friendly method of all those tested.

The analysis of the sampling data clearly indicated that if all that was required from the sampling data was to be able to detect stem nematode and provide a reasonable estimate of the mean nematode infestation present, then the number of sampling points per 4 ha could be systematically reduced from 400 to 100 without significantly altering the mean. However, the SADIE analysis (Table 3) clearly shows that as sampling intensity is reduced, the nematode aggregation (patchiness) evident when 400 sampling points are used is lost (i.e. the distribution appears more random).

There was little to choose between the hand sampling regime and the systematic pneumatic sampling protocol. Indeed, hand sampling recovered stem nematode from two sites where pneumatic sampling system did not. However, at both these sites the level of nematode infestation was very low. This result is surprising as examination of the in-field distribution of stem nematode suggested that a systematic sampling grid was probably the best option for detecting stem nematode. A number of factors may have contributed to this result. It is possible that nematodes were uniformly distributed in most fields so the W-shaped path was as effective as the sampling grid. There was insufficient resource in this project to undertake intensive sampling of the validation fields so it is unclear how nematodes were distributed. It is also possible that the random sample happened to hit a patch of nematodes. Finally the volume of soil taken with the hand sampler may have increased the chances of finding nematodes. Despite there being little difference between hand sampling and the systematic sampling protocol and in view of the potential patchy distribution of stem nematodes in soil it is likely that a systematic sampling grid of 100 points covering 4 ha will give the best chance of detecting the pest.

Further work would be useful to help clarify the relative benefits of a systematic sampling grid and a random sampling pattern. This would concentrate on fields with moderate to low infestations of stem nematode where the aim of soil sampling would be to detect the pest

Conclusions

- The Seinhorst two-flask technique was the most effective extraction method for stem nematode and most other nematode groups.
- Growers should demand that analytical laboratories use the Seinhorst two-flask extraction to optimise chances of recovering stem nematode from soil samples.
- Centrifugation was the least user-friendly extraction method of those tested.
- It is likely that a systematic sampling grid of 100 points covering a 4 ha area will be the best compromise between sampling accuracy and economy of sampling effort. A standard 25 core sample along a W pattern across the same area may not be sufficient to identify low populations particularly if the nematodes have a patchy distribution. Further work would be necessary to clarify the relative benefits of a systematic sampling grid and a random sampling pattern.

- Pneumatic sampling tended to result in higher nematode counts than hand sampling although the differences were not always statistically significant.

Technology transfer

The project has featured in all of the following

HDC News May 2008, Plant, feed, spray and weed with accuracy

HDC News July 2008, How to find a needle in a haystack

Precision Farming Event, Springfields Spalding, March 26 2008

Article in Vegetable Farmer, October 2007

BOPA R&D group meeting at PG Rix, Stourgarden, 29 January 2008

BCGA Carrot meeting PGRO 21 February 2008

Elsoms Onion Conference, November 2008

HDC News, October 2009

Glossary

Cluster index – within SADIE (see below), a method for describing a region of either relatively large counts close to one another in two-dimensional space (i.e. a patch), or of relatively small counts (i.e. a gap).

Differential GPS – a means of accurately locating sampling points using the Global Positioning System satellite network linked to known reference points on the ground to obtain very high (sub-metre) accuracy.

Endoparasite – a parasite which spends most of its time within its host

Index of aggregation (I_a) – the degree to which a set of counts is identified as being aggregated spatially as defined using SADIE (see below).

Punctual krigging – a statistical method for interpolating point data sets into a contour map format.

SADIE – Spatial Analysis by Distance IndicEs. A statistical technique for assessing the degree of aggregation in a spatially-referenced set of counts.

References

Caubel, G. (1975). Estimation of *Ditylenchus dipsaci* populations in soil. Technical note. Sciences Agronomiques Rennes. Rennes, E.N.S.A., France: 1975. 97-99.

- Flegg, J J M. (1967). Extraction of *Xiphinema* and *Longidorus* species from soil by a modification of Cobb's decanting and sieving technique. *Annals of Applied Biology* **60**: 429-437.
- Jenkin, W R. (1964). A rapid centrifugal-flotation technique for separating nematodes from soil. *Plant Disease Reporter* **48**: 692.
- Muller, J, Knuth, P, Sturhan D. (1993). Requirements for the assessment of soil population densities of the stem nematode *Ditylenchus dipsaci*. *Nachrichtenblatt des Deutschen Pflanzenschutzdienstes* **45**: 73-77.
- Perry, J N. (1995). Spatial analysis by distance indices. *Journal of Animal Ecology* **64**: 303-314.
- Perry, J N, Winder, L, Holland, J M, Alston, R D. (1999). Red-blue plots for detecting clusters in count data. *Ecology Letters* **2**: 106-113.
- Seinhorst, J W. (1955). Een eenvoudige methode voor het afschieden van aaltjes uit grond. *Tijdschr. PIZiekt* **58**: 103-108.
- Whitehead, A G, Hemming, J R. (1965). A comparison of some quantitative methods of extracting small vermiform nematodes from soil. *Annals of Applied Biology* **55**: 25-38.